

Analysis Of Several Subtropical Cyclones By Means Of The High-Resolution HARMONIE-AROME Model

Lara Quitián Hernández¹, Daniel Santos-Muñoz², Juan Jesús González Alemán³, Javier Diaz-Fernandez⁴, Sergio Fernández-González², Pedro Bolgiani⁴, Mariano Sastre⁵, Francisco Valero⁶, and Maria Luisa Martin⁷

¹Complutense University of Madrid

²State Meteorological Agency (AEMET), Leonardo Prieto Castro, 8. 28040 Madrid. Spain.

³UCLM (University of Castilla-La Mancha), Faculty of Environmental Sciences and Biochemistry

⁴Department of Astrophysics and Atmospheric Sciences. Faculty of Physics. Complutense University of Madrid, Ciudad Universitaria s/n. 28040 Madrid. Spain.

⁵Universidad Complutense de Madrid

⁶Dpto. Astrofísica y Física de la Atmósfera. Facultad de Física. Universidad Complutense de Madrid. Ciudad Universitaria s/n. 28040 Madrid. Spain.

⁷Dpto. Matemática Aplicada. E. de Ingeniería Informática. Universidad de Valladolid. Pza. Alto de los Leones, 1. 40005 Segovia. Spain.

November 26, 2022

Abstract

Subtropical cyclones (STCs) are characterized by a hybrid structure sharing tropical and extratropical features. The impacts of this kind of cyclones are similar to the generated by tropical storms or even hurricanes, leading to widespread social damage and great economic losses. Carrying out accurate simulations becomes key for a forecast improvement of these extreme events. In this study, several STCs that hit the northeastern Atlantic coasts and experimented a transition into tropical storms (Delta 2005) or even hurricanes (Vince 2005 and Ophelia 2017) are assessed by means of simulations using the high-resolution HARMONIE-AROME model. This model is developed and operated at 2.5 km resolution through the collaboration of the 10 European National Meteorological Services (NMS) that are part of the international research program HIRLAM together with the 16 countries that comprise the ALADIN consortium. The HARMONIE-AROME model has a convection-permitting configuration and uses a non-hydrostatic spectral dynamical core with a semi-Lagrangian and semi-implicit discretization of the equations. In this way, more realistic results are obtained, which allows providing an added value to the study of tropical transitions.

Analysis of several subtropical cyclones by means of the high resolution HARMONIE-AROME model

Lara Quitián-Hernández¹, María Luisa Martín², Daniel Santos-Muñoz³, Juan Jesús González-Alemán⁴, Sergio Fernández-González³, Javier Díaz Fernández¹, Pedro Bolgiani¹, Mariano Sastre¹ and Francisco Valero Rodríguez¹

¹Dpt. of Astrophysics and Atmospheric Science. Faculty of Physics. Complutense University of Madrid, Ciudad Universitaria s/n. 28040 Madrid. Spain.

²Dpt. of Applied Mathematics. Faculty of Computer Engineering. University of Valladolid, Plza. de la Universidad, 1. 40005 Segovia. Spain.

³State Meteorological Agency (AEMET), Leonardo Prieto Castro, 8. 28040 Madrid. Spain.

⁴Enviromental Science Institute. University of Castilla-La Mancha, Avda. Carlos III s/n. 45071 Toledo. Spain.

L. Quitián-Hernández email: lquitian@ucm.es



Introduction and motivation

Subtropical Cyclones (STCs) are characterized by a hybrid structure sharing **tropical and extratropical features**^[3]. The impacts of this kind of atmospheric systems are similar to the generated by tropical storms of even hurricanes, leading to widespread social damage and great economic losses. Carrying out accurate simulations becomes key for a forecast improvement of these extreme events.

In the current work, several STCs that occurred over the North Atlantic Ocean and experimented a transition into tropical storms (Delta, 2005) or even hurricanes (Vince, 2005 and Ophelia, 2017) are assessed by means of the **high-resolution HARMONIE-AROME model**.

The HARMONIE-AROME model

This model is developed and operated at **2.5 km resolution** through the collaboration of the **10 European National Meteorological Services (NMS)** that are part of the international research program **HIRLAM**, together with the **16 countries** that comprise the **ALADIN** consortium.




Figure 1. Short Range Numerical Weather Prediction (SRNWP) Network

The **HARMONIE-AROME model** has a **convection-permitting configuration** and uses a **non-hydrostatic spectral dynamical core** with a **semi-Lagrangian** and **semi-implicit discretization of the equations**. In this way, more realistic results are obtained, which allows providing an added value to the study of tropical transitions.

This model comprises a **data assimilation system** which substantially **improves the quality of the forecasts**, since it allows to evaluate the future state of the atmosphere starting from the best possible initial state.

Global Forecast System (GFS) vs Integrated Forecast System (IFS) vs HARMONIE-AROME model

Tabla 1: FV3-GFS vs ECMWF-IFS vs HARMONIE-AROME model configuration

		Dynamical Core	Assimilation Algorithm	Resolution	Initialization Freq.
GLOBAL	FV3-GFS	Non-Hydrostatic Spectral Finite Volume (FV)	Hybrid 4D Ensemble-Var 0.875/0.125 (3-hr window)	Horizontal: 13 km Vertical levels: 64 Time step: 225 s	6 hr (cycled)
GLOBAL	ECMWF-IFS	Hydrostatic Spectral Semi-Lagrangian (SL) and Semi-Implicit (SI)	4DVar (12-hr window)	Horizontal: 9/16 km Vertical levels: 137 Time step: 450 s	12 hr (cycled)
LAM	HARMONIE-AROME	Convection-Permitting Non-Hydrostatic Spectral Semi-Lagrangian (SL) and Semi-Implicit (SI)	3DVAR (1-hr window)	Horizontal: 2.5 km Vertical levels: 65 (12m-10hPa) Time step: 75 s	3 hr (cycled)

Results

Ophelia STC stage: 07 Oct, 2017 at 1200 UTC




Figure 2. (a) HARMONIE-AROME model Total Cloud Cover (%). (b) Meteosat SEVIRI for VIS channel for hurricane Ophelia at its STC stage (1200UTC). (Source: Dundee Satellite Receiving Station).




Figure 4. (a) GFS, (b) ECMWF-IFS and (c) HARMONIE-AROME model Precipitable Water (Kg-m⁻²) simulation for hurricane Ophelia at its STC stage (1200UTC).

Delta STC stage: 22 Nov, 2005 at 1800 UTC




Figure 3. (a) HARMONIE-AROME model Total Cloud Cover (%). (b) Meteosat SEVIRI for VIS channel for hurricane Delta at its STC stage (1800UTC). (Source: Dundee Satellite Receiving Station).




Figure 5. (a) GFS, (b) ECMWF-IFS and (c) HARMONIE-AROME model Wind Speed (ms⁻¹) simulation for hurricane Delta at its STC stage (1800UTC).

Conclusions

➤ HARMONIE-AROME is the high-resolution operational Limited Area Model (LAM) of the HIRLAM and ALADIN consortia.

➤ Is a semi-Implicit (SL), semi-Lagrangian (SL), non-hydrostatic model which explicitly resolve the convection (convection-permitting).

➤ It has a 2.5 km horizontal resolution with 65 hybrid vertical levels and a model time-step of 75 s.

➤ It is stressed the good results in the prediction of convection and extreme rainfall events (included electric shocks and hail), wind speed and 2m temperature.^[1]

References

- [1] Bengtsson, L., Andrae, U., Aspelien, T., Batrak, Y., Calvo, J., deRooy, W. & Ødegard Koltzow, M. (2017). The HARMONIE-AROME model configuration in the ALADIN-HIRLAM NWP system. *Monthly Weather Review*, 145, 1919-1935 <https://doi.org/10.1175/MWR-D-16-0417.1>
- [2] ECMWF (2014). IFS Documentation, CY40R1.
- [3] Evans, J.L., & Guishard, M.P. (2009). Atlantic subtropical storms. Part I: Diagnostic criteria and composite analysis. *Monthly Weather Review*, 137, 2065-2080, doi:10.1175/2009MWR2468.1.
- [4] González-Alemán, J.J., Valero, F., Martín-León, F., & Evans, J.L. (2015). Classification and synoptic analysis of subtropical cyclones within the northeastern Atlantic Ocean. *Journal of Climate*, 28, 3331-3352. doi: 10.1175/JCLI-D-14-00276.1.
- [5] Guishard, M.P., Evans, J.L., & Hart, R.E. (2009). Atlantic subtropical storms. Part II: Climatology. *Journal of Climate*, 22, 3574-3594, doi:10.1175/2008JCLI2346.1.
- [6] Gustafsson, N., & Coauthors. (2018). Survey of data assimilation methods for convective-scale numerical weather prediction at operational centres. *Quarterly Journal of the Royal Meteorological Society*, 144, 1218-1256. <https://doi.org/10.1002/qj.3179>.
- [7] Hortal, M. (2002). The development and testing of a new two-timelevel semi-Lagrangian scheme (SETTLS) in the ECMWF forecast model. *Quarterly Journal of the Royal Meteorological Society*, 128, 1671-1687.
- [8] Laprise R. (1992). The Euler Equations of motion with hydrostatic pressure as independent variable. *Monthly Weather Review*, 120, 197-207.
- [9] Quitián-Hernández, L., Martín, M.L., González-Alemán, J.J., Santos-Muñoz, D., & Valero, F. (2016). Identification of a subtropical cyclone in the proximity of the Canary Islands and its analysis by numerical modeling. *Atmospheric Research*, 178-179, 125-137. Doi:10.1016/j.atmosres.2016.03.008
- [10] Steppeler, J., R. Hess, U. Schättler, & L. Bonaventura. (2003). Review of numerical methods for nonhydrostatic weather prediction models. *Meteorology and Atmospheric Physics*, 82, 287-301.

Acknowledgments

This work was partially supported by research projects PCIN-2014-013-C07-04, PCIN2016-080 (UE ERA-NET Plus NEWA Project), CGL2016-78702-C2-1-R and CGL2016-78702-C2-2-R, by the Instituto de Matemática Interdisciplinar (IMI) of the Universidad Complutense and by the ECMWF special projects (SPESMART and SPESVALE). The authors also thank the European Centre for Medium-Range Weather Forecasts (ECMWF) for providing the analysis databases (ECMWF).