

Crop-soil organic phosphorus cycling – a key knowledge gap for sustainable food and water resources

Victoria Janes-Bassett¹, Jessica Davies¹, Ed Rowe², Edward Tipping², and Philip Haygarth¹

¹Lancaster University

²Centre for Ecology and Hydrology

November 24, 2022

Abstract

Phosphorus is a critical nutrient in sustaining food production. In agricultural systems, application of P fertilizers has significantly increased since the green revolution to become common practice globally, contributing to increased productivity. However, excess use of P fertilizer does not only pose a cost to farmers, but costs for society in the form of water quality problems and environmental degradation. Furthermore, rock phosphates from which these fertilizers are derived are a finite resource, which brings into question the long-term sustainability of this resource and the food production it supports. Soils play a critical role in hosting the P cycle, and organic forms of P (monoesters, diesters) often represent a significant portion of soil P stocks, that are so often overlooked. The mineralization of organic P by phosphatase enzymes is recognized as a key mechanism for converting organic to inorganic forms, which can then be potentially used for P uptake by plants. However, quantification of their contribution still remains a significant challenge. In order to sustainably meet growing food production demands over the next century and reduce the impacts of P fertilizers on waterways, there is a need to understand the extent that soil organic P is available, or can be made available for plants. Here, we present recent findings from a soil-plant biogeochemical model of integrated carbon-nitrogen-phosphorus cycling in agricultural environments. Comparison of observational yield data taken from various long-term experimental sites with model simulations indicate a gap in current scientific understanding of P sources. Whilst yields in the experiments are maintained under low P addition conditions, the model indicates yield declines due to exhaustion of available P resources. We use the model to explore the missing links: potential P inputs, processes and pathways. Finally, we discuss the need for additional empirical evidence to support understanding of organic P cycling, and development of models to include these processes to inform future land management and ensure long-term food security and sustainable water resources.

Crop-soil organic phosphorus cycling

A key knowledge gap for sustainable food and water resources

Victoria Janes-Bassett¹, Jessica Davies¹, Ed C Rowe², Edward Tipping³ and Philip M Haygarth¹


1.Lancaster University, Lancaster Environment Centre, Lancaster, UK, 2.Centre for Ecology and Hydrology, Bangor, UK, 3.Centre for Ecology and Hydrology, Lancaster, UK.

B11H-2217

Phosphorus, soils and food production

Phosphorus is essential for agricultural production, global average P fertilizer use is around 1.2 g P m⁻² yr⁻¹[1]. Yet the vast majority of this P is not taken up by plants, and excess application poses issues to water quality. Organic P represents a significant portion of global P stocks; 33% representing up to 117 years of P for agricultural production[2].

Here we present findings from a plant-soil model of integrated C-N-P cycling, applied across natural and agricultural systems. Our results indicate a gap in current understanding of P cycling in agriculture, and the potential role of organic P sources in these environments.




Modelling soil P

N14CP model

We used the N14CP-Agri model[3]. This model*:

- Requires no site-specific calibration
- Uses readily available input data
- Operates over long timescales
- Simulates land use change.



The model was applied across natural sites[4], and agricultural experimental sites. Using observational data (topsoil C, N, C:N) the results demonstrate the ability of the model to capture magnitudes and variation of key variables in both natural and agricultural land uses, and across a range of management practices.

Missing phosphorus

The model works well when applied to natural (n=88) and the majority of agricultural land uses (n=52 includes control plots, N&P addition, and P only addition).


In agricultural settings with N fertilizer only (n=10) yields were underestimated by 77%.

Net Primary Productivity (NPP) is simulated assuming Liebig's law of minimum (controlled by the most limiting of temperature, precipitation, N or P availability). Prior to fertilizer addition all plots were N limited. N fertilizer increases NPP and P demand. Simulated available P declines and is then cleaved from organic pools until a maximum C:P ratio is reached, at which point P severely limits NPP and crop yields. However, observed yields do not show such a decline.


Potential sources of the missing P explored but unable to explain this deficit:

- Atmospheric deposition – up to 0.027 g m⁻² yr⁻¹ [5 – Global ave' value]
- Weathering – up to 0.085 g m⁻² yr⁻¹ [6]
- Flexible plant stoichiometries.

Extra P required to simulate observed yields for these plots ranged from 0.15-2.5 g m⁻² yr⁻¹ which is a similar magnitude to fertilizer application rates.




Applying the model: Issue with P limited plots



Knowledge gap: Organic P

Organic Phosphorus – a resource?

It's clear the model, and our current understanding of P cycling in agricultural environments is omitting a significant process and/or source of P. Our representation of plant access to organic P is basic. Whilst the importance of phosphatase enzymes as a mechanism for P assimilation in P limiting environments is recognised, quantification of their contribution is a significant challenge.



Our research therefore highlights a key gap in our understanding of organic P cycling in soils, and the sustainability of agricultural systems given the finite nature of rock sources of P fertilizer.