Spatial and temporal variability of rainfall towards watershed water resources management

Letícia Guarnier¹ and Gilberto Fonseca Barroso²

¹University Federal of Espírito Santo ²Universidade Federal do Espirito Santo

November 22, 2022

Abstract

Analysis of multi-temporal and spatial trends of rainfall in a river basin is an effective tool for water resources management based on delimited hotspots of water scarcity and flooding risks. The present study aims to characterize rainfall distribution patterns in a coastal watershed, Santa Maria da Vitória River Basin–SMVRB (Southeastern, Brazil), based on 42 meteorological stations, using geostatistical approach (i.e., kriging), from 2004 to 2017. Anisotropy effects on rainfall were observed throughout the year and were related to regional/continental climate processes. Trends in rainfall, from 1970 to 2017, were computed using the RClimDex package with eleven climate extreme indices. The results show that rainfall had an irregular distribution and droughts have become more persistent in recent years causing water scarcity to sustain crops and threatening water supply. In the lower basin, where part of the Great Vitória metropolitan area is located, flooding risks increase in the response of intensive short-term rainfall events. Knowledge of rainfall patterns contributes to assure water security and subsidize adaptative responses to extreme hydrological events.

Hosted file

tables.docx available at https://authorea.com/users/539376/articles/599898-spatial-and-temporal-variability-of-rainfall-towards-watershed-water-resources-management

Spatial and temporal variability of rainfall towards watershed water resources management

- 3 Letícia Guarnier¹ and Gilberto Fonseca Barroso²
- ¹Graduate Program of Environmental Oceanography, Federal University of Espírito
 Santo, Vitória, Espírito Santo, Brazil

²Department of Oceanography and Ecology, Federal University of Espírito Santo,
 Vitória, Espírito Santo, Brazil

8 Key points

1

2

- Rainfall spatial and temporal patterns in a coastal river basin were assessed with a geostatistical approach and climate indices.
- Extreme climate indices indicate that drought events in the upper basin threaten water
 security for agriculture and water supply.
- Because of extreme rainfall events with short-term duration, the lower basin becomes
 more prone to flooding.

15 Abstract

16 Analysis of multi-temporal and spatial trends of rainfall in a river basin is a useful 17 tool for water resources management based on delimited hotspots of water scarcity and flooding risks. The present study aims to characterize rainfall distribution in a coastal 18 19 watershed, Santa Maria da Vitória River Basin-SMVRB (Southeastern, Brazil). Data from 42 meteorological stations, from 2004 to 2017, were analyzed using kriging as a geostatistic 20 21 tool. Anisotropy effects on rainfall were observed throughout the year and were related to 22 regional/continental climate processes. From 1970 to 2017, trends in rainfall were computed 23 using the RClimDex package with eleven extreme climate indices. The results have shown 24 spatial and temporal rainfall variability, with drought events becoming more persistent in 25 recent years. Water shortage causes water scarcity to sustain crops and threatening water 26 supply. In the lower river basin, where part of the Great Vitória metropolitan area is located, flooding risks increase in the response of intensive short-term rainfall events. Knowledge of 27 28 rainfall variability in the river basin is required to assure water security, and subsidize 29 adaptative responses to extreme hydrological events.

30 **1 Introduction**

31 Climate oscillations can shift equilibrium states and cause the loss of stability of 32 ecosystems and regional landscapes. The disturbance of shifting baseline influences 33 biogeochemistry, biological productivity, biodiversity composition, and, ultimately, 34 ecosystem services. The hydrological cycle is a critical biogeochemical process of a nested 35 network of natural and social systems. Consequently, assuring water security for ecosystems, 36 drinking water, economic development, and natural hazards is essential, particularly under 37 persistent long term climate shifts further than discrete weather events. The resilience of socio-hydrological systems affected by climate oscillations at local and regional scales, and 38 39 long-term changes at a global scale, have been compromised by intensive land use and land 40 cover - LULC changes. LULC changes can amplify the frequency and intensity of climate 41 extremes, resulting in highly severe environmental and socioeconomic impacts (Pan et al., 42 2015; Zhao et al., 2018; Gogoi et al., 2019). As a result, it has been recognized to assure that 43 water security is crucial to achieving a better understanding of climate patterns and dynamics 44 (Dore, 2005; Swaney et al., 2012).

45 The assessment of water availability and hydrological risks and hazards for 46 sustainable water resources management must be based on sound information emphasizing spatial and temporal rainfall distribution in watersheds (Badr & Zaitchik, 2014; Shamir et al., 47 48 2015; Hartmann et al., 2016; Jepson et al., 2017; Vogel, 2017; Lima et al., 2018; Andreu et al., 2019; Gunda et al., 2019; Qin et al., 2019; Shikangalah & Mapani, 2019; Yang et al., 49 2019; Vera et al., 2020). Changes in rainfall patterns with extended dry periods reduce river 50 51 discharge with effects on water supply, hydroelectricity production (Gohar & Cashman, 52 2016; Su et al., 2019), and disturbances on downstream ecosystems structure and functioning

53 (Sabater et al., 2018).

River discharge reductions on lower sections of tropical and warm temperate coastal basins, under low and/or highly seasonal rainfall and fluvial discharge regulation, can lead to estuarine and groundwater salinization (Alber, 2002; Tweedley et al., 2019). Ecological effects can be related to disturbing nutrient dynamics, impairing feeding, mating and nesting habitats, and reducing the biodiversity of aquatic biological communities (Atrill & Power, 2000; Alber, 2002; Dai et al., 2006; Yin et al., 2014; Corbari et al., 2016; Costa et al., 2018; Lund-Hansen, 2018; Tweedley et al., 2019).

A concise understanding of climate dynamics requires the definition of standard
 indices for the assessment of climate data. In this respect, the World Meteorological
 Organization–WMO assembled 27 core indices of extreme climate in the RClimeDex
 software package for indices calculation and climate trends outputs (Zhang & Yang, 2004).

65 Geographic information systems (GIS) and geostatistical analysis have become powerful tools for rainfall spatial modeling and water resources management (Goovaerts, 66 1999; Mir et al., 2017; Passarella et al., 2020). GIS can handle analog, and digital 67 georeferenced data spatially indexed in a database for problem-solving using spatial analysis. 68 69 Geostatistical analysis has been considered as a practical approach for modeling climate 70 variables due to high accuracy and low errors of spatial models. Experimental variograms allow the most suitable model fit for spatial data dependence, including directional effects 71 72 (i.e., anisotropy) on regionalized variables, improving variable predictions and with low costs 73 (Holdaway, 1996; Holawe & Dutter, 1999; Noori et al., 2014; Mendez & Calvo-Valverde, 74 2016; Ozturk & Kilic, 2016).

This study aims to analyze the spatial and temporal variability and trends of rainfall data using meteorological stations, from 1970 to 2017, at a coastal river basin in southeastern Brazil, Santa Maria da Vitória River Basin–SMVRB, to promote effective water resources management of the fluvial-estuarine hydrological system.

79 **2 Materials and Methods**

80 2.1 Study area

The Santa Maria da Vitória River Basin–SMVRB drains an area of 1,799.6 km² 81 82 comprising the municipalities of Santa Maria de Jetibá, Santa Leopoldina and part of 83 Cariacica, Serra, and Vitória in Espírito Santo state, southeastern Brazil (Figure 1a). The 84 SMVRB drains to Vitória Bay estuarine system–VBES (42.9 km²), which mangrove forests cover 57.9 % (24.8 km²) of the area under a microtidal (<2.0 m) and semidiurnal regime. The 85 VBES watershed (1,921 km²), which embraces the other nine river basins, has a mean 86 estimated discharge of 65.14 m³/s (2.0×10^9 m³/yr) with SMRVB contributing with 80% of 87 88 fluvial discharge (Teubner Jr. et al., 2018).

The rugged SMVRB relief embraces elevations from sea level at Quaternary coastal plains, in the lower basin, to 1,421 m at the headwaters, where pre-Cambrian gneisses and

- 91 granitic rocks predominate. The average slope of SMVRB is 12.8%, with significant
- 92 elevation changes in the middle basin section. The regional climate is seasonal, with a hot
- 93 wet summer and dry, mild winter, with a mean annual rainfall of 1,300 mm (Teubner Jr. et
- al., 2018). Koppen climate classes are Cfb, Cfa, and Am and account to 42.3, 22.8, and 32.4
- of the basin areas at the upper, middle, and lower sections, respectively (Alvares et al., 2013).
 The Aw class accounts for 2.5% of the SMVRB area within 200 and 400 m elevation in the
- 96 The Aw class accounts for 2.5% of the SMVRB area within 200 and 400 m elevation in the97 middle basin section.
- The SMVRB has three main water reservoirs, with two of them in operation for
 hydropower generation in the main river course: Rio Bonito (2.4 km² of surface area and 22.5
 MW of hydropower capacity) with annual storage regulation; and Suiça (0.14 km² of surface
- area and 33.5 MW of hydropower capacity) with daily regulation (Teubner Jr. et al., 2018).
- 102 In 2012 the LULC of SMVRB was characterized by native Atlantic forest (45%)
- 103 followed by pastureland (16%), croplands (14%), and Eucalyptus forestry (7%). Urban areas
- account for 4% of the river basin area and are primarily concentrated in the lower basin
- section (Figure 1c). In 2010, population counting for the whole VBES was 1,159,350
- 106 inhabitants, most of them living in the urban Great Vitória Metropolitan Area at the coastal
- 107 plain. SMVRB is the primary source of water for urban supply (Teubner Jr. et al., 2018).



108

Figure 1. Study area: a) SMVRB digital elevation model; b) location of SMVRB and
meteorological stations for rainfall data in the Espírito Sato State, Southeastern Brazil; c)
2012 land use and land cover–LULC (Restinga: coastal plain herbaceous and shrub
vegetation), according to GEOBASES (2015).

113 2.2 Rainfall data

Rainfall data from 2004 to 2017 were obtained from 42 meteorological stations at 114 115 Hidroweb, a database of hydrologic information of the National Water Agency (ANA) 116 (Figure 1b). The dataset from 2018 to 2019 was not considered because of the incomplete records of several meteorological stations. When data gaps and inconsistencies were found in 117 temporal series, gaps were filled, considering a threshold of 10% of monthly data. Multiple 118 119 linear regressions considering a set of three meteorological stations, spatially nearer and 120 within the same geomorphological feature, were applied as a filling method. Regressions with $r^2 > 0.7$ were considered, while months with lower coefficients were discarded from temporal 121 122 series.

- 123 2.3 Interpolation of rainfall data and processing of continuous surface models 124 Continuous spatial distributions of rainfall data were interpolated for each month from 125 2004 to 2017. Geostatistical analysis with ordinary kriging with a spherical variogram model was applied to rainfall data using the GIS package ESRI ArcGIS 10.7 with UTM coordinate 126 system and SIRGAS 2000 datum. As a statistical requirement for monthly means, outliers 127 128 and non-normal distribution data (Naoum and Tsanis, 2004), assessed with the Shapiro-Wilk 129 test and R software, were excluded. Rainfall predictions for the SMVRB were inferred with spatial correlations of meteorological station data through experimental variograms. As a 130 131 geostatistical requirement, the spatial data structure analysis was based on the spatial 132 correlation between samples and their distances (Goovaerts, 1999).
- 133 Interpolation results were evaluated considering reports of variograms parameters, 134 cross-validation errors, linear regressions between predicted and measured values, rainfall 135 prediction maps, and maps of associated errors. Adjustments on variograms parameters are 136 required for high kriging capacity for climate datasets spatialization, including regions with
- 137 low data densities (Campling et al., 2001; Aalto et al., 2012; Choudhury et al., 2015).
- Additionally, kriging procedures can consider, in variogram analysis, data anisotropy, which is a common feature of climate data (Holdaway, 1996).
- Monthly mean rainfall models were developed with ESRI ArcGIS 10.7 raster algebra
 routines. Time series of annual rainfall for the same period, from 2004 to 2017, was
 computed considering annual precipitation. Descriptive statistics were acquired using ESRI
 ArcGIS 10.7 raster calculation routines for each continuous surface model of annual rainfall.
- 144 2.4 Cl

2.4 Climate Extreme Indices-CEI

145 The evaluation of climate change effects was computed using the software RClimDex with 11 rainfall related Climate Extreme Indices - CEI from the 27 core indices. The Expert 146 147 Team proposed the RClimDex on Climate Change Detection Monitoring and Indices -ETCCDMI (Zhang & Yang, 2004) (Table 1). After the evaluation of discontinuities and gaps, 148 a rainfall dataset of 34 meteorological stations, from 1970 to 2017, was selected for CEIs 149 150 computation. Datasets with more than 25% of annual gaps and monthly datasets with more 151 than three gaps were excluded. For each computed index, RClimDex provides as outputs 152 statistical reports, linear trends based on least squares, statistical trend significance levels, 153 according to the Fischer method, and standard error estimates (Zhang & Yang, 2004). In the present study, the confidence level for CEIs were 90 and 95%. Ordinary kriging with 154 155 spherical models was applied to interpolate CEI data using ArcGIS.

156	Tab	ole 1					
157	Selected indices of extreme climate applied in the study area						
	ID Name		Definition	Unity			
	RX1day	1 day maximum rainfall Maximum 1 day monthly rainfall					
	RX5day	5 days maximum rainfall	Maximum 5 consecutive days monthly rainfall	mm			
	SDII	Simple daily index	The ratio of annual rainfall and rainy days (≥1 mm)	mm/day			
	R10mm	Days with intense rainfall	Days with rainfall $\geq 10 \text{ mm}$	days			
	R20mm	Days with high intense rainfall	Days with rainfall $\geq 20 \text{ mm}$	days			
	R50mm	Days with rainfall higher than 50 mm	Days with rainfall \geq 50 mm	days			
	CDD	Consecutive days of droughts	Maximum consecutive days of rainfall < 1 mm	days			
	CWD	Consecutive rainfall days	Maximum consecutive days of rainfall \geq 1 mm	days			
	R95p	Very wet days	Annual total PRCP when RR>95 th percentile	mm			
	R99p	Extremely wet days	Annual total PRCP when RR>99 th percentile	mm			
	PRCPTOT	Annual total wet day precipitation	Annual total PRCP in wet days (RR≥1mm)	mm			

158

159 **3 Results and Discussion**

160 3.1 Rainfall temporal variability

161 The interpolation of monthly rainfall data, using spheric kriging, produced 168 162 continuous surface models with 30 m of spatial resolution and the associated cross-validation 163 errors. Figures 2 show the mean monthly rainfall and related RMS. Lowest RMS were 164 associated with lower data variability of dryer periods, principally after 2014. Seasonality of 165 wet months (>120 mm) comprise from late spring to early fall, and dry months (<60 mm) 166 from late fall to middle spring (Figure 2a and Table 2). December and November account for 167 35% of annual precipitation.





Figure 2. Boxplots of rainfall and estimated errors (RMS) in mm in the study area: a)
 mean monthly rainfall; b) mean RMS of monthly rainfall.

172 **Table 2**

173 Descriptive statistics of monthly rainfall (mm) for 2004 and 2017. Data from month 174 rainfall spacial models

Month	Minimum	Maximum	Mean	Standard deviation
January	3.91	336.75	150.39	112.74
February	30.66	369.16	117.61	95.87
March	32.03	399.46	179.62	115.55
April	55.03	211.19	107.81	50.90
May	0.93	205.52	63.98	57.25
June	14.70	184.99	62.81	41.65
July	3.93	149.49	52.09	40.24
August	9.91	189.03	46.46	46.28
September	23.94	115.80	51.00	24.58
October	34.40	386.32	108.65	88.94
November	65.59	425.36	219.57	95.59
December	75.97	710.63	239.33	161.38

175

176 Mean annual rainfall along the 13 years was $1,339.4 \pm 367.4$ mm. 2005 and 2013 177 were the wettest years with 2,044.3 and 1,959.5 mm, respectively. For the Considering 178 monthly averages from 2004 to 2017, the highest positive anomalies were concentrated in 179 February, May, and June of 2005. The anomalous precipitation recorded for the year 2013

180 was concentrated in December, reaching three times the monthly average.

The driest years were 2015, 2007, and 2014, with 789.1, 928.0, and 949.8 mm of
rainfall, respectively. However, the drought of 2014 and 2015 seems to be the most severe,
considering the lower rainfall variability along the 14 years.

The highest rainfalls of November and December are related to the South Atlantic 184 185 Convergence Zone (SACZ). SACZ is an important climatological feature of the austral 186 summer in South America, with high precipitation extending NW to SE, from the Amazon toward southeastern Brazil, and occurring between October to March with a period of at least 187 three days (Carvalho et al., 2004; Abrizzi & Ferraz, 2013). Conversely, the dry season, which 188 189 extends from June to August, is primarily controlled by semi-permanent meteorological 190 phenomena of South Atlantic, such as cold fronts (CF) and the South Atlantic Subtropical 191 High (SASH) (Santos et al., 2019).

The anisotropic pattern of rainfall variograms indicates an NW trend for the wettest months. In contrast, for the dry season, the tendency is SW and S (Figure 3). Rainfall in the dry season is driven by the dominant CF and SASH atmospheric systems. During this season, the SASH is displaced from the middle South Atlantic Ocean towards the continent, where lower temperatures prevent low-pressure systems, decreasing rainfall (Santos et al., 2019).

197 Anisotropy is a parameter obtained by analyzing the structure of the data in the 198 experimental variogram. It indicates a directional tendency of natural phenomena. Thus, the 199 data show the best estimates of predictions with lower associated errors in a particular 200 direction. Nevertheless, predictions of different anisotropic and isotropic variograms may not be significant (Haberlandt, 2007). Anisotropy is a common phenomenon associated to 201 202 climate variables since its relation with other factors, whether climatic as preferential to wind 203 direction, orogenic to geomorphology, geographic as the distance from the sea, among others 204 (Arbia & Lfratta, 2002; Niemi et al., 2014; Wadoux et al., 2017).

205 The analysis of the experimental variograms of the mean monthly directional trends, 206 for the period from 2004 to 2017, shows a relative homogenization of the anisotropy 207 throughout the year (Figure 3). The mean annual anisotropy of rainfall in the study area, 208 based on 168 monthly rainfall measurements, is oriented towards SW-S direction (133.7°) 209 (Figure 3m). For the driest months, between June and August, the directional trend is oriented 210 towards SW-S (Figure 3f-h). The wettest months, November and December, show 211 preferential anisotropy towards NW. From January to April, anisotropy in the BHSMV is in 212 the NE direction. From June to August, the direction follows S-SW.

The Espírito Santo central region is characterized by a marked change in elevation from NW to SE, from the hinterland towards the coast. This rouged relief influence the directional trends of rainfall anisotropy as a spatial dependent factor. A more refined understanding of rainfall anisotropy variability in the river basin would require a more spatially dense network of meteorological stations and extensive time series.



218

Figure 3. Variograms preferential directions of mean monthly (a-l) and annual
rainfall (m) in the study area for the period of 2004 to 2017 (n = 168): a) January; b)
February; c) March, d) April; e) May; f) June; g) July; h) August; i) September; j) October; k)
November; l) December; (m) annual.

223 3.2 Rainfall spatial variability

Throughout the year, the spatial distribution of rainfall in the SMVRB is geographically heterogeneous, with lower precipitation in the upper and west basin sectors during the dry season. In the wettest month, December, the amount of rainfall is higher in the west part of the river basin. During December, the rainfall is related to the SACZ (Figure 3L; Figure 4L).



229

Figure 4. SMVRB mean monthly rainfall (mm) for the period of 2004 to 2017: a)
January; b) February; c) March, d) April; e) May; f) June; g) July; h) August; i) September; j)
October; k) November; l) December.

The spatial patterns of annual rainfall are shown in Figure 5. For the years 2005 and 234 2013, the wettest years, the precipitation was concentrated in the middle basin section, where 235 the relief is more accentuated. The accumulation of rainfall in the middle section can also be 236 observed in other years. This pattern demonstrates the influence of maritime and 237 continentality on rainfall and associated regional instability lines.

In this sense, the distance from the ocean towards the continent, exert a wide range of influences on regional climate conditions. The hilly relief of the middle SMVRB, geomorphic, dominated the south Espírito Santo staggered levels. The foothills of mountain range concentrate and trap part of precipitation, producing a rain shadow with low rainfall

towards the upper basin section where the Caparaó massive dominates. Therefore, the

mountain range increases precipitation associated with frontal rains at foothills of staggering
levels. In the plateau, rainfall decreases toward the driest upper west part of the basin.



245

Figure 5. SMVRB annual rainfall (mm) for the period of 2004 to 2017.

247 3.3 Extreme Climate Indices

The analysis of time series trends is associated with its non-stationarity throughout its development to an average. Several factors that driven non-stationarity, are commonly related to factors inherent to the studied phenomenon, presenting as seasonal components, trend, or heterogeneity of variances (Milly et al., 2008; Milly et al., 2015). However, the nonstationarity of the data is a sophisticated statement that requires that the variable distribution depends on the time associated with a deterministic statistical function (Serinaldi & Kilsby, 2015; Serinaldi et al., 2018). Climate indices have been commonly applied in the analysis of climate features, mainly for rainfall variables, due to the relevance of knowing the temporal distribution of rainfall for socioeconomic and environmental issues. Figure 6 shows the annual spatial features of CEI associated with rainfall at SMVRB.



259

Figure 6. Rainfall indices at SMVRB computed with RClimDex from 1970 to 2017: 260 261 a) CDD-consecutive dry days; b) CWD-consecutive wet days; c) RX1day-maximum rainfall amount in one day; d) RX5day-maximum rainfall amount in five days; e) RX10mm-number 262 of days in a year with rainfall amount higher than 10 mm; f) RX20mm-number of days in a 263 264 year with rainfall amount higher than 20 mm; g) RX50mm-number of days in a year with 265 rainfall amount higher than 50 mm; h) R95p-extremely wet days with total annual higher than 95 percentiles; i) R99p-extremely wet days with total annual higher than 99 percentiles; 266 j) SDII-simple day intensity index for rain; k) PRCPTOT-total annual rainfall. 267

268 The PRCPTOT index indicates that rainfall for rainy days (>1 mm) shows a negative 269 trend for 71.4% meteorological stations. The PRCPTOT index shows a trend for rainfall below time-series average and drought conditions along half of the 60s, 80s, and 90s (Figure 270 271 7a). Spatial analysis of PRCPTOT presents a decrease towards W and NE, while towards E, 272 in the coastal basin section, the amount of rainfall increase (Figure 6k). Overall, there is a 273 trend for rainfall decrease in the upper basin sector, which is the contributing area with the 274 complex runoff processes for water production in the basin. Moreover, the upper basin sector shows high water demands for crop irrigation, which could impair agriculture production and 275 276 other economic activities if the negative trends of the PRCPTOT index are verified.

The analysis of consecutive dry days, computed with the CDD index, indicates a tendency of increase for a significant part of the basin, as shown for 82.3% climate stations, 37.9% of them with statistical significance. The trend of dryness is shown with the increase of the CDD index after 2014 (Figure 7b) and a sharp decrease of PRCPTOT (Figure 7a). The CDD index indicates a robust positive tendency for the lower basin section, with increasing rainfall intensity but concentrated in a few days (Figure 6a). The lower basin, particularly the coastal compartment, has been most urbanized, concentrates a significant part of the population (Teubner Jr. et al., 2018), and is vulnerable to SACZ events of high rainfall with
short duration. Urban impacts, such as flooding, landslides, and homeless inhabitants, have
been reported in the Great Vitória after SACZ torrential rainfalls (Marchioro et al., 2016).

The statistical significance associated with high positive tendencies for the CWD index, which indicates consecutive wet days, were found for 14.7% of meteorological stations. A decreasing trend was evident in the temporal series (Figure 7c). The spatial trends of CWD for the coastal sector of SMVRB are following CDD and PRCCPTOP results (Figure 6b). Therefore, the tendency for increasing precipitation in a short time interval is indicated for the highly populated coastal sector. This trend increases the risks of flooding and impacts related to rainfall drainage (Marchioro et al., 2016).

294 The R10mm index shown a negative tendency in 85,3% stations, with 43.3% of them 295 statistically significant. In general, the R10mm index follows previous indices' results with 296 the tendency of rainfall reduction through the increase of consecutive dry days and 297 concentrated rainfall in consecutive wet days in the lower section of the SMVRB. The 298 R20mm index, which has shown negative trends, while the R50mm index is showing positive 299 trends for 82.3% stations and a shift with increasing years with intensive rainfall (>50 mm) 300 (Figure 6e-g and Figure 7d-e). Monthly or annual thresholds of rainfall indices are essential 301 to identify different types of events throughout the historical series. This type of approach is commonly used in the climatological analysis to verify the frequency of days with extreme 302 303 precipitation values.

The SDII has shown positive trends for daily rainfall intensity at 76.5% stations, with 305 30.8% of them with statistical significance, as shown in Figure 7f. Following the CDD and 306 PRCPTOT indexes, SDII also shows a trend for high daily rainfall occurring in rare events in 307 the coastal sector. Maximum daily rainfall rates are also in conformity to the coastal rainfall 308 increase (Figure 6j).

The RX1day and RX5day indexes show positive trends for 76.5% climate stations, 34.6% of them with statistical significance, and 91.2% stations with 19,3% of them with statistical significance, respectively. These two indexes are following the other indexes analyzed with the trend of extended dry periods and short periods of intense rainfall, verified with more intensity in the eastern portion of SMVRB (Figure 6c-d).

For the long term, the R95p and R99p, which indicate extremely wet days, with total annual higher than 95 and 99 percentiles and positive trends for 85.3% and 76.5% climate stations, respectively, show a trend for high rainfall intensity in the short term, particularly in the coastal section. Trends of increasing very wet days and extremely wet days were evident in Figure 7g-h. The increasing rainfall on consecutive days, as indicated with the RX5day index, can lead to hydrological hazards in the coastal section.



320

Figure 7. Linear trends of rainfall indices for Cachoeira Suíça a climate station for
1961 to 2017: a) PRCPTOT; b) CDD; c) CWD; d) R10mm; e) R50mm; f) SDII; g) R95p, and
h) R99p. The continuous line is the linear regression trend, and the dashed line is the locally
weighted scatterplot smoothing.

325 The consecutive droughts of the late 80s and early 90s can be associated to intense El 326 Niño-Southern Oscillation (ENSO) events of 1987-1988, 1991-1992 and 1993-1994 (Sobral 327 et al., 2019) and related positive anomalies of the sea surface temperature (SST) in the Tropical North Atlantic (Araújo & Brito, 2011). The ENSO is associated with changes in 328 329 oceanic and atmospheric systems with the warming of Pacific Ocean SST affecting the 330 climate system with the decrease of equatorial trade winds intensity and, consequently, the 331 pattern of rainfall distribution in South America (Cane, 2005). As a cold phase of El Niño, La 332 Niña (LN) is the cooling of Equatorial Pacific Ocean SST, decreasing rainfall in Southern 333 South America. It is the extreme LN event of early 2007 and middle 2008 (Sobral et al., 334 2019).

Regardless of ENSO and LN interannual variability, the intensity and frequency of these climate oscillations have been increasing. This scenario can be aggravated with global warming of 1.5°C above the preindustrial levels as it has been predicted in several climate change models (Yeh, 2009; Cai et al., 2014; Wang et al., 2017).

Because of water scarcity scenarios, climate extreme indices based on precipitation have been applied to Brazilian arid and semiarid regions (Santos and Oliveira, 2017; Bezerra et al., 2018) and Southeastern region (Tafarello et al., 2016; Mohor & Mendiondo, 2017). Nevertheless, these indices are functional tools for analysis of positive trends of short-term intensive rainfall and, therefore, associated with hydrological hazards.

The climate extremes indices have indicated water scarcity at the SMVRB upper section, where agriculture and poultry activities are concentrated. The water shortage in the upper basin can also threaten the water regulation capacity of the Rio Bonito Reservoir (RBR), which regulates the fluvial discharge of River Santa Maria da Vitória and compromising the water supply of a part of Great Vitória metropolitan area. In September 2016, the water volume of RBR was below 30% of its storage capacity $(13.6 \times 10^6 m^3)$.

350 Thus, the understanding of rainfall characteristics and patterns can be an effective 351 way to address water security and subsidize the formulation of public policies (Guzmam et al., 2017). The prediction of a 45% increase of irrigated cropland areas in Brazil for 2030 352 353 requires the best irrigation practices, improved reservoirs management, and adequate water 354 resources grants to assure downstream water security (ANA, 2017). Water use grants issued 355 by a public agency are a crucial tool for water resources regulation in Brazil. The granting 356 system for water uses rights aims to assure river basin water security, through the analysis of 357 quali and quantitative water availability.

358 Changes in rainfall patterns can lead to a reduction of river discharge and consequently, water scarcity in river basins impairing several economic activities (Marengo 359 360 and Bernasconi, 2014; Escobar, 2015; Nobre et al., 2016). This scenario demands an evaluation of effective public policies and measures to ensure the storage of rainfall water in 361 362 the fluvial system. Water storage can be achieved through construction and operation of 363 reservoirs, and structural best management practices, such as small sediment basins (Strauch 364 et al., 2013; Saad et al., 2018), as well as adopting more integrative approaches such as 365 ecosystem-based management (Vörösmarty et al., 2018). In respect to the last issue, the payment for ecosystem services (PES) approaches recently adopted in Brazil, such as the 366 Espirito Santo State Program for Increase of Forest Cover. The Reflorestar Program aims to 367 368 restore the hydrological cycle through the restoration of forest cover in key river basin 369 headwaters for water production with landowners' payment (SEAMA, 2019).

The integration of science and governance to foster public policies is complicated since the arrangement of several issues, including available technology and social behavior, and the lack of understanding of holistic approaches for implementing sustainable policies (Liu et al., 2015; Rose & Parsons, 2015; Mercure et al., 2016; Mercure et al., 2019). Understanding spatial and temporal rainfall distribution through spatial modeling is essential for the sustainable development of river basins.

376 4 Conclusions

Kriging was a practical approach to model rainfall data, through spatial interpolation,
at SMVRB. The spatial analysis of rainfall datasets from 2004 to 2017 has shown the effect
of basin geomorphology, with orographic retention of precipitation in the steep middle basin.
Anisotropy of rainfall orientation was related to continental and regional climate processes.

381 Scenarios of water scarcity have been verified in recent years associated with climate 382 events on a global scale. Rainfall-based extreme climate indices have shown a trend for high-383 intensity rains with short duration in the coastal section of SMVRB, which is located in the 384 Great Vitória Metropolitan Area. Conversely, a tendency for rainfall reduction in the upper 385 basin section may lead to a water shortage, which can be aggravated because of damming 386 fluvial tributaries for intensive irrigation of croplands.

387 The spatial analysis and rainfall indices have shown the heterogeneity of distribution and trends of precipitation. We identify hotspots that need different approaches for water 388 389 resources management. In upper section basin strategies for mitigating the drought associated 390 with intensive agriculture is necessary. On the other hand, in the lower section, the trends for 391 flooding in urban areas and the scarce water supply in some years show that it is essential to 392 adopt adaptive measures to mitigate the socio-hydrological effects of changes on rainfall 393 patterns. It is suggested that further studies emphasizing the extremes of climatic conditions 394 must be carried out using a more extensive range of meteorological stations with statistical 395 significance above 90%. Future studies using seasonal indices can also identify other patterns 396 and trends not verified by the annual and monthly indices applied in the present work.

397 Develop the adoption of sustainability in public management of water resources
 398 should be in a participatory and decentralized scheme, involving diverse and representative
 399 stakeholders, integrating governance, science and society, presenting clear articulations
 400 between these spheres with the focus of rainfall distribution and water resources availability.

401

402 Acknowledgments

403 This study was financed in part by the Coordenação de Aperfeiçoamento de Pessoal
404 de Nível Superior–Brasil (CAPES)–Finance Code 001. The research was stored in an
405 institutional repository: http://limnolab.ufes.br/data-repository>.

406

407 **References**

- 408 Aalto, J., Pirinen, P. & Heikkinen, J. (2012). Spatial interpolation of monthly climate data for
 409 Finland: comparing the performance of kriging and generalized additive models.
 410 *Theoretical and Applied Climatology*, *112*, 99–111. https://doi.org/10.1007/s00704411 012-0716-9
- Alber, M.A. (2002). Conceptual model of estuarine freshwater inflow management.
 Estuaries, 25, 1246–1261. https://doi.org/10.1007/BF02692222
- Alvares, C.A., Stape, J.L., Sentelhas, P.C., De Moraes, G., Leonardo, J. & Sparovek, G.
 (2013). Köppen's climate classification map for Brazil. *Meteorologische Zeitschrift*, 22(6), 711-728. https://doi.org/10.1127/0941-2948/2013/0507
- Ambrizzi, T. & Ferraz, S.E.T. (2013). An objective criterion for determining the South
 Atlantic Convergence Zone. *Frontiers of Environmental Sciences*, *17*, 1-9.
 https://doi.org/10.3389/fenvs.2015.00023
- ANA–AGÊNCIA NACIONAL DAS ÁGUAS. (2017). Atlas Irrigação: uso da água na
 agricultura irrigada. Superintendência de Planejamento de Recursos Hídricos (SRH).
 Brasília–DF.

Andreu, A., Dube, T., Nieto, H., Mudau, A.E., González-Dugo, M., Guzinski, R. & Hülsmann, S. (2019). Remote sensing of water use and water stress in the African

- 425 savanna ecosystem at local scale–Development and validation of a monitoring tool.
 426 *Physics and Chemistry of the Earth, 112*, 154–164.
- 427 https://doi.org/10.1016/j.pce.2019.02.004
- Araújo, W.S & Brito, J.I.B. (2011). Indices of trends of climatic changes for the states of the
 Bahia and Sergipe by means of daily precipitacion indices and its relation with SST's
 of the Pacific and Atlantic. *Revista Brasileira de Meteorologia*, 26 (4), 541-554. .
 https://doi.org/10.1590/S0102-77862011000400004
- Arbia, G. & Lafratta, G. (2002). Anisotropic spatial sampling designs for urban pollution.
 Applied Statistics. 51(2), 223–234. https://doi.org/0035–9254/02/51223
- Atrill, M.J. & Power, M. (2000). Modelling the effect of drought on estuarine water quality.
 Water Research, 34(5),1584-1594. https://doi.org/10.1016/S0043-1354(99)00305-X
- Badr, H.S. & Zaitchik, B.F. (2014). Application of Statistical Models to the Prediction of
 Seasonal Rainfall Anomalies over the Sahel. *Journal of Applied Meteorology And Climatology*, *53*(3), 614–636. 2014. https://doi.org/10.1175/JAMC-D-13-0181.1
- Bezerra, B.G., Silva, L.L., Silva, C.M.S. & Carvalho, G.G. (2018). Changes of precipitation
 extremes indices in São Francisco River Basin, Brazil from 1947 to 2012. *Theoretical and Applied Climatology*, *135*, 565–576. https://doi.org/10.1007/s00704-018-2396-6
- Cai, W., Borlace, S., Lengaigne, M., Rensch, P., Collins, M., Vecchi, G., Timmermann, A.,
 Santoso, A., Mcphaden, M.J., Wu, L., England, M.H., Wang, G., Guilyardi, E. & Jin,
 F.-F. (2014). Increasing frequency of extreme El Niño events due to greenhouse
 warming. *Nature Climate Change*, *4*, 111–116.
 https://doi.org/10.1038/NCLIMATE2100
- 447 Campling, P., Gobin, A. & Feyen, J. (2001). Temporal and spatial rainfall analysis across a
 448 humid tropical catchment. *Hydrological Process*, *15*, 359-375.
 449 https://doi.org/10.1002/hyp.98
- Cane, M.A. (2005). The evolution of El Niño, past and future. *Earth and Planetary Science Letters*, 230, 227–240. https://doi.org/10.1016/j.epsl.2004.12.003
- 452 Carvalho, L.M.V., Jones, C. & Liebmann, B. (2004). The South Atlantic Convergence Zone:
 453 intensity, form, persistence, and relationships with iIntraseasonal to interannual
 454 activity and extreme rainfall. *Journal of Climate, 17*, 88-108.
 455 https://doi.org/10.1175/1520-0442(2004)017<0088:TSACZI>2.0.CO;2
- Choudhury, N. H., Rahman, A. & Ferdousi, S. (2015). Kriging Infill of Missing Data and
 Temporal Analysis of Rainfall in North Central Region of Bangladesh. *Journal of Climatology & Weather Forecasting*, 3(3). https://doi.org/10.4172/23322594.1000141
- 460 Corbari, C., Lassini, F. & Mancini, M. (2016). Effect of intense short rainfall events on
 461 coastal water quality parameters from remote sensing data. *Continental*462 *Shelf Research*, 123, 18–28. https://doi.org/10.1016/j.csr.2016.04.009
- 463 Costa, C.R., Costa, M.F., Dantas, D.V. & Barletta, M. (2018). Interannual and seasonal
 464 variations in estuarine water quality. *Frontiers in Marine Science*, *5*, 301
 465 https://doi.org/10.3389/fmars.2018.00301

466 Dai, M., Guo, X., Zhai, W., Yuan, L., Wang, B., Wang, L., Cai, P., Tang, T. & Cai, W. J. 467 (2006). Oxygen depletion in the upper reach of the Pearl River estuary during a winter

468 drought. Marine Chemistry, 102, 159-169. 469 https://doi.org/10.1016/j.marchem.2005.09.020 470 Dore, M.H.I. (2005). Climate change and changes in global precipitation patterns: What do 471 we know? Environment International, 31, 1167-1181. 472 https://doi.org/10.1016/j.envint.2005.03.004 473 Escobar, H. (2015). Water security. Drought triggers alarms in Brazil's biggest metropolis. 474 Science, 347(6224), 812. https://doi.org/10.1126/science.347.6224.812 GEOBASES, Mapeamento 2012-2015-Uso e cobertura do solo. Available in 475 476 https://geobases.es.gov.br/links-para-mapes1215 477 Gogoi, P. P., Vinoj, V., Swain, D, Dash, J. & Tripathy, S. (2019). Land use and land cover 478 change effect on surface temperature over Eastern India. Scientific Reports 9, 8859 479 https://doi.org/10.1038/s41598-019-45213-z 480 Gohar, A.A. & Cashman, A. (2016). A methodology to assess the impact of climate 481 variability and change on water resources, food security and economic welfare. 482 Agricultural Systems, 147, 51-64. http://dx.doi.org/10.1016/j.agsv.2016.05.008 0308-483 521X 484 Goovaerts, P. (1999). Using elevation to aid the geostatistical mapping of rainfall erosivity, 485 Catena, 34, 227-242. https://doi.org/10.1016/S0341-8162(98)00116-7. 486 Gunda, T., Hess, D., Hornberger, G. M. & Worland, S. (2019). Water security in practice: 487 The quantity-quality-society nexus. Water Security, 6:100022. 488 https://doi.org/10.1016/j.wasec.2018.100022 489 Guzmán, D.A., Mohor, G.S., Taffarello, D. & Mendiondo, E.M. (2017). Economic impacts of 490 drought risks for water utilities through Severity-Duration-Frequency framework 491 under climate change scenarios. Hydrology and Earth System Sciences Discuss., 492 https://doi.org/10.5194/hess-2017-615 493 Haberlandt, U. (2007). Geostatistical interpolation of hourly precipitation from rain gauges 494 and radar for a large-scale extreme rainfall event. Journal of Hydrology, 332, 144-495 157. https://doi.org/10.1016/j.jhydrol.2006.06.028 496 Hartmann, H., Snow, J., Su, B. & Jiang, T. (2016). Seasonal predictions of precipitation in 497 the Aksu-Tarim River basin for improved water resources management. Global and 498 Planetary Change, 147, 86–96. https://doi.org/10.1016/j.gloplacha.2016.10.018 499 Holawe, F. & Dutter, R. (1999). Geostatistical study of precipitation series in Austria: time 500 and space. Journal of Hydrology, 219, 70-82. https://doi.org/10.1590/0001-501 3765201620150103 502 Holdaway, M.R. (1996). Spatial modeling and interpolation of monthly temperature using 503 kriging. Climate Research, 6, 215-225. https://www.jstor.org/stable/24864559 504 Jepson, W., Budds, J., Eichelberger, L., Harris, L., Norman, E., O'reilly, K., Pearson, A., Shah, S., Shinn, J., Staddon, C., Stoler, J., Wutich, A. & Young, S. (2017). Advancing 505 506 human capabilities for water security: A relational approach. Water Security, 1, 46-507 52.. https://doi.org/10.1016/j.wasec.2017.07.001 508 Lima G. N., Lombardo, M. A. & Magaña, V. (2018). Urban water supply and the changes in 509 the precipitation patterns in the metropolitan area of São Paulo-Brazil. Applied 510 Geography, 9, 223–229. https://doi.org/10.1016/j.apgeog.2018.03.010

- Liu, J., Mooney, H., Hull, V., Davis, S. J., Gaskell, J., Hertel, T., Lubechenco, J., Seto, K. C.,
 Gleick, P., Kremen, C. & Li, S. (2015). Systems integration for global sustainability. *Science*, 347(6225), 1.258.832. https://doi.org/10.1126/science.1258832
- Lund-Hansen, L.C., Jensen, K.T., Andersen, T.J., Nielsen, M.H., Doan-Nhu, H. & NguyenNgoc, L. (2018). Impacts and effects of a historical high and ENSO linked freshwater
 inflow in the tropical estuary Nha Phu, southeast Vietnam. *Regional Studies in Marine Science*, 17, 28–37. https://doi.org/10.1016/j.rsma.2017.11.012
- Marchioro, E., Silva, G.M. & Correa, W.S.C. (2016). A zona de convergência do Atlântico
 sul e a precipitação pluvial do município de Vila Velha (ES): repercussões sobre as
 inundações. *Revista do Departamento de Geografia, 31*, 101-117.
 http://dx.doi.org/10.11606/rdg.v31i0.108447
- Marengo, J.A. & Bernasconi, M. (2014). Regional differences in aridity/drought conditions
 over Northeast Brazil: present state and future projections. *Climatic Change*, 129,
 103-115. http://dx.doi.org/10.1007/s10584-014-1310-1
- Mendez, M. & Calvo-Valverde, L. (2016). Assessing the performance of several rainfall
 interpolation methods as evaluated by a conceptual hydrological model. *Procedia Engineering*, 154, 1050-1057. https://doi.org/10.1016/j.proeng.2016.07.595
- Mercure, J.-F., Pollitt, H., Bassi, A.M., Viñuales, J. E. & Edwards, N.R. (2016). Modelling
 complex systems of heterogeneous agents to better design sustainability transitions
 policy. *Global Environmental Change*, *37*, 102–115.
 http://dx.doi.org/10.1016/j.gloenvcha.2016.02.003
- Mercure, J.-F., Paim, M.A., Bocquillon, P., Lindner, S., Salas, P., Martinelli, P., Berchin, I.I.,
 Andrade Guerra, J.B.S.O., Derani, C., Albuquerque Junior, C.L., Ribeiro, J.M.P.,
 Knobloch, F., Pollitt, H., Edwards, N.R., Holden, P.B., Foley, A., Schaphoff, S.,
 Faraco, R.A. & Vinuales, J.E. (2019). System complexity and policy integration
 challenges: The Brazilian Energy-Water-Food Nexus. *Renewable and Sustainable Energy Reviews*, 105, 230–243. https://doi.org/10.1016/j.rser.2019.01.045
- Milly, P.C D., Betancourt, J., Falkenmark, M., Hirsch, R.M., Kundzewics, Z.W.,
 Lettenmaier, D. P. & Stouffer, R. (2008). Stationarity is dead: Whither water
 management? *Climate Change*, *139*, 573–574. https://doi.org/10.1126
 /science.1151915
- Milly, P.D.C., Betancourt, J., Falkenmark, M., Hirsch, R.M., Kundzewics, Z.W.,
 Lettenmaier, D. P., Stouffer, R. Dettinger, M.D. & Krysanova, V. (2015). On
 Critiques of "Stationarity is dead: whither water management?" *Water Resources Research*, *51*(9), 7785-7789. https://doi.org/10.1002/2015WR017408
- 546 Mir, A., Piri, J. & Kisi, O. (2017). Spatial monitoring and zoning water quality of Sistan
 547 River in the wet and dry years using GIS and geostatistics. *Computers and Electronics*548 *in Agriculture, 135*, 38–50. http://dx.doi.org/10.1016/j.compag.2017.01.022
- Mohor, G.S. & Mendiondo, E.M. (2017). Economic indicators of hydrologic drought
 insurance under water demand and climate change scenarios in a Brazilian context.
 Ecological Economics, 140, 66–78. https://doi.org/10.1016/j.ecolecon.2017.04.014
- Naoum, S. & Tsanis, I.K. (2004). Ranking spatial interpolation techniques using a GIS-based
 DSS. *Global Nest Jornal*, 6(1), 1-20. https://doi.org/10.30955/gnj.000224

554 Niemi, T.J., Kokkonem, T. & Seed, A.W. (2014). A simple and effective method for quantifying spatial anisotropy of time series of precipitation fields. Water Resources 555 Research, 50 (7), 5906-5925. https://doi.org/10.1002/2013WR015190 556 557 Nobre, C.A., Marengo, J.A., Seluchi, M.E., Cuartas, L.A. & Alves, L.M. (2016). Some 558 characteristics and impacts of the drought and water crisis in Southeastern Brazil 559 during 2014 and 2015. Journal of Water Resource and Protection, 8(2), 252-262. 560 https://doi.org/10.4236/jwarp.2016.82022 561 Noori, M.J., Hassan, H.H. & Mustafa, Y.T. (2014). spatial estimation of rainfall distribution and its classification in Duhok Governorate using GIS. Journal of Water Resource 562 and Protection, 6, 75-82. http://dxdoi.org/10.4236/jwarp.2014.62012 563 Ozturk, D. & Kilic, F. (2016). Geostatistical approach for spatial interpolation of 564 565 meteorological data. Anais da Academia Brasileira de Ciências 88(4), 2121-2136. 566 http://dx.doi.org/10.1590/0001-3765201620150103 567 Pan T., Wu S. & Liu, Y. (2015). Relative Contributions of Land Use and Climate Change to 568 Water Supply Variations over Yellow River Source Area in Tibetan Plateau during 569 the Past Three Decades. PLoS ONE, 10(4), e0123793. http://dx.doi.org/10.1371/journal.pone.0123793 570 571 Passarella, G., Bruno, D., Lay-Ekuakille, A., Maggi S., Masciale, R. & Zaccaria, D. (2020). 572 Spatial and temporal classification of coastal regions using bioclimatic indices in a 573 Mediterranean environment. Science of the Total Environment, 700, 134415. 574 https://doi.org/10.1016/j.scitotenv.2019.134415 575 Qin, K., Liu, J., Yan, L. & Huang, H. (2019). Integrating ecosystem services flows into water 576 security simulations in water scarce areas: Present and future. Science of the Total 577 Environment 670, 1037-1048. https://doi.org/10.1016/j.scitotenv.2019.03.263 0048-578 9697 579 Rose, N. A. & Parsons, E. C. M. (2015). "Back off, man, I'm a scientist!": When marine 580 conservation science meets policy. Ocean & Coastal Management, 115, 71-76. 581 https://doi.org/10.1016/j.ocecoaman.2015.04.016 582 Saad, S.I., Mota da Silva, J., Silva, M.L.N., Guimarães, J.L.B., Sousa Júnior, W.C., 583 Figueiredo, R.D.O. & Rocha, H.R. (2018). Analyzing ecological restoration strategies 584 for water and soil conservation. PLoS ONE, 13(2), e0192325. 585 https://doi.org/10.1371/journal.pone.0192325 586 Sabater, S., Bregoli, F., Acuña, V., Barceló, D., Elosegi, A., Ginebreda, A., Marcé, R., 587 Muñoz, I., Sabater-Liesa, L. & Ferreira, V. (2018). Effects of human-driven water 588 stress on river ecosystems: a meta-analysis. Scientific Reports, 8, 11462. 589 https://doi.org/10.1038/s41598-018-29807-7 590 Santos, V., Gastmans, D., Sánchez-Murillo, R., Gozzo, L. F., Batista, L. V., Manzione, R.L. 591 & Martinez, J. (2019). Regional atmospheric dynamics govern interannual and 592 seasonal stable isotope composition in southeastern Brazil. Journal of Hydrology, 593 579, 124136. https://doi.org/10.1016/j.jhydrol.2019.124136 594 Santos, C.A.C. & Oliveira, V.G. (2017). Trends in extreme climate indices for Pará State, 595 Brazil. Revista Brasileira de Meteorologia, 32(1), 13-24. 596 http://dx.doi.org/10.1590/0102-778632120150053

597 SEAMA-Secretaria de Estado do Meio Ambiente e Recursos Hídricos. Diário Oficial dos Poderes do Estado: Portaria Nº 005-R, de 22 de março de 2019. Available in 598 599 https://seama.es.gov.br/o_que_e_reflorestar Access in Jan 2020. Serinaldi, F. & Kilsby, C.G. (2015). Stationarity is undead: uncertainty dominates the 600 601 distribution of extremes. Advances in Water Resources, 77, 17-36. 602 http://dx.doi.org/10.1016/j.advwatres.2014.12.013 603 Serinaldi, F., Kilsby, C.G. & Lombardo, F. (2018). Untenable nonstationarity: an assessment 604 of the fitness for purpose of trend tests in hydrology. Advances in Water Resources, 605 111, 132-155. http://dx.doi.org/10.1016/j.advwatres.2017.10.015 606 Shamir, E., Megdal, S.B., Carillo, C., Castro, C.L., Chang, H.I., Chief, K., Corkhill, F. E., 607 Eden, S., Georgakakos, K. P., Nelson, K. M. & Prietto, J. (2015). Climate change and water resources management in the Upper Santa Cruz River, Arizona. Journal of 608 609 Hydrology, 521, 18-33. http://dx.doi.org/10.1016/j.jhydrol.2014.11.062 Shikangalah, R.N. & Mapani, B. (2019). Precipitation variations and shifts over time: 610 611 Implication on Windhoek city water supply. Physics and Chemistry of the Earth, 112, 612 103-112. https://doi.org/10.1016/j.pce.2019.03.005 613 Sobral, B. S., Oliveira-Júnior, J. F., Gois, G., Pereira-Júnior, E. R., Terassi, P. M. B., Muniz-614 Júnior, G. R., Lyra, G. B. & Zeri, M. (2019). Drought characterization for the state of Rio de Janeiro based on the annual SPI index: trends, statistical tests and its relation 615 616 with ENSO. Atmospheric Research, 220, 141-154. 617 https://doi.org/10.1016/j.atmosres.2019.01.003 618 Strauch, M., Lima, J. E. F. W., Volk, M., Lorz, C. & Makeschin, F. (2013). The impact of 619 Best Management Practices on simulated streamflow and sediment load in a Central Brazilian catchment. Journal of Environmental Management, 127, S24-S36. 620 621 http://dx.doi.org/10.1016/j.jenvman.2013.01.014 622 Su, Y., Gao, W. & Guan, D. (2019). Integrated assessment and scenarios simulation of water 623 security system in Japan. Science of the Total Environment, 671, 1269–1281. https://doi.org/10.1016/j.scitotenv.2019.03.373 0048-9697 624 625 Swaney, D. P., Humborg, C., Emeis, K., Kannen, A., Silvert, W., Tett, P., Pastres, R., Solidoro, C., Yamamuro, M., Hénocque, Y. & Nicholls, R. (2012). Five sale 626 questions of scale for the coastal zone. Estuarine, Coastal and Shelf Science, 96, 9-21. 627 https://doi.org/10.1016/j.ecss.2011.04.010 628 629 Taffarello, D., Mohor, G.S., Calijuri, M.C. & Mendiondo, E.M. (2016). Field investigations 630 of the 2013–14 drought through quali-quantitative freshwater monitoring at the 631 headwaters of the Cantareira System, Brazil. Water International, 41(5), 776-800. 632 http://dx.doi.org/10.1080/02508060.2016.1188352 Teubner Jr., F. J., Lima, A. T. M. & Barroso, G. F. (2018). Emission rates of nitrogen and 633 634 phosphorus in a tropical coastal river basin: a strategic management approach. Environmental Monitoring and Assessment, 190, 747 https://doi.org/10.1007/s10661-635 018-7101-9 636 637 Tweedley, J. R., Dittmann, S. R., Whitfield, A. K., Withers, K., Hoeksema, S. D. & Potter, I. 638 C. (2019). *Hypersalinity: Global distribution, causes, and present and future effects* 639 on the biota of estuaries and lagoons. Wolanski, E., Day, J. W., Elliott, M. & 640 Ramachandran, R. Coasts and estuaries: the future, Elsevier, 523-546. 641 https://doi.org/10.1016/B978-0-12-814003-1.00030-7

- Vera, J F.R., Mera, Y.E.Z. & Pérez-Martín, M.A. (2020). Adapting water resources systems
 to climate change in tropical areas: Ecuadorian coast. *Science of the Total Environment.* 703, 135554. https://doi.org/10.1016/j.scitotenv.2019.135554
- Vogel, R. M. (2017). Stochastic watershed models for hydrologic risk management. *Water Security*, 1, 28-35. 2017. https://doi.org/10.1016/j.wasec.2017.06.001.
- Vörösmarty, C. J., Osuna, V. R., Cak, A. D., Bhaduri, A., Bunn, S. E., Corsi, F.,
 Gastelumendi, J., Green, P., Harrison, I., Lawford, R., Marcotullio, P. J., McClain,
 M., McDonald, R., McIntyre, P., Palmero, M., Robarts, R. D., Szöllösi-Nagy, A.,
 Tessler, Z. & Uhlenbrook, S. (2018). Ecosystem-based water security and the
 Sustainable Development Goals (SDGs). *Ecohydrology & Hydrobiology*, *18*(4), 317333. https://doi.org/10.1016/j.ecohyd.2018.07.004
- Wadoux, A.M.J-C, Brus, DJ., Rico-Ramirez, M.A. & Heuvelink, G.B.M. (2017). Sampling
 design optimization for rainfall prediction using a non-stationary geostatistical model. *Advances in Water Resources, 107*, 126–138.
 https://doi.org/10.1016/j.advwatres.2017.06.005
- Wang, G., Gan, B., Cai, W. & Wu. L. (2017). Continued increase of extreme El Niño
 frequency long after 1.5°C warming stabilization. *Natural Climate Change*, 7, 568573. https://doi.org/10.1038/NCLIMATE3351
- Yang, H., Xiao, H., Guo, C. & Sun, Y. (2019). Spatial-temporal analysis of precipitation
 variability in Qinghai Province, China. *Atmospheric Research*, 228, 242–260.
 https://doi.org/10.1016/j.atmosres.2019.06.005
- Yeh, S.W, Kug, J. S. Dewite, B., Kwon, M. H., Kirtman, B.P. & Jin, F.F. (2009). El Niño in a changing climate. *Nature 461*, 511-515. https://doi.org/10.1038/nature08316
- Yin, K., Lin, Z. & Ke, Z. (2004). Temporal and spatial distribution of dissolved oxygen in the
 Pearl River Estuary and adjacent coastal waters. *Continental Shelf Rearch*, 24, 19351948. https://doi.org/10.1016/j.csr.2004.06.017
- Zhang, X. & Yang, F. (2004). *RClimDex (1.0) user manual*. Environment Canada, 23pp.
 [available at: http://etccdi.pacificclimate.org/RClimDex/RClimDexUserManual.doc]
- Kata Z., Liu, G., Liu, Q., Huang, C. & Li, H. (2018). Studies on the spatiotemporal
 variability of river water quality and its relationships with soil and precipitation: a
 case study of the Mun river basin in Thailand. *International Journal of Environmental Research and Public Health*, 15, 2466. https://doi.org/10.3390/ijerph15112466

Figure 1.



Figure 2.



Figure 3.



Figure 4.



Figure 5.



Figure 6.



Figure 7.

