

Significant Improvement of Cloud Representation in MRI-ESM2

Hideaki Kawai¹, Seiji Yukimoto¹, Tsuyoshi Koshiro¹, Naga Oshima², Taichu Tanaka³,
Hiromasa Yoshimura¹, and Ryoji Nagasawa¹

¹Meteorological Research Institute

²Meteorological Research Institute, Japan Meteorological Agency

³Japan Meteorological Agency

November 23, 2022

Abstract

The representations of clouds in climate model MRI-ESM2 (Yukimoto et al. 2019, JMSJ) used in CMIP6 simulations are significantly improved from the previous version MRI-CGCM3 used in CMIP5 simulations. The score of the spatial pattern of 2m temperature for MRI-ESM2 is better than any of the 47 CMIP6 models. The score for radiative fluxes at the top of the atmosphere for MRI-ESM2 is ranked in the top seven models among these CMIP6 models. We will introduce comprehensively various modifications related to clouds, which contribute to the dramatically improved cloud representation, and their main impacts (Kawai et al. 2019, GMD). The modifications cover various schemes and processes including the cloud scheme, turbulence scheme, cloud microphysics processes, the interaction between cloud and convection schemes, resolution issues, cloud radiation processes, aerosol properties, and numerics. In addition, we would like to mention what data we used during the model development, how we evaluated our model, how useful the intercomparison studies were, and what data we need for further model development.

Significant Improvement of Cloud Representation in MRI-ESM2

Hideaki Kawai, Seiji Yukimoto, Tsuyoshi Koshiro, Naga Oshima, Taichu Tanaka, Hiromasa Yoshimura, and Ryoji Nagasawa

Meteorological Research Institute, JMA




Kawai, H., et al., 2019: Significant Improvement of Cloud Representation in Global Climate Model MRI-ESM2. *Geosci. Model Dev.*, **12**, 2875-2897.

Improvement in clouds in MRI-ESM2

Old model: MRI-CGCM3

New model: MRI-ESM2




TCC bias



* much less over the Southern Ocean

* Much Improved over the SO and off Peru

TOA SW bias
(upward)




* Too much SW reflection over Tropics

* Improved over Tropics

* Much less SW reflection over the SO

* Improved over the SO and off Peru

Error in Net Radiation (SW+LW, TOA)



New model MRI-ESM2: Best among CMIP5 models

7th among 47 CMIP6 models (5th for 2m Temperature)


(SW is also the best among CMIP5 models. Taylor diagrams show similar results.)

Various improvements related to clouds...

Kawai et al. (2019, *GMD*)

- i. Stratocumulus parameterization (turbulence scheme)
- ii. Cloud microphysics
- iii. Vertical resolution
- iv. Convection scheme (shallow convection)
- v. Cloud overlap scheme for radiation
- vi. Radiation process
- vii. Bug
- viii. Aerosol mode radii
- ix. Cloud ice fall calculation


Impact of Stratocumulus parameterization



Not only temperature but also water vapor profile is taken into account.

$$ECTEI = EIS - \beta \frac{L}{c_p} (q_{\text{surf}} - q_{700})$$

Highest correlation!



Cloud Obs. Data:
Meteo. Field Data:


EECRA Ship Obs. Climatology
ERA40
Using Global Ocean Data

Impact of introduction of new stratocumulus scheme

Cloud top mixing is suppressed for $ECTEI > -2$ K.
($ECTEI$ -LCC relationship is not used explicitly.)

New stratocumulus scheme — Old scheme


Low Cloud Cover



LCC is increased over the SO, the North Pac., and west coast of the continents

(Closer to Obs.)

TOA SW radiation (upward)




SW reflection is increased over these regions

(Closer to obs.)

10-yr mean

Impact of modified cloud microphysics

Liquid Water Ratio



Observation: High liquid water ratio


Modified treatment of WBF effect in the model cloud microphysics:

Supercooled water was increased. ➡ Optical thickness of clouds increased.
(Cloud droplets are smaller than ice crystals.)

Impact of modified treatment of WBF effect on SW radiation

TOA SW radiation (upward)

New WBF effect — Old WBF effect



SW reflection is increased over the Southern Ocean
(Closer to obs.)

10-yr mean


Suppression of Shallow Convection

Increased Vertical Resolution

Cross sections of cloud fraction along 20°S


Shallow Cu conditionally off

L80




Shallow Cu on

L80



Shallow Cu on

L48



Low-level cloud transition from stratocumulus to cumulus became more realistic when shallow convection is suppressed over the area where stratocumulus forms.

Geometrically thin boundary layer clouds became more realistic in L80.


10-yr mean

Impact of new cloud overlap scheme

closer to Maximum-random ovlp

closer to random ovlp

New cloud overlap scheme — Old scheme



Too much reflection is reduced over the tropics.

(Closer to Obs.)


New scheme: PICA (Nagasawa 2012)

10-yr mean

Impact of considering fine maritime aerosols


CCN from maritime aerosols is doubled to consider fine maritime aerosols

Column cloud droplet number concentration (ratio)



N_c is increased over the Southern Ocean

TOA SW radiation (upward)



SW reflection is increased over the SO

(Closer to obs.)

Lower bound of N_c that affects a lot 20C radiation forcing by aerosols (Hoose et al. 2009) is not used in MRI-ESM2.

10-yr mean

Various improvements related to clouds...

Kawai et al. (2019, *GMD*)

- i. Introduction of a new **stratocumulus parameterization** based on CTE (cloud top entrainment) criterion (Kawai 2013, Kawai et al. 2017)
Cloud shortage over the Southern Ocean & Northern Pacific was alleviated.
- ii. The modification of the treatment of the Wegener–Bergeron–Findeisen process in **cloud microphysics**, etc.
Supercooled water was increased. Then, cloud optical thickness also increased.
- iii. Increased **vertical resolution** from L48 to L80 (Especially in BL.)
Geometrically thin boundary layer clouds became more realistic.
- iv. Suppression of **shallow convection** under condition of stratocumulus occurrence
Low-level cloud transition from stratocumulus to cumulus became more realistic.
- v. Improvement of a **cloud overlap scheme** (introduction of PICA; Nagasawa 2012)
An excess reflection of shortwave radiation over the tropics was drastically alleviated.

Various improvements related to clouds...

Kawai et al. (2019, *GMD*)


- vi. Abolishment of spatially reduced calculation of a **radiation process**
The low-level clouds in the subtropics and mid-latitudes slightly increased.
- vii. A **bug** associated with the prognostic equations of number concentrations of cloud particles was fixed.
Too large number concentrations of cloud particles, particularly, for Sc and St were dissolved.
- viii. Modification of **aerosol mode radii** based on recent observations
Number concentrations of cloud particles became more appropriate.
- ix. Improved calculation of **cloud ice fall** (based on Kawai 2005)
The calculation became more realistic & the time-step dependency of IWC was alleviated.

Others * Improvement in the aerosol model (MASINGAR) etc.


Tuning for total performance (in a convention scheme etc...)

Contribution from various modifications (based on on-off exp.)

Contribution to LCC




Contribution to TOA SW radiation



- Stratocumulus scheme
- Cloud microphysics (WBF effect)
- Fine maritime aerosols
- Radiation spatial resolution
- Cloud overlap
- Shallow Conv. off over Sc condition

Kawai et al. (2019, *GMD*)

TOA SW radiation bias (upward)



- ❑ The representations of clouds in climate model MRI-ESM2 (for CMIP6) are significantly improved from the previous version MRI-CGCM3 (for CMIP5).
- ❑ The significant improvement is not attributed to the introduction of a new advanced scheme but to the cumulative effect of many "minor" modifications.

Acknowledgements

These improvements were achieved owing to the valuable information obtained from multiple model intercomparison studies and projects.

This work was partly supported by the Integrated Research Program for Advancing Climate Models (TOUGOU) Grant Number JPMXD0717935561 from the Ministry of Education, Culture, Sports, Science and Technology (MEXT), Japan. It was also supported by the Japan Society for the Promotion of Science (JSPS) KAKENHI Grant Numbers JP18H03363, JP19K03977, and JP19H05699.

References

- Hoose, C., et al., 2009: Constraining cloud droplet number concentration in GCMs suppresses the aerosol indirect effect, *Geophys. Res. Lett.*, **36**, L12807.
- Kawai, H., 2005: Improvement of a Cloud Ice Fall Scheme in GCM. *CAS/JSC WGNE Research Activities in Atmospheric and Oceanic Modelling/WMO*, **35**, 4.11-4.12.
- Kawai, H., 2013: Improvement of a Stratocumulus Scheme for Mid-latitude Marine Low Clouds. *CAS/JSC WGNE Research Activities in Atmospheric and Oceanic Modelling/WMO*, **43**, 4.03-4.04.
- Kawai, H., S. Yukimoto, T. Koshiro, N. Oshima, T. Tanaka, H. Yoshimura, and R. Nagasawa, 2019: Significant Improvement of Cloud Representation in Global Climate Model MRI-ESM2. *Geosci. Model Dev.*, **12**, 2875-2897.
- Kawai, H., T. Koshiro, and M. J. Webb, 2017: Interpretation of Factors Controlling Low Cloud Cover and Low Cloud Feedback Using a Unified Predictive Index. *J. Climate*, **30**, 9119-9131.
- Nagasawa, R., 2012: The Problem of Cloud Overlap in the Radiation Process of JMA's Global NWP Model. *CAS/JSC WGNE Research Activities in Atmospheric and Oceanic Modelling/WMO*, **42**, 4.15-4.16.
- Yukimoto, S., et al., 2019: The Meteorological Research Institute Earth System Model version 2.0, MRI-ESM2.0: Description and basic evaluation of the physical component. *J. Meteor. Soc. Japan*, **97**, 931-965.