

Options for increased soil carbon storage and water holding capacity through sustainable agriculture and forestry: modelling results from showcase regions in Austria

Dagmar Henner¹, Gottfried Kirchengast², Melannie Hartman³, and Clara Hohmann²

¹Wegener Center for Climate and Global Change, University of Graz, Austria

²University of Graz

³Natural Resource Ecology Laboratory, Colorado State University, USA

November 22, 2022

Abstract

Sustainable agriculture and forestry is an important topic under climate change. This is a potential route for increasing long-term soil and biomass carbon storage, soil water retention capacity and for reducing water and wind erosion risks. This study uses the Styrian Raab and Enns catchment regions in Southeastern Austria as showcase regions for exploring sustainable whole-system options for climate change adaptation and mitigation under increased hot-dry conditions in agriculture and forestry. Based on dense data of the WegenerNet observing network and further hydrometeorological data, combined with hydrological modelling (WaSiM), the current hydrological disturbance potential in the Southeastern Austria focus regions is assessed. Furthermore, downscaled IPCC climate change scenarios are used for future projections and the results are evaluated for increasing heat and drought risks. This work provides the hydrological context for modelling the soil water and carbon storage enhancement options that farming, forestry and land-use practices might apply. A first key study aspect in this context is the sustainable potential of bioenergy crops. Using the local-scale WegenerNet data combined with Harmonized World Soil Database (HWSD) soil data, potential yields for bioenergy from lignocellulosic biomass (forest and Miscanthus, willow and poplar) is modelled using MiscanFor, SalixFor, PopFor and DAYCENT models for representative local areas in the showcase regions. Using DAYCENT biogeochemical modelling with different agricultural, forest management, and land use practices under climate change, sustainable system options under different future climate change scenarios are developed. These results will be used in turn to develop whole-system options, namely to jointly achieve increase of soil carbon and robustness of soil water retention capacity, increase of soil quality, reduction of soil erosion and degradation, reduced compaction, stabilisation of slopes, sustainability and resilience in the soil as well as the agricultural and forest production system. Sustainable whole-system options for farmers and forest managers are the major study focus and will supply data for local estimates in the chosen representative areas in the showcase regions as well as for upscaling to country level.

Options for increased soil carbon storage and water holding capacity through sustainable agriculture and forestry: modelling results from a showcase region in Austria



Options for increased soil carbon storage and water holding capacity through sustainable agriculture and forestry: modelling results from a showcase region in Austria

Authors: Dagmar Nadja Henner (1,2), Gottfried Kirchengast (1, 2, 3), Melannie D. Hartman (4), Clara Hohmann (1,2)

(1) Wegener Center for Climate and Global Change, University of Graz, Austria, (2) FWF-DK Climate Change, University of Graz, Austria, (3) Institute for Geophysics, Astrophysics, and Meteorology/Institute of Physics, University of Graz, Austria, (4) Natural Resource Ecology Laboratory, Colorado State University, USA



Welcome, Poster Presentation and Background

Welcome

AGU Fall Meeting 2020 poste...

OPEN

Location and Data

The Styrian Raab catchment in Southeastern Austria was used as a showcase region for exploring sustainable whole-system options for climate change adaptation and mitigation under increased hot-dry conditions.

Information about **Styria**




Fig. 1: Upper panel: The red cross shows the location of Hofstätten an der Raab. The map of the Raabtal region (RTR) located in Southeastern Austria (see Inset left inset) includes the Styrian Raab catchment (red outline) with major rivers (blue lines) and cities (black dots).

OPEN

The Sustainability Potential of Bioenergy Crops

Potential yields (as primary use) for bioenergy from lignocellulosic biomass (*Miscanthus*, willow and poplar) are modelled using **MiscanFor**, **SalixFor**, **PopFor** and **DayCent** models for Hofstätten bioenergy site. The synergistic potential of bioenergy crops, especially the increase of soil carbon, is shown in the following figures. This so-called "secondary use" is the primary focus of this research. For comparison with the potential of *Miscanthus*, willow and poplar, the usual agricultural rotation of maize and soybean is modelled with DayCent at the same site.




Fig. 3: Aboveground live and dead biomass carbon stored in *Miscanthus* (misc), willow and poplar at Hofstätten bioenergy site modelled with DayCent using the fert1 scenario. The bioenergy crops were planted in 2007 and the graph shows the development until the end of 2019. The different harvest schedule for *Miscanthus* (yearly) and willow or poplar (every fourth year) is obvious, and biomass stored in live and dead plant parts follows this schedule.

OPEN

Discussion and Conclusions

This research has shown that there is a large potential for soil carbon increase by lignocellulosic bioenergy crops compared with conventional agricultural rotation. Figure 6 shows the development of soil organic matter carbon under bioenergy crops and conventional agricultural rotation. While soil organic matter under agriculture slowly decreases over time, there is a significant increase under the bioenergy crops in this research. This is mostly caused by less frequent harvest, decreased tillage and by the growth pattern of bioenergy crops themselves (e.g. extensive root system). Especially *Miscanthus* has a high potential for stabilizing soils because it is normally harvested while dormant, which stabilizes soil carbon but also the carbon stored in plant material.

Methods and References

Methods

DayCent was set up using the local-scale WegenerNet data combined with measured soil data (pH, soil structure) from the Austrian Chamber of Agriculture. All DayCent simulations (fert1 scenario) started with the same initial conditions on January 1, 2007, and ended on December 31, 2019. The agricultural system before was run as a continuous maize system because maize is the dominant agricultural crop in the research area. For *Miscanthus*, no fertilizer was added, and we assumed that the crop was harvested every year at day 350. For poplar and willow, 5.6 gN/m² fertilizer were added each year and harvest was scheduled every fourth year. For the maize and soybean rotation, in the year 2007 maize was planted in spring and 2.8 gN/m² and later in the year again 3.04 gN/m² fertilizer were added.

OPEN

Outlook and Contact Information

Outlook

The DayCent model will allow computing a GHG budget for all land use options and potential new systems of land use. These results will be used in turn to develop whole-system options, namely to jointly achieve an increase of soil carbon and robustness of soil water retention capacity, an increase of soil quality, reduction of soil erosion and degradation, reduced compaction, stabilization of slopes, sustainability and resilience in the soil as well as the agricultural and forest

OPEN

Authors: Dagmar Nadja Henner (1,2), Gottfried Kirchengast (1, 2, 3), Melannie D. Hartman (4), Clara Hohmann (1,2)
 (1) Wegener Center for Climate and Global Change, University of Graz, Austria, (2) FWF-DK Climate Change, University of Graz, Austria, (3) Institute for Geophysics, Astrophysics, and Meteorology/Institute of Physics, University of Graz, Austria, (4) Natural Resource Ecology Laboratory, Colorado State University, USA

Presented at:



Welcome, Poster Presentation and Background

Welcome

[VIDEO] <https://www.youtube.com/embed/Ah6IBjaiV4c?rel=0&fs=1&modestbranding=1&rel=0&showinfo=0>

Abstract

Sustainable agriculture and forestry is an important topic under climate change. This is a potential route for increasing long-term soil and biomass carbon storage, soil water retention capacity and for reducing water and wind erosion risks. This study uses the Styrian Raab and Enns catchment regions in Southeastern Austria as showcase regions for exploring sustainable whole-system options for climate change adaptation and mitigation under increased hot-dry conditions in agriculture and forestry. Based on dense data of the WegenerNet observing network and further hydrometeorological data, combined with hydrological modelling (WaSiM), the current hydrological disturbance potential in the Southeastern Austria focus regions is assessed. Furthermore, downscaled IPCC climate change scenarios are used for future projections and the results are evaluated for increasing heat and drought risks. This work provides the hydrological context for modelling the soil water and carbon storage enhancement options that farming, forestry and land-use practices might apply. A first key study aspect in this context is the sustainable potential of bioenergy crops. Using the local-scale WegenerNet data combined with Harmonized World Soil Database (HWSD) soil data, potential yields for bioenergy from lignocellulosic biomass (forest and Miscanthus, willow and poplar) is modelled using MiscanFor, SalixFor, PopFor and DAYCENT models for representative local areas in the showcase regions. Using DAYCENT biogeochemical modelling with different agricultural, forest management, and land use practices under climate change, sustainable system options under different future climate change scenarios are developed. These results will be used in turn to develop whole-system options, namely to jointly achieve increase of soil carbon and robustness of soil water retention capacity, increase of soil quality, reduction of soil erosion and degradation, reduced compaction, stabilisation of slopes, sustainability and resilience in the soil as well as the agricultural and forest production system. Sustainable whole-system options for farmers and forest managers are the major study focus and will supply data for local estimates in the chosen representative areas in the showcase regions as well as for upscaling to country level.


Background

Sustainable agriculture and forestry are important topics under climate change. This is a potential route for increasing long-term soil and biomass carbon storage, soil water retention capacity and for reducing water and wind erosion risks. Bioenergy crops, especially second-generation crops, like *Miscanthus*, willow and poplar, are considered to be a potential option for climate mitigation and for meeting the targets of the Paris Agreement in Europe.

Methods

Methods

DayCent was set up using the local-scale WegenerNet data combined with measured soil data (pH, soil structure) from the Austrian Chamber of Agriculture. All DayCent simulations (fert1 scenario) started with the same initial conditions on January 1, 2007, and ended on December 31, 2019. The agricultural system before was run as a continuous maize system because maize is the dominant agricultural crop in the research area. For *Miscanthus*, no fertilizer was added, and we assumed that the crop was harvested every year at day 350. For poplar and willow, 5.6 gN/m² fertilizer were added each year and harvest was scheduled every fourth year. For the maize and soybean rotation, in the year 2007 maize was planted in spring and 2.8 gN/m² and later in the year again 3.04 gN/m² fertilizer were added. In 2008, maize was grown following the prescribed pattern. In 2009, soybean was planted, and no fertilizer added. This by-yearly pattern was followed until the end of the simulations.




Location and Data

The Styrian Raab catchment in Southeastern Austria was used as a showcase region for exploring sustainable whole-system options for climate change adaptation and mitigation under increased hot-dry conditions.

Information about [Styria](#)

Fig. 1: Upper panel: The red cross shows the location of Hofstätten an der Raab. The map of the Raabtal region (RTR) located in Southeastern Austria (see lower left inset), including the Styrian Raab catchment (red; with upper, middle, and lower sub-basins in different colors). Digital elevation model topography and other information elements such as station locations are shown as background and complementary information (see legend; Southeastern Styria (SES) and Feldbach region (FBR, WegenerNet) are intensively studied sub-regions; Kabas et al. 2011; Kirchengast et al. 2014). Lower panels: summer (JJA) temperature (left) and precipitation (right) anomaly time series (relative to the 1971–1990 average) and corresponding trends over 1971 to 2016, for different sub-regions and climate-quality stations in the catchment region. (Hohmann et al. 2018)



Bioenergy test site Hofstätten an der Raab




Fig- 2: Hofstätten bioenergy site (Google, CNES/Airbus, Maxar Technologies 2020)

WegenerNet data

Based on dense data of the WegenerNet observing network and further hydrometeorological data, combined with hydrological modelling ([WaSiM](#)), the current hydrological disturbance potential in the Southeastern Austria focus regions is assessed.

WegenerNet [home](#)

WegenerNet [data portal](#)

WegenerNet 3D - [Video](#) Installing Precipitation Radar FURUNO WR 2120 on ORS mast Stradnerkogel

The Sustainability Potential of Bioenergy Crops

Potential yields (as primary use) for bioenergy from lignocellulosic biomass (*Miscanthus*, willow and poplar) are modelled using [MiscanFor](#), [SalixFor](#), [PopFor](#) and [DayCent](#) models for Hofstätten bioenergy site. The synergetic potential of bioenergy crops, especially the increase of soil carbon, is shown in the following figures. This so-called “secondary use” is the primary focus of this research. For comparison with the potential of *Miscanthus*, willow and poplar, the usual agricultural rotation of maize and soybean is modelled with DayCent at the same site.




Fig. 3: Aboveground live and dead biomass carbon stored in *Miscanthus*(misc), willow) and poplar at Hofstätten bioenergy site modelled with DayCent using the fert1 scenario. The bioenergy crops were planted in 2007 and the graph shows the development until the end of 2019. The different harvest schedule for *Miscanthus* (yearly) and willow or poplar (every fourth year) is obvious, and biomass stored in live and dead plant parts follows this schedule.




Fig. 4: Biomass carbon from stems and leaves harvested from Miscanthus, willow and poplar at Hofstätten bioenergy site modelled with DayCent using the fert1 scenario over the years 2007-2019. Miscanthus allows for a regular harvest of biomass carbon while willow and poplar harvest events are spiking events.




Fig. 5: Cumulative carbon removed during tree management events from Miscanthus, willow and poplar at Hofstätten bioenergy site modelled with DayCent using the fert1 scenario for the years 2007-2019.




Fig. 6: Soil organic matter carbon (SOMC) stored under Miscanthus, willow and poplar at Hofstätt bioenergy site modelled with DayCent. The bioenergy crops were planted in 2007 and the graph shows the development until the end of 2019. The situation at the beginning of the graph represents the historic agricultural rotation before the establishment of bioenergy crops. SOMC slowly decreases over time in the agricultural rotation and increases with bioenergy crops.




Fig. 7: Comparison of the N₂O fluxes from agricultural crops, Miscanthus, willow and poplar at Hofstätten bioenergy site modelled with DayCent for the years 2007-2019. Agricultural N₂O fluxes are intermediate while fluxes from Miscanthus are sometimes higher. This corresponds with larger rain events (2018). Willow and poplar stabilize the N₂O fluxes most effectively which may correspond with the fact that they need larger amounts of water.

Discussion and Conclusions

This research has shown that there is a large potential for soil carbon increase by lignocellulosic bioenergy crops compared with conventional agricultural rotation. Figure 6 shows the development of soil organic matter carbon under bioenergy crops and conventional agricultural rotation. While soil organic matter under agriculture slowly decreases over time, there is a significant increase under the bioenergy crops in this research. This is mostly caused by less frequent harvest, decreased tillage and by the growth pattern of bioenergy crops themselves (e.g. extensive root system). Especially *Miscanthus* has a high potential for stabilizing soils because it is normally harvested while dormant, which stabilizes soil carbon but also the carbon stored in plant material.

Outlook and Contact Information

Outlook

The DayCent model will allow computing a GHG budget for all land use options and potential new systems of land use. These results will be used in turn to develop whole-system options, namely to jointly achieve an increase of soil carbon and robustness of soil water retention capacity, an increase of soil quality, reduction of soil erosion and degradation, reduced compaction, stabilization of slopes, sustainability and resilience in the soil as well as the agricultural and forest production system. Sustainable whole-system options for farmers and forest managers are the major study focus. These will supply data for local estimates in the chosen representative areas in the showcase regions as well as for upscaling to country and European level.

Contact information

Dagmar Henner dagmar.henner@uni-graz.at

Author Information

www.linkedin.com/in/dagmar-henner-09580352



Der Wissenschaftsfonds.

This work was funded by the Austrian Science Fund (FWF) under Research Grant W1256 (Doctoral Programme Climate Change: Uncertainties, Thresholds and Coping Strategies)

REFERENCES

Clifton-Brown, J., Harfouche, A., Casler, M. D., Jones, H. D., Macalpine, W. J., Murphy-Bokern, D., ... Lewandowski, I. (2019). Breeding progress and preparedness for mass-scale deployment of perennial lignocellulosic biomass crops switchgrass, miscanthus, willow and poplar. *GCB Bioenergy*, 11(1), 118–151. <https://doi.org/10.1111/gcbb.12566>

Del Grosso, S. J., Ahuja, L. R., Parton, W. J., Del Grosso, S., Ahuja, L., & Parton, W. (2016). Modeling GHG Emissions and Carbon Changes in Agricultural and Forest Systems to Guide Mitigation and Adaptation: Synthesis and Future Needs. In *Advances in Agricultural Systems Modeling*. American Society of Agronomy, Inc., Crop Science Society of America, Inc., and Soil Science Society of America, Inc. <https://doi.org/10.2134/advagricsystmodel6.2013.0008>

Dondini, M., Abdalla, M., Aini, F. K., Albanito, F., Beckert, M. R., Begum, K., ... Smith, P. (2018). Chapter 9—Projecting Soil C Under Future Climate and Land-Use Scenarios (Modeling). In B. K. Singh (Ed.), *Soil Carbon Storage* (pp. 281–309). Academic Press. <https://doi.org/10.1016/B978-0-12-812766-7.00009-3>

- Fuchsberger, J., Kirchengast, G., Bichler, C., Leuprecht, A., and Kabas, T.: WegenerNet climate station Network Level 2 data version 7.1 (2007 - 2018), Tech. rep., Wegener Center for Climate and Global Change, University of Graz, Graz, Austria, <https://doi.org/10.25364/WEGC/WPS7.1:2019.1>
- Hastings, A., Clifton-Brown, J., Wattenbach, M., Mitchell, C. P., & Smith, P. (2009a). The development of MISCANFOR, a new *Miscanthus* crop growth model: Towards more robust yield predictions under different climatic and soil conditions. *GCB Bioenergy*, 1(2), 154–170. <https://doi.org/10.1111/j.1757-1707.2009.01007.x>
- Hastings, A., Clifton-Brown, J., Wattenbach, M., Mitchell, C. P., Stampfl, P., & Smith, P. (2009b). Future energy potential of *Miscanthus* in Europe. *GCB Bioenergy*, 1(2), 180–196. <https://doi.org/10.1111/j.1757-1707.2009.01012.x>
- Hastings, A., Mos, M., Yesufu, J. A., McCalmont, J., Schwarz, K., Shafei, R., ... Clifton-Brown, J. (2017). Economic and Environmental Assessment of Seed and Rhizome Propagated *Miscanthus* in the UK. *Frontiers in Plant Science*, 8. <https://doi.org/10.3389/fpls.2017.01058>
- Hastings, A., Tallis, M. J., Casella, E., Matthews, R. W., Henshall, P. A., Milner, S., ... Taylor, G. (2014). The technical potential of Great Britain to produce ligno-cellulosic biomass for bioenergy in current and future climates. *GCB Bioenergy*, 6(2), 108–122. <https://doi.org/10.1111/gcbb.12103>
- Hastings, A., Yeluripati, J., Hillier, J., & Smith, P. (2013). Biofuel Crops and Greenhouse Gases. In B. P. Singh (Ed.), *Biofuel Crop Sustainability* (pp. 383–405). John Wiley & Sons, Ltd. Retrieved from <http://onlinelibrary.wiley.com/doi/10.1002/9781118635797.ch12/summary>
- Henner, D. N. (2019). The potential for second-generation bioenergy crops in Europe and their impact on soil carbon changes and erosion. (PhD Thesis). University of Aberdeen, Aberdeen, UK.
- Henner, D. N., Hastings, A., Pogson, M., McNamara, N. P., Davies, C. A., & Smith, P. (2020). PopFor: A new model for estimating poplar yields. *Biomass and Bioenergy*, 134, 105470. <https://doi.org/10.1016/j.biombioe.2020.105470>
- Hohmann, C., Kirchengast, G. & Birk, S. Alpine foreland running drier? Sensitivity of a drought vulnerable catchment to changes in climate, land use, and water management. *Climatic Change* 147, 179–193 (2018). <https://doi.org/10.1007/s10584-017-2121-y>
- Kirchengast G, Kabas T, Leuprecht A et al (2014) WegenerNet: a pioneering high-resolution network for monitoring weather and climate. *Bull Am Meteorol Soc* 95:227–242. <https://doi.org/10.1175/BAMS-D-11-00161.1>

Parton, W. J., Schimel, D. S., Cole, C. V., & Ojima, D. S. (1987). Analysis of Factors Controlling Soil Organic Matter Levels in Great Plains Grasslands. *Soil Science Society of America Journal*, 51(5), 1173–1179. <https://doi.org/10.2136/sssaj1987.03615995005100050015x>

Parton, William J., Gutmann, M. P., Merchant, E. R., Hartman, M. D., Adler, P. R., McNeal, F. M., & Lutz, S. M. (2015). Measuring and mitigating agricultural greenhouse gas production in the US Great Plains, 1870–2000. *Proceedings of the National Academy of Sciences*, 201416499. <https://doi.org/10.1073/pnas.1416499112>

Parton, William J., Hartman, M., Ojima, D., & Schimel, D. (1998). DAYCENT and its land surface submodel: Description and testing. *Global and Planetary Change*, 19(1–4), 35–48. [https://doi.org/10.1016/S0921-8181\(98\)00040-X](https://doi.org/10.1016/S0921-8181(98)00040-X)

Richards, M., Pogson, M., Dondini, M., Jones, E. O., Hastings, A., Henner, D. N., ... Smith, P. (2017). High-resolution spatial modelling of greenhouse gas emissions from land-use change to energy crops in the United Kingdom. *GCB Bioenergy*, 9(3), 627–644. <https://doi.org/10.1111/gcbb.12360>

Schulla J (2015) Model description WaSiM. Technical report, www.wasim.ch