

# The geologically supervised spectral investigation as a key methodology for identifying volcanically active areas on Venus

Piero D’Incecco<sup>1</sup>, Justin Filiberto<sup>2</sup>, Iván López<sup>3</sup>, Dmitry Gorinov<sup>4</sup>, Goro Komatsu<sup>1</sup>, Alexey Martynov<sup>5</sup>, and Pavel Pisarenko<sup>5</sup>

<sup>1</sup>Università d’Annunzio

<sup>2</sup>Lunar and Planetary Institute

<sup>3</sup>Universidad Rey Juan Carlos - Campus de Móstoles

<sup>4</sup>Space Research Institute of Russian Academy of Sciences (IKI)

<sup>5</sup>Scientific and Production Association. S.A. Lavochkin

November 26, 2022

## Abstract

Combining geologic mapping and stratigraphic reconstruction of lava flows at Sapas, Maat and Ozza Montes, three potentially young volcanic structures of Atla Regio on Venus, with analysis of the spectral signature (radar emissivity anomalies) characterizing each mapped flow, Brossier et al. (2021) conclude that some of the lava flows at Maat Mons may be geologically recent (~25 Ma). The lava flows of Sapas and Ozza Montes are more consistent with weathered lava flows forming chlorapatite and some perovskite oxides. We discuss the reasons why, besides the importance of the results they obtained, the methodology they used can be very valuable for future investigations with higher resolution datasets. The importance of combining geologic interpretation with spectral analysis in the reconstruction of the volcanic history of Venus Considering its size, gravity and the presence of an atmosphere, Venus is typically considered as the twin sister of the Earth, but despite the apparent similarities with our planet, Venus is notably different because it is characterized by its extreme surface environment. With 90 bars and 475 °C, its surface is a very inhospitable place for life as we know it. Venus does not show evidence for a present plate tectonics-like activity, having a major part of its surface volcanic deposits younger than 300 Ma. It has been hypothesized that Venus underwent a catastrophic event of global resurfacing about 300 Ma ago, which may have almost entirely rejuvenated its surface (Schaber, 1992; Nimmo and McKenzie, 1998; Romeo and Turcotte, 2010; Strom et al., 1994; Turcotte et al., 1999). Some other studies instead favor a more equilibrium resurfacing model of the surface (Phillips et al, 1992; Phillips and Hansen, 1994; Bjonnes et al., 2012; O’Rourke and Korenaga, 2015). It is also possible that the past volcanic history of Venus somehow reflected an intermediate situation between these two end-member scenarios. Related to this topic, there has been a subject of debate whether or not the volcanism on Venus is currently evolving toward an equilibrium stage, with occurrences of smaller and more frequent localized eruptions. In this regard, it is vital to identify areas with current or recent volcanism, to measure the actual rate and volume of the most recent volcanic eruptions. The geologic interpretation and analysis of spectral signatures (both in radar and infrared wavelengths) can help us constraining the age of surface volcanic deposits on Venus. In geology, the so called “cross-cutting interrelationships” can constrain the relative age of two lava flows as it has been applied to young, possibly very recent lava flows and tectonic features on

## **The geologically supervised spectral investigation as a key methodology for identifying volcanically active areas on Venus.**

P. D’Incecco<sup>1,2</sup>, J. Filiberto<sup>3</sup>, I. López<sup>4</sup>, D.A. Gorinov<sup>5</sup>, G. Komatsu<sup>1,6</sup>, A. Martynov<sup>7</sup>, P. Pisarenko<sup>7</sup>

<sup>1</sup> *Dipartimento di Ingegneria e Geologia (INGEO), Università d'Annunzio, Viale Pindaro 42, 65127 Pescara, Italy*

<sup>2</sup> *Arctic Planetary Science Institute (APSI), Rovaniemi, Finland.*

<sup>3</sup> *Lunar and Planetary Institute, USRA, 3600 Bay Area Blvd Houston, TX 77058*

<sup>4</sup> *Universidad Rey Juan Carlos, Madrid, Spain*

<sup>5</sup> *Space Research Institute of the Russian Academy of Sciences, Profsoyuznaya st., 84/32 Moscow, Russia*

<sup>6</sup> *International Research School of Planetary Sciences, Università d'Annunzio, Viale Pindaro 42, 65127 Pescara, Italy*

<sup>7</sup> *Lavochkin Association, Leningradskaya 24, Khimki, Russia.*

### **Abstract**

Combining geologic mapping and stratigraphic reconstruction of lava flows at Sapas, Maat and Ozza Montes, three potentially young volcanic structures of Atla Regio on Venus, with analysis of the spectral signature (radar emissivity anomalies) characterizing each mapped flow, Brossier et al. (2021) conclude that some of the lava flows at Maat Mons may be geologically recent (~25 Ma). The lava flows of Sapas and Ozza Montes are more consistent with weathered lava flows forming chlorapatite and some perovskite oxides. We discuss the reasons why, besides the importance of the results they obtained, the methodology they used can be very valuable for future investigations with higher resolution datasets.

### **The importance of combining geologic interpretation with spectral analysis in the reconstruction of the volcanic history of Venus**

Considering its size, gravity and the presence of an atmosphere, Venus is typically considered as the twin sister of the Earth, but despite the apparent similarities with our planet, Venus is notably different because it is characterized by its extreme surface environment. With 90 bars and 475 °C, its surface is a very inhospitable place for life as we know it. Venus does not show evidence for a present plate tectonics-like activity, having a major part of its surface volcanic deposits younger than 300 Ma. It has been hypothesized that Venus underwent a catastrophic event of global resurfacing about 300 Ma ago, which may have almost entirely rejuvenated its surface (Schaber, 1992; Nimmo and McKenzie, 1998; Romeo and Turcotte, 2010; Strom et al., 1994; Turcotte et al., 1999). Some other studies instead favor a more equilibrium resurfacing model of the surface (Phillips et al, 1992; Phillips and Hansen, 1994; Bjonnes et al., 2012; O’Rourke and Korenaga, 2015). It is also possible that the past volcanic history of Venus somehow reflected an intermediate situation between these two end-member scenarios. Related to this topic, there has been a subject of debate whether or not the volcanism on Venus is currently evolving toward an equilibrium stage, with occurrences of smaller and more frequent localized eruptions. In this regard, it is vital to identify areas with current or recent volcanism, to measure the actual rate and volume of the most recent volcanic eruptions. The geologic interpretation and analysis of spectral signatures (both in radar and infrared wavelengths) can help us constraining the age of surface volcanic deposits on Venus.

In geology, the so called “cross-cutting interrelationships” can constrain the relative age of two lava flows as it has been applied to young, possibly very recent lava flows and tectonic features on

Venus (i.e., Figure 6 in D’Incecco et al., 2020). The spectral analysis can provide some additional constraints on the ages of surface volcanic materials. We know that on Venus recently erupted lava flows become rapidly altered when they come in contact with the thick and chemical active atmosphere. This process is called chemical weathering. In general, unweathered materials are characterized by high emissivity anomalies in the infrared band and high emissivity anomalies in the radar band (at certain altitudes) and such materials can be considered as geologically recent. Recent laboratory analyses on the oxidation rate of igneous minerals showed that such chemical weathering on Venus may act on the order of weeks or months (Fegley et al. 1995; Berger et al. 2019; Filiberto et al., 2020; Cutler et al., 2020; Treiman et al., 2021). This implies that areas on Venus with high infrared emissivity anomalies (and high radar emissivity) may be volcanically active at the present day.

The present manuscript presented by Brossier et al. (2021) combines geologic interpretation and consequent stratigraphic reconstruction with spectral analysis of radar emissivity anomalies observed at a number of lava flows, using a peculiar technique which may be defined as a geologically supervised spectral investigation. The authors find spatial correlations between many lava flow units and radar emissivity excursions at different altitudes, over three volcanic structures of Atla Regio; Sapas, Maat and Ozza Montes (Figure 1). At the three volcanic structures, for a given altitude and temperature, low emissivity excursions do not occur uniformly over all the mapped units and surface materials. This implies that the observed low emissivity excursions are strongly controlled by the presence of distinct ferroelectric minerals with high dielectric constant.

The authors propose that a group of ferroelectric minerals can explain the observed low emissivity excursions. In particular, chlorapatite and four perovskites can account for any of the emissivity excursions observed at the three volcanoes. They indicate that ferroelectric (high dielectric) minerals can be intrinsic of a lava flow in the form of direct crystallization or, alternatively, can be produced by the surface-atmosphere chemical interactions over the time. Furthermore, the authors assume that – for a given composition and atmospheric condition - the low radar emissivity excursions can be then used as a chronometer for estimating the relative ages of the lava flow units. Sharp low emissivity excursions (high dielectric constants) will indicate older (more weathered) surface materials. Building on the degradation model of dark haloes surrounding some impact craters on Venus, the authors also provide further constraints in terms of absolute ages of the volcano-tectonic activity in the study area.

The investigation conducted with this work, which correlates morphologic mapping with radar properties of surface materials on Venus is elegant and innovative. The methodology used could provide a significant contribution to the key debate regarding the style of resurfacing on Venus. Their results demonstrate what can be still achieved using the (relatively low resolution) Magellan radar dataset but it also shows what could be potentially be obtained by higher resolution radar data hopefully obtained from future missions, such as European Space Agency’s EnVision mission (Ghail et al., 2012, 2020) or NASA’s Venus Emissivity, Radio Science, InSAR, Topography & Spectroscopy (VERITAS) mission (Smrekar et al., 2015), both are currently in the conceptual and technological development. The Deep Atmosphere of Venus Investigation of Noble gases, Chemistry and Imaging, Plus (DAVINCI+) (Glaze et al., 2017; 2018; Garvin et al., 2020) and Venera-D (Senske et al., 2017; Zasova et al., 2019) mission concepts plan instead to provide more detailed data on the structure and thermal profiles of the Venusian atmosphere. DAVINCI+ will also image the surface below the cloud deck, while Venera-D will obtain and analyze a sample of the surface material at the landing site. In this regard, new experimental data obtained in the laboratory will help interpret the data observed through remote sensing. In addition to those mentioned above, a Venus Flagship mission concept is currently being developed, with the main


goal of unveiling new clues about the geologic history of Venus (Bullock et al., 2009; Gilmore et al., 2019).

### Data Availability Statement

For this commentary article, no new data were used. The data we commented in this article come from the previously published research by Brossier et al. (2021), on this journal.

### Acknowledgements

P. D’Incecco thanks the European Union for the financial support through the “Programma Operativo Nazionale” (PON) Attraction and International Mobility (AIM) grant AIM1892731. This is APSI contribution no. 15.



**Figure 1.** Sketch maps of lava flow units within three volcanoes: (top-left) Sapas, (top right) Maat, and (bottom) Ozza. Crosses (+) indicate the summit(s) of each volcano. Black box indicates an isolated patch of O2 unit. From Brossier et al. (2021).

## References

- Bjornes, E.E., Hansen, V.L., James, B., Swenson, J.B., 2012. Equilibrium resurfacing of Venus: Results from new Monte Carlo modeling and implications for Venus surface histories. *Icarus* 217, pp. 451-461. <https://doi.org/10.1016/j.icarus.2011.03.033>
- [Brossier, J., Gilmore, M.S., Toner, K., Stein, A.J., 2020. Supporting information for distinct mineralogy and age of individual lava flows in Atla Regio, Venus derived from Magellan radar emissivity. Figshare. https://doi.org/10.6084/m9.figshare.13331210.v2](https://doi.org/10.6084/m9.figshare.13331210.v2)
- Brossier, J., Gilmore, M.S., Toner, K., Stein, A.J., 2021. Distinct mineralogy 1 and age of individual lava flows in Atla Regio, Venus derived from Magellan radar emissivity. *J. Geophys. Res.* <https://doi.org/10.1029/2020JE006722>.
- Bullock, M., et al., 2009. Venus Flagship Mission Study: Report of the Venus Science and Technology Definition Team. *Geography*.
- Cutler, K.S., Filiberto, J., Treiman, A.H., Trang, D., 2020. Experimental Investigation of Oxidation of Pyroxene and Basalt: Implications for Spectroscopic Analyses of the Surface of Venus and the Ages of Lava Flows. *Planet. Sci. J.*, 1, 21. <https://doi.org/10.3847/psj/ab8faf>
- D’Incecco, P., López, I., Komatsu, G., Ori, G.G., Aittola, M., 2020. Local stratigraphic relations at Sandel crater, Venus: Possible evidence for recent volcano-tectonic activity in Imdr Regio. *Earth Planet. Sci. Lett.* 546. <https://doi.org/10.1016/j.epsl.2020.116410>.
- Filiberto, J., Trang, D., Treiman, A.H., Gilmore, M.S., 2020. Present-day volcanism on Venus as evidenced from weathering rates of olivine. *Sci. Adv.* 6, 1. <https://doi.org/10.1126/sciadv.aax7445>
- Garvin, J., Arney, G., Getty, S., Johnson, N., Kiefer, W., Lorenz, R., Ravine, M., Malespin, C., Webster, C., Campbell, B., 2020. DAVINCI+: Deep Atmosphere of Venus Investigation of Noble Gases, Chemistry, and Imaging Plus. 51st Lunar and Planetary Science Conference, Abstract #2326.
- Gilmore, M.S., Beauchamp, P.M., and the 2019 Venus Flagship Mission Study Team. Proposed Venus Flagship mission. Abstracts of the tenth Moscow Solar System Symposium, 10MS3-VN-10, 2019, pp. 84–86.
- Glaze, L.S., Garvin, J.B., Robertson, B., Johnson, N.M., Amato, M.J., Thompson, J., Goodloe, C., Everett, D., 2017. DAVINCI: Deep atmosphere venus investigation of noble gases, chemistry, and imaging, in: *IEEE Aerospace Conference Proceedings*. <https://doi.org/10.1109/AERO.2017.7943923>

- Glaze, L.S., Wilson, C.F., Zasova, L.V., Nakamura, M., Limaye, S., 2018. Future of Venus Research and Exploration. *Space Science Reviews* 214, 89.
- Nimmo, F., McKenzie, D., 1998. Volcanism and tectonics on Venus. *Annu. Rev. Earth Planet. Sci.* 26, pp. 23-51. <https://doi.org/10.1146/annurev.earth.26.1.23>
- O'Rourke, J.G., Korenaga, J., 2015. Thermal evolution of Venus with argon degassing. *Icarus* 260, pp. 128-140. <https://doi.org/10.1016/j.icarus.2015.07.009>
- Phillips, R.J., Hansen, V.L., 1994. Tectonic and magmatic evolution of Venus. *Annu. Rev. Earth Planet. Sci.* 22, pp. 597-654. <https://doi.org/10.1146/annurev.earth.22.1.597>
- Phillips, R.J., Raubertas, R.F., Arvidson, R.E., Sarkar, I.C., Herrick, R.R., Izenberg, N., and Grimm, R.E., 1992. Impact craters and Venus resurfacing history, *J. Geophys. Res.*, 97(E10), pp. 15923– 15948, doi:10.1029/92JE01696.
- Romeo, I., Turcotte, D.L., 2010. Resurfacing on Venus. *Planet. Space Sci.* 58, 10, pp. 1374-1380. <https://doi.org/10.1016/j.pss.2010.05.022>
- Schaber, G.G., 1992. Geology and distribution of impact craters on Venus: what are they telling us? *J. Geophys. Res.* <https://doi.org/10.1029/92je01246>.
- Senske, D.A., Zasova, L. V., Ignatiev, N.I., Korablev, O., Eismont, N., Gerasimov, M., Ivanov, M.A., Martynov, M., Khatuntsev, I. V., Limaye, S.S., Jessup, K.L., Economou, T., Esposito, L.W., 2017. Venera-D: Expanding our Horizon of Terrestrial Planet Climate and Geology through the Comprehensive Exploration of Venus, Report of the Venera-D Joint Science Definition Team.
- Strom, R.G., Schaber, G.G., Dawson, D.D., 1994. The global resurfacing of Venus. *J. Geophys. Res.* <https://doi.org/10.1029/94je00388>
- Turcotte, D.L., Morein, G., Roberts, D., Malamud, B.D., 1999. Catastrophic Resurfacing and Episodic Subduction on Venus. *Icarus* 139, 1, pp. 49-54. <https://doi.org/10.1006/icar.1999.6084>
- Zasova, L.V., Gorinov, D.A., Eismont, N.A. et al., 2019. Venera-D: A Design of an Automatic Space Station for Venus Exploration. *Sol Syst Res* 53, pp. 506–510. <https://doi.org/10.1134/S0038094619070244>