

# A Seismo-Acoustic Investigation of a Localized Crater Terrace Collapse at Stromboli Volcano

Bryan Rosenblatt<sup>1,1</sup>, Jeffrey Johnson<sup>1,1</sup>, Kristina Rossavik<sup>1,1</sup>, Ulrich Kueppers<sup>2,2</sup>, Tullio Ricci<sup>3,3</sup>, and Daniele Andronico<sup>4,4</sup>

<sup>1</sup>Boise State University

<sup>2</sup>Ludwig Maximilian University of Munich

<sup>3</sup>National Institute of Geophysics and Volcanology

<sup>4</sup>Istituto Nazionale di Geofisica e Vulcanologia

November 30, 2022

## Abstract

Interpreting seismo-acoustic signals is critical for assessing and characterizing changes in volcanic vents and has implications for interpreting volcanic unrest. This is especially relevant for Stromboli volcano (Italy), an active stratovolcano with a complex plumbing system, continuous activity, and recurring paroxysms. Stromboli is known for its consistent Strombolian style of eruption, multiple active vents on its crater terrace, and for occasional structural modifications including explosive excavation and/or collapsing craters due to near-surface changes to the plumbing system. This study addresses a single localized collapse of the crater terrace, occurring in May of 2019, when one of Stromboli's vents changed from a pronounced hornito to a pit crater, resulting in a shift in eruption style at this vent from jetting to Strombolian. The days before and after this transition were recorded with eight infrasound sensors and three seismic geophones located on the crater terrace. We investigate the seismo-acoustic timing of these signals as well as the ratio between seismic and acoustic energy to identify changes associated with eruptive signals and associated variations in location of the eruptive sources. This work highlights the effectiveness of seismo-acoustic data analysis, provides insight into Stromboli's structural modifications, and builds a foundation for focused analysis of seismo-acoustic signals associated with Stromboli and other open-vent volcanic systems.


# A Seismo-Acoustic Investigation of a Localized Crater Terrace Collapse at Stromboli Volcano

Bryan B. Rosenblatt<sup>1</sup>, Jeffrey B. Johnson<sup>1</sup>, Kristina Rossavik<sup>1</sup>, Ulrich Kueppers<sup>2</sup>, Tullio Ricci<sup>3</sup>, Daniele Andronico<sup>4</sup>

1.) Department of Geoscience, Boise State University 2.) Experimental & Physical Volcanology, Ludwig-Maximilians-Universität München (LMU)  
3.) Istituto Nazionale di Geofisica e Vulcanologia, Sezione Roma 1, Rome, Italy 4.) Istituto Nazionale di Geofisica e Vulcanologia, Etna Observatory

## INTRODUCTION/OBJECTIVES

Stromboli volcano is an active island stratovolcano with a complex plumbing system and recurring paroxysms. It is known for its consistent Strombolian eruption style, multiple active vents (fig right), and regular vent evolutions<sup>3</sup> to its crater terrace. This study focuses on a single localized collapse of the crater terrace (15 May 2019) where the S1 vent changed from a spatter cone to a pit crater. This collapse resulted in a change in eruption style at this vent from jetting to Strombolian. We investigate the seismo-acoustic arrival times and energy ratios between our co-located acoustic and seismic signals before and after the collapse to identify changes associated with eruptive signal and source location.




## THE COLLAPSE

14-May 2019: jetting spatter cone 15-May 2019: Strombolian pit crater



## SEISMO-ACOUSTIC TIMING


To investigate the timing information across the seismic and acoustic signals, we beamstack, or sum, all events of the same type. As seen in the plot below (fig below), the timing information from the S1 vent between the co-located geophone and *InfraBSUs* cumulative signals does not seem to change much after the May 15<sup>th</sup> collapse. Prior to the collapse, the acoustic signal reached the sensor ~204 samples (~0.51 seconds) after the initial onset of the seismic signal. Post-collapse, the acoustic signal reached the sensor ~202 samples (~0.505 seconds) after the initial onset of the seismic signal. This result shows an insignificant change in the arrival times of ~2 samples (~0.005 seconds). Thus, we assume the source for the S1 vent is non-changing throughout this period. This also leads to the reasonable assumption that there is no significant change to the locations of the seismic and acoustic sources.



## DATA AND METHODS

Data from the terrace of Stromboli was collected May 11-15, 2019, with two 3-channel infrasound arrays (*InfraBSUs*) and six co-located vertical geophones about 200-250 m from the S1 vent. Sensor data and GPS locations were logged via the Omnirecs DATA-CUBE. Acoustic and seismic data were both sampled at 400 Hz.

Acoustic signals are high-passed above 1 Hz and seismic signals are low-passed below 4 Hz. Using time lags and high correlation values across the arrays, ten events for each eruption style (jetting spatter cone, 14 May; Strombolian pit crater, 15 May) are selected. All acoustic events are shifted to align their initial pressure onsets. Each seismic event is then shifted by the same amount as its acoustic counterpart (figs below). We compare the cumulative event arrival times across the co-located sensors as well as changes to these co-located sensors' volcano acoustic seismic ratio (VASR) from before to after the collapse.



## SEISMO-ACOUSTIC ENERGY RATIO


In order to properly characterize the acoustic and seismic energy of discrete eruptive events, we integrate over a time window spanning 5 seconds prior to the event onset to a time where the acoustic and seismic signals reach background levels. Using formulas along with standard density and propagation speed values detailed by Johnson & Aster<sup>1</sup> (Eq 1 & 2), we calculate an approximate energy release (kJ) for each individual event. These summed energies are then divided by their co-located counterpart to produce VASR values (Eq 3, fig left). In general, there is more acoustic energy and a similar amount of seismic energy post-collapse.

$$E_{acoustic} \propto \frac{2\pi r^2}{\rho_{atmos} c_{atmos}} \int \Delta P(t)^2 dt \quad (1)$$

$$E_{seismic} \propto 2\pi r^2 \rho_{Earth} c_{Earth} \int V(t)^2 dt \quad (2)$$

$$VASR = \frac{E_{acoustic}}{E_{seismic}} \quad (3)$$

We find lower VASR values before the localized collapse (fig right). Since the spatter cone's vent is narrower<sup>2</sup> and there's more acoustic energy after the collapse, it is reasonable to assume that the spatter cone vent released more elastic energy (seismic) into the earth resulting in a lower VASR (Eq 3).



## CONCLUSIONS

This study investigates a single localized collapse of the crater terrace of Stromboli volcano, which resulted in a change of eruption style from jetting to Strombolian. The moments before and after this collapse were recorded with co-located acoustic and seismic sensors. We found that there was no significant change in the seismo-acoustic timing from before to after the collapse, indicating that the overall source as well as the source location for both signals is non-changing. In addition, we found that the VASR increased from before to after the collapse indicating that more elastic energy was released into the earth (as seismic waves) before the collapse. This study only includes a total of 20 events from this expedition. More research as well as more events are needed to produce a robust outcome from this dataset.

## REFERENCES/ACKNOWLEDGMENTS

- References:**
- Johnson, J. B., & Aster, R. C. (2005). Relative partitioning of acoustic and seismic energy during Strombolian eruptions. *Journal of Volcanology and Geothermal Research*, 148(3-4), 334-354. <https://doi.org/10.1016/j.jvolgeores.2005.05.002>
  - Schmid et al. (2021). Characterising vent and crater shape changes at Stromboli: Implications for risk areas. *Volcanica*, 4(1), 87-105. <https://doi.org/10.30909/vol.04.01.87105>
  - Schmid et al. (2020). Release characteristics of overpressurised gas from complex vents: implications for volcanic hazards. *Bulletin of Volcanology*, 82(11). <https://doi.org/10.1007/s00445-020-01407-2>
  - Suckale et al. (2016). Flow-to-fracture transition in a volcanic mush plug may govern normal eruptions at Stromboli. *Geophysical Research Letters*, 43(23), 12,071-12,081. <https://doi.org/10.1002/2016GL071501>
- Acknowledgments:**  
Scott Gauvain, Amanda Kuhlman, Armando Pineda, Ashley Bosa, Jerry Mock, Jacob Anderson  
NSF grant number 1830976