

# What controls the ice nucleating ability of sea spray aerosols at cirrus temperatures?

Ryan Patnaude<sup>1</sup>

<sup>1</sup>Colorado State University

November 21, 2022

## Abstract

Cirrus clouds are frequently observed in the upper troposphere and have a strong impact on Earth's radiative balance. However, significant uncertainties remain regarding the role aerosols play in cirrus cloud formation and evolution. Due to their abundance in the atmosphere sea spray aerosols (SSA) have recently been identified as potential ice nucleating particles in cirrus clouds. Using a continuous flow diffusion chamber (CFDC), the ice nucleating ability of size-selected SSA and sodium chloride (NaCl) particles is investigated at temperatures  $< 235$  K. Above  $\sim 220$  K, the majority of SSA and NaCl particles fully deliquesce and freeze via homogeneous nucleation near water relative humidities RH<sub>w</sub> of  $\sim 95\%$ . Below 220 K, the onset freezing RH<sub>w</sub> is much lower at  $\sim 75\%$  for SSA and NaCl, and strong heterogeneous freezing of 10% of the aerosol population is observed. The similar heterogeneous freezing behavior between SSA and NaCl, which occurs near their expected deliquescence RH<sub>w</sub> indicates the freezing of SSA at the lowest temperatures is controlled by the inorganic salt components. Lastly, ice nucleation active surface site densities are calculated and show that particle size does not dictate the heterogeneous freezing efficiency for SSA and NaCl. This study demonstrates SSA as a potentially significant source of ice nucleating particles in cirrus clouds, and may contribute further climate impacts if SSA emissions and transport change in the future.




# What controls the ice nucleating ability of sea spray aerosols at cirrus temperatures?

Ryan Patnaude, Russell Perkins, Sonia Kreidenweis, Paul DeMott  
Colorado State University, Fort Collins, CO





## Motivation and background



- Cirrus clouds are one of the most abundant cloud types composed entirely of ice below  $-38^{\circ}\text{C}$ .
- Sea spray aerosols (SSA) generated by wave breaking may be lofted to cirrus levels through deep convection and detrainment in anvils.
- Wagner et al., (2018) showed that mixed-phase solid/liquid SSA freeze heterogeneously via immersion freezing below 220 K.
- Schill & Tolbert (2014) also observed heterogeneous freezing behavior, which they could not determine as being either deposition nucleation or immersion freezing
- CSU CFDC modified to go down to lower temperatures  $< -70^{\circ}\text{C}$ , making cirrus studies possible.

## Methodology

- Continuous Flow diffusion chamber (CFDC) is a concentric cylindrical column with a “warm” outer wall (OW) and “cold” inner wall (IW)
- Linear temperature gradient between walls + nonlinear saturation vapor pressure produces supersaturation in focused lamina between ice-covered walls.
- Operates from ice saturation to water-supersaturated conditions depending on the wall temperature difference




### Experimental setup

- Used natural seawater collected at the Scripps Pier in La Jolla, CA for SSA and 3.5% wt NaCl solution.
- SSA and NaCl particles are generated by an aerosol nebulizer, and dried at room temperature.
- Used monodisperse aerosol distribution of  $\sim 150$  nm and  $\sim 600$  nm
- Particle phase trajectories (left) help describe the mode of freezing at cold temperatures

## Results

- Homogeneous freezing at temperatures  $> 220$  K.
- Heterogeneous freezing transition  $< 220$  K; Lower onset  $\text{RH}_w$  for SSA particles.
- Similar freezing onsets for both aerosol sizes.



- Higher ice active surface site density ( $n_s$ ) values for smaller particles.
- Regardless of size or surface area, nucleation onset remains the same  $\rightarrow$  violates the  $n_s$  concept and suggests it is not applicable for low temperature SSA freezing.
- Two possible pathways to explain heterogeneous nucleation onset and lack of size dependence.
  1. Immersion freezing
  2. Pore condensation freezing
- Intersection of two lines coincides with onset of heterogeneous freezing.

## Conclusions and atmospheric implications

1. NaCl and SSA experiments using CSU CFDC able to replicate heterogeneous nucleation results at cirrus temperatures, similar to previous cloud chamber studies.
2. SSA freeze via heterogeneous nucleation at  $\text{RH}_w \sim 75\%$  below 220 K at high fractions (10%).
3. Multiple possible freezing mechanisms for NaCl and SSA particles depending on T and RH.
4. SSA particles may represent a significant source of INPs at cirrus levels (see Patnaude et al., 2021).

**Acknowledgements:** This project was funded by the National Science Foundation (NSF) through the NSF Center for Aerosol Impacts on Chemistry of the Environment (NSF-CAICE), Award CHE-1801971

**References:** Schill & Tolbert (2014). *Journal of Physical Chemistry C*, 118(50), 29234–29241. Wagner et al., (2018). *Journal of Geophysical Research: Atmospheres*, 123(5), 2841–2860. Patnaude et al., (2021). *ACS Earth and Space Chemistry*, 5(9), 2196–2211.