Formulation of a Consistent Multi-Species Canopy Description for Hydrodynamic Models Embedded in Large-Scale Land-Surface Representations of Mixed-Forests

Bohrer Gil¹ and Missik Justine²

¹Ohio State University ²The Ohio State University

November 16, 2022

Abstract

The plant hydrodynamic approach represents a recent advancement to land surface modeling, in which stomatal conductance responds to water availability in the xylem rather than in the soil. To provide a realistic representation of tree hydrodynamics, hydrodynamic models must resolve processes at the level of a single modelled tree, and then scale the resulting fluxes to the canopy and land surface. While this tree-to-canopy scaling is trivial in a homogeneous canopy, mixed-species canopies require careful representation of the species properties and a scaling approach that results in a realistic description of both the canopy and individual-tree hydrodynamics, as well as leaf-level fluxes from the canopy and their forcing. Here, we outline advantages and pitfalls of three commonly used approaches for representing mixed-species forests in land surface models, and present a new framework for scaling vegetation characteristics and fluxes in mixed-species forests. The new formulation scales fluxes from the tree- to canopy-level in an energy- and mass-conservative way and allows for a consistent multi-species canopy description for hydrodynamic models.

1 2	Formulation of a Consistent Multi-Species Canopy Description for Hydrodynamic Models Embedded in Large-Scale Land-Surface Representations of Mixed-Forests
3	G. Bohrer ¹ , J. E. C. Missik ¹
4	¹ Ohio State University, Department of Civil, Environmental & Geodetic Engineering.
5	
6	Corresponding author: Gil Bohrer (bohrer.17@osu.edu)
7	
8	Key Points:
9 10	• New generation of plant-hydrodynamic land surface models require resolution representative of characteristic individual trees.
11 12	• Tree-level flux representation diverges from the common area-based representation, especially in mixed plots of multiple species or types.
13 14 15	• We provide a formulation using common crown characteristics for consistent scaling of leaf- and tree-level fluxes to area-based plot level.

16 Abstract

The plant hydrodynamic approach represents a recent advancement to land surface modeling, in 17 which stomatal conductance responds to water availability in the xylem rather than in the soil. To 18 provide a realistic representation of tree hydrodynamics, hydrodynamic models must resolve 19 processes at the level of a single modelled tree, and then scale the resulting fluxes to the canopy 20 21 and land surface. While this tree-to-canopy scaling is trivial in a homogeneous canopy, mixedspecies canopies require careful representation of the species properties and a scaling approach 22 that results in a realistic description of both the canopy and individual-tree hydrodynamics, as 23 well as leaf-level fluxes from the canopy and their forcing. Here, we outline advantages and 24 pitfalls of three commonly used approaches for representing mixed-species forests in land 25 surface models, and present a new framework for scaling vegetation characteristics and fluxes in 26 27 mixed-species forests. The new formulation scales fluxes from the tree- to canopy-level in an

energy- and mass-conservative way and allows for a consistent multi-species canopy description

- 29 for hydrodynamic models.
- 30

Introduction – why tree-level fluxes are inconsistent with current surface flux representation

Representation of canopy and tree function in earth system models (ESMs) has gone 33 through rapid advancements in the last decade. The classical big-leaf representation of the forest 34 canopy [Farquhar et al., 1989; Sellers et al., 1992] in land surface models (LSMs) is being 35 expanded and replaced by more complex canopy representations. The complexity of canopy 36 representation in LSMs is advancing in multiple dimensions. One dimension of canopy 37 38 complexity is functional diversity. Models vary in the number and exact definitions of functional types they can represent. Nonetheless, almost every modern LSM can handle the representation 39 of structural/functional diversity of the ecosystem through sets of pre-calibrated parameters 40 describing land-cover classes and plant functional types (PFTs), equivalent to or more detailed 41 than the International Geosphere–Biosphere Programme (IGBP) land-cover classification system 42 [Loveland and Belward, 1997]. More complex canopy models can represent age/size stages 43 within functional types [see review by Fisher et al., 2018]. A second dimension of complexity 44 includes the spatial heterogeneity of the canopy, which can be represented by multiple "plots" 45 within a model grid-cell with different canopy height and leaf area characteristics. Finally, a third 46 dimension of complexity is the vertical structure of the canopy. Major advances have been made 47 in the representation of structural complexity of canopies along the vertical dimension, by 48 including vertically detailed representations of leaf density with between two canopy-leaf layers 49 [representing light and shade leaves, as in the CLM model, Dai et al., 2004] to n canopy-leaf 50 layers [for example in an advanced version of CLM, Bonan et al., 2021; Bonan et al., 2018]. 51 Advancements in remote sensing and ground-based optical techniques using high resolution lidar 52 and hyperspectral images make it possible to interpret tree-level information from remote 53 sensing data. These data can be used for resolving horizontal heterogeneity and tree-crown 54 characteristics [e.g., Detto et al., 2013; Garrity et al., 2012; Simard et al., 2011], 55 functional/species/trait identity and diversity [e.g., Schweiger et al., 2018; Serbin et al., 2014; 56 57 Sothe et al., 2019], and vertical canopy structure [e.g., Atkins et al., 2018; Lefsky et al., 2002; Yu *et al.*, 2017]. 58

The plant hydrodynamic approach represents an advancement to land surface modeling, 59 and LSMs and ESMs have recently started including a hydrodynamic version [Christoffersen et 60 al., 2016; Kennedy et al., 2019; Lawrence et al., 2019; Li et al., 2021; Xu et al., 2016]. Under the 61 hydrodynamic approach, stomata respond to water availability (represented as water content or 62 water potential) in the xylem, rather than directly responding to soil water availability. The result 63 is a model that includes the stem and/or stem xylem as potential spaces for water storage, and as 64 conduits for water. The stem and xylem characteristic dimensions (area, diameter, volume, etc.) 65 are needed by such models for determining the flow and storage of water in the stem/xylem. 66

Current, non-hydrodynamic LSMs represent vegetation at the site-level, corresponding to 67 the discretized spatial pixel of the atmospheric column model, and/or at the plot-level, 68 corresponding to the horizontally averaged representation of the vegetation of a particular land 69 cover or PFT. Different models use different terminology to the describe the smallest resolved 70 spatial unit (e.g., site, pixel, grid-cell), and the PFT-specific sub-grid scale unit (e.g., plot, tile, 71 patch). Throughout this manuscript we will use 'site' for a single resolved land unit and 'plot' for 72 the PFT-specific sub-grid-scale unit within a site. Consistent with the per ground area resolution 73 of fluxes and the atmospheric column, the surface and canopy characteristic dimensions are 74 either normalized per ground area (e.g., leaf area index, LAI) or provided in units of length (e.g., 75 rooting depth, roughness length, canopy height). Hydrodynamic models require a revised 76 approach to discretization of model canopies to the tree level, a distinctly different scale than the 77 site or plot levels. 78

79 As demonstrated by Bohrer et al. [2005], tree hydrodynamics are strongly dependent on the allometry of tree height, stem diameter, xylem area, and taper function (i.e., how the stem 80 diameter/area change with height). Furthermore, the stem xylem acts as a combined water 81 storage space and a central conduit to water transport from the soil, which serves the entire 82 crown. Therefore, for the purpose of resolving realistic tree hydrodynamics, it is imperative to 83 preserve the relationships between the total leaf area supported by a single crown, the height of 84 85 the crown, the xylem area and volume, and the vertical profiles of xylem area and leaf area throughout the crown, which requires a set of intrinsic characteristic length scales describing an 86 individual tree (Fig. 1). For an individual tree, a crown with a large ground-projection area is 87 88 supported by a stem with a conductive active xylem cross-sectional area. The stem and xylem 89 cross-section areas are much smaller than the crown projection area. Like the crown, the roots explore a certain volume with a horizontal projection area that is much larger than the stem. A 90 91 challenge for all hydrodynamic approaches is that diameter, height, area, and volume do not scale linearly [see example for allometric scaling functions relating height, crown and DBH, 92 93 Garrity et al., 2012]. Furthermore, the equations that describe the hydraulic relationships between water content and conductivity to water flow are strongly non-linear and depend both on 94 the volume and the conductive area of a stem element. For example, the combined crown area of 95 five small trees with stems with a diameter at breast height, DBH (the common measure of stem 96 97 size) of 10 [cm DBH], and a corresponding basal cross-section area of 314.16 [cm²] could be much larger (or could be smaller) than the crown area of a single tree with stem diameter of 50 98 cm (corresponding with basal area of 7853.98 $[cm^2]$), or a stem cross-section area of 5×314.16 = 99 1570.80 [cm²] (corresponding with a DBH of 22.36). Furthermore, for a particular tree species, 100 the characteristic height of a tree with a DBH of 10 [cm] is different than that of a 50 [cm DBH]-101 and 22.36 [cm DBH]-trees. There are infinite permutations of trees with different characteristic 102 sizes but the same leaf area index (m² leaves per m² ground area). Because both the conductive 103 area (for transpiration and sap flow) and the hydrostatic water potential in the xylem (a function 104

105 of volumetric water content) are important to the dynamics of moving water through the

106 conductive system of the tree, and because crown area, leaf area, and height are important to the

strength of evaporative demands, adding x% more xylem volume by growing x% taller and

- maintaining a constant stem and crown cross-sectional areas will have very different
- 109 consequences than adding the same x% more crown and stem-xylem areas while maintaining 110 constant height.

A trivial, intuitive scaling from the tree to the plot level involves multiplying the tree-111 level (total per tree) flows by the stand density (number of trees per unit plot area). However, 112 when scaling up from the tree to the canopy scale, there is no guarantee that the tree crowns 113 continuously and uniformly cover the whole plot area. In sparse forests, the average distance 114 between trees may be larger than the crown diameter, whereas in dense forests, individual tree 115 crowns may overlap and the mean distance between trees may be smaller than the crown 116 117 diameter. This leads to a discrepancy between the meaning of LAI in hydrodynamic models (at the tree scale) and general land surface models (at the plot scale). Hydrodynamic models must 118 explicitly represent all the transpiration from the entire crown, as all transpiration from a crown 119 flows through a single stem. That means that evaporative demands and transpiration rates are 120 considered per the area relevant to tree, i.e., the ground area projected under the crown, rather 121 than the ground area of the whole plot (Fig. 1). As the total area under the crowns (crown area \times 122 123 number of trees in the site) does not necessarily equal the plot area, special considerations must be given to scaling LAI and tree-level fluxes, which despite having the same physical units, in a 124 hydrodynamic tree-level model are provided per ground area under the crown, whereas the land 125 surface model requires fluxes per ground area in the plot. 126

127 The overall plot-level leaf area index (LAI_p) is currently the most observed and 128 commonly used site-level characteristic of vegetation canopies. Worldwide observations of LAI_p 129 exist from remote sensing products and plot-level optical measurements. LAI_p represents the 130 average leaf area per ground area for an entire plot [m² leaf/m² ground]. In a mixed forest it 131 includes leaves of multiple species (sp), such that

132

$$LAI_p = \sum_{sp=1}^{N_{sp}} LAI_p^{(sp)} \tag{1}$$

where $LAI_p^{(sp)}$ [m² leaf of (sp) /m² ground] is the species-specific leaf area index per unit 133 ground area in the plot and N_{sp} is the number of species in the plot. $LAI_n^{(sp)}$ is typically measured 134 using leaf-litter traps and could also be characterized by remote sensing using high-resolution 135 multi-spectral images combined with spectral characterization of different tree types [Ju and 136 Bohrer, 2022; Sothe et al., 2019; Yu et al., 2017]. Currently, $LAI_p^{(sp)}$ is available in some plots but 137 is not a very commonly reported property of forest plots. Splitting $LAI_p^{(sp)}$ to further distinguish 138 among size categories is harder and not commonly done. Nonetheless, any modeling approach 139 that resolves two different species requires $LAI_p^{(sp)}$, and therefore current modeling efforts 140 already must make some assumptions about $LAI_p^{(sp)}$ values at any given plot. 141

142 A different approach to view species-specific leaf area index is the ground accumulated 143 leaf area per unit ground-projection under the tree crowns of each species, $LAI_c^{(sp)}$ [m² leaf of 144 (sp) / m² ground under (sp) crowns]. Importantly, this is the leaf area that is supported by 145 individual crowns, and therefore it is $LAI_c^{(sp)}$ that characterizes the virtual hydrodynamic model

tree crowns for each type. While it is possible to measure $LAI_c^{(sp)}$ in many individual crowns 146 using ground-based optical methods with a narrow upward cone, or by combining lidar and 147 multispectral images, it is rarely done or reported as plot characteristic. Nonetheless, we can use 148 the relationship between $LAI_c^{(sp)}$ and $LAI_p^{(sp)}$ to determine a plot-characteristic crown-level 149 $LAI_{c}^{(sp)}$: 150

151
$$LAI_{p}^{(sp)} = LAI_{c}^{(sp)} \times A_{c}^{(sp)} \times 0.0001SD^{(sp)}$$
(2)

where $A_c^{(sp)}[m^2]$ is the characteristic ground-projection area of crowns of (sp) (Fig. 1), $SD^{(sp)}$ [ha⁻¹] is the species-specific stand density (i.e. the number of tree crowns of (sp) per unit 152

153 plot area in hectares), and 0.0001 is the conversion factor from hectare⁻¹ to m⁻². 154

155



156

Figure 1. Schematic illustration of the characteristic length scales and areas for species-specific 157 158 crowns and whole-plot descriptions of canopies.

159

The difference between the tree-level description of the hydrodynamic canopy and the 160

- horizontally averaged description of the canopy in the plot-scale surface-flux and radiation-161
- exchange modules leads to further specific challenges in mixed canopies that are composed of 162 trees belonging to more than one species, or trees of the same species that belong to different size 163

- classes, and therefore have different hydrodynamics. Imagine a mixed canopy, checkerboard 164
- pattern of canopy-dominant trees: 50% of trees of species A and 50% of species B, with an LAIp 165
- of 4. Assume species A and species B trees have the same crown-level leaf area index $LAI_c^{(sp)}$ 166
- (i.e., $LAI_c^{(A)} = LAI_c^{(B)} = 4$ [m² leaves of (sp) / m² ground under (sp) crowns]). The plot-scale, 167
- species-specific leaf area index $LAI_p^{(sp)}$ of both species A and B is 2 [m² leaves of (sp) / (m² 168
- ground)]. Imagine that species A trees are taller [10 m] than species B [8 m] and species A have 169
- thicker stems [DBH_A=20 cm] than species B [DBH_B=10 cm]. This hypothetical setup is 170
- 171 illustrated in Figure 2 (top panel).



172

- Figure 2. The "checkerboard-forest" conundrum how to represent the canopy and crowns 173
- characteristics of a mixed forest site composed of two species of trees with different crown 174
- characteristics and hydraulic traits (top panel), in crown-level hydraulic models that need to 175 represent the combination of crowns that make up the entire site? Three approaches, namely the
- 176 averaged-plot approach, mixed-plot approach, and multi-plot approach, which are currently
- 177
- utilized in models, are represented in the bottom panels. 178
- 179

How could such a mixed forest (Fig. 2, top panel) be represented in hydrodynamic 180 model? Current models take one of three approaches to solve this: (1) The averaged-plot 181 approach resolves a single plot with a single characteristic tree type that represents the average of 182 the two species in the original canopy (Fig. 2, bottom left). CLM-ml [Bonan et al., 2018], CLM5 183 [Song et al., 2020], and ELMv0 [Liang et al., 2019] are examples of this approach. (2) The 184 mixed-plot approach resolves two tree species co-filling the same plot, each with the assumed 185 plot-scale properties of the tree species (Fig. 2, bottom center). ED2 [Xu et al., 2016] and CLM-186 FATES [Koven et al., 2020] are examples of this approach. (3) The multi-plot approach resolves 187 a separate sub-site plot for each species, such that each plot is filled with a canopy composed of 188 trees of only one species. Each sub-site plot occupies half of the area of the site (Fig. 2, bottom 189 right). Most land surface models allow a multi-plot setup (as in approach 3) to represent different 190 PFTs. As an example of this approach, Cai et al. [2019] demonstrates a multi-plot application of 191 ELMv1 (the same model which Liang et al. 2019, referenced above, used in a single plot setup). 192 Each of these approaches result in some bias due to the assumptions of scaling from tree to site 193 scale. In the averaged-plot approach (1), the characteristics of the hydrodynamic model tree are 194 not realistic and match none of the two species. In the mixed-plot approach (2), assuming that the 195 number of stems is determined by a plot-level stand density, the stems of each of the tree species 196 support half the leaf area that their crowns support in reality. The two species compete for light, 197 which in the example above results in the taller species A always shading species B. In some 198 199 cases, especially involving forest stands of mixed ages where the different species represent different age/size categories, this is a realistic description of the relationship between the two 200 species. However, in a forest where the dominant canopy trees are of mixed species and grow 201 side by side, both species are exposed to full sunlight, especially around noontime, despite 202 species B being shorter on average. The multi-plot approach (3) results in hydrodynamically 203 realistic trees; however, the environmental drivers of evaporative demand that each species 204 experiences (e.g., temperature, humidity, and aerodynamic conductance) are calculated based on 205 canopy characteristics particular to a single species rather than a realistic mixed canopy. The 206 multi-plot approach is realistic when a forest is composed of large clusters of similar types/sizes 207 of trees, as is often the case in silvicultural and heavily managed forests but is not a realistic 208 representation of a mixed-canopy forest. In reality, scalar profiles, aerodynamic resistance, and 209 the resulting wind profile inside a mixed canopy are horizontally mixed at spatial scales smaller 210 than the turbulence eddy correlation length scale, and thus are characteristic of the mixed 211 canopy, not a "pure" canopy of species-A or species-B crowns. In addition, the multi-plot 212 approach also does not provide a realistic representation of water movement and water 213 availability in the soil since these are influenced by the combined root characteristics of the 214 different species in a mixed-canopy forest. 215

We propose a fourth approach that combines the advantages of the three approaches above. We will present a formulation that scales vegetation characteristics and the resulting fluxes from trees to forests in an energy and mass-conservative way and allows for a smooth and consistent multi-species canopy description for hydrodynamic models in mixed-forests.

220

221 2 Materials and Methods

222

2.1 Formulation - Canopy construction

We present the formulation for scaling hydrodynamic simulations of N_{sp} species of trees 223 in a mixed forest. For simplicity, we refer to the categories of trees representing distinct 224 hydraulic traits in the model as "species", but these could also represent different PFTs and/or 225 different size categories within a PFT. These N_{sp} species co-occupy a single plot in a mixed 226 227 canopy with assumed no (or low degree of) clustering among trees of the same species. In the model representation, characteristic tree-crown level dimensions will be used to specify N_{sp} 228 individual virtual trees, each representing one species (sp). Each of these individual virtual 229 crowns is resolved with multiple vertical layers, and for numerical simplicity, we assume that an 230 231 identical vertical discretization of space is used for all species of trees in the canopy.

Within the model plot, the plant area density in the virtual tree crowns, a_{cz} [m² plant area / m³ unit volume inside the crown], is composed of two components: the leaves, l_{cz} , and the stem area density, b_{cz} (i.e., $a_{cz} = l_{cz} + b_{cz}$). Plant area density can be measured directly with 3-D detail from airborne, spaceborne, or ground-based lidar. To represent the full crowns of multiple species co-occupying the same plot, a_{cz} values are organized in an array, \mathbf{a}_{cz} , with one column per species. Each column, $\mathbf{a}_{cz}^{(sp)}$, represents one model characteristic tree, and each row of the full matrix \mathbf{a}_{cz} represents one vertical layer, z, throughout the vertical dimension of the canopy:

239
$$a_{cz_{i}}^{(sp)} \in \mathbf{a}_{cz}^{(sp)} = \begin{bmatrix} a_{z_{Nz}} \\ \vdots \\ a_{0} \end{bmatrix}^{(sp)} \in \mathbf{a}_{cz} = \begin{bmatrix} a_{cz_{Nz}}^{(1)} & \dots & a_{cz_{Nz}}^{(N_{sp})} \\ \vdots & \ddots & \vdots \\ a_{c0}^{(1)} & \dots & a_{c0}^{(N_{sp})} \end{bmatrix}$$
(3)

such that the full array of plant area densities \mathbf{a}_{cz} is composed of N_{sp} species-specific column 240 vectors describing the vertical profiles of plant area density of each species, $a_{cz}^{(sp)}$, which are 241 discretized to elements, $a_{cz_i}^{(sp)}$, in [m² leaf/m³], describing the plant area density at the height 242 above ground, z_i , of each vertical layer within the canopy for each crown type. The vertical 243 discretization of the canopy is conducted such that the vertical spacing of each grid layer is 244 constant, i.e., $z_{i+1} - z_i = \Delta z$. It is also uniform across all crown types; i.e., the number of 245 vertical layers, N_{zz} is such that the maximal height of the top vertical layer is at the height of the 246 tallest crown type, i.e., $z_{Nz} = max\{h^{(1)}: h^{(Nsp)}\}$, where $h^{(sp)}$ is the crown-top height (commonly 247 referred to as "canopy height") of each species. sp. 248

249 It is common to ignore the stem area density by assuming: $b_{cz} = 0$, at which point the plant area density profiles become identical to the leaf area density, $a_{cz} = l_{cz}$. For the purposes of 250 calculating drag and light attenuation, ignoring stem/branch density is appropriate. However, b_{cz} 251 is required for applications that use a plant hydrodynamic approach. Plant hydrodynamic models 252 resolve the water movement and water potential through the xylem and thus require the 253 conductive cross-section area of the xylem throughout the tree system. The species-specific 254 active xylem conductive cross-section area, $\mathbf{bx}_{cz}^{(sp)}$ [m²] is typically calculated using species-specific allometric equations of \mathbf{b}_{cz} [*Matheny et al.*, 2014], which, in turn, is usually calculated 255 256 using an allometric equation of $DBH^{(sp)}$, $A_c^{(sp)}$, and z. For deciduous species, \mathbf{b}_{cz} can be directly 257 observed by lidar measurements in the dormant season. 258

In mixed forests, lidar can be used to observe the plot-level leaf area density profile l_{pz} 259 but, because it is impossible to distinguish species within a lidar point cloud, it cannot be used to 260 directly observe $\mathbf{l}_{cz}^{(sp)}$. Ground-based lidar observations conducted under individual crowns are 261 sometimes available, but typically an assumption must be made about the typical crown shape of 262 each species. This assumption will be represented as a unit-normalized vertical profile of leaf 263 density (i.e., when integrated vertically will sum to 1) and can be de-normalized to represent any 264 tree of that type by multiplying the z axis of the normalized profile by $h^{(sp)}$ and multiplying the 265 x axis by $LAI_c^{(sp)}/h^{(sp)}$. Note that by definition, 266

267
$$LAI_{c}^{(sp)} = \int_{0}^{z_{Nz}} l_{cz}^{(sp)} dz \simeq \Delta z \sum_{i=0}^{Nz} l_{cz_{i}}^{(sp)}$$
(4)

This approach can be used to combine plot-level observations of $LAI_c^{(sp)}$ and assumed allometric shapes of the crowns for each *sp* into a complete \mathbf{l}_{cz} array, which is also consistent across scales from the resolved vertical layer to the whole crown. Furthermore, a plot-level leaf area density profile that represents the mean canopy characteristic leaf area per ground are over all tree types, \mathbf{l}_{pz} (single column vector), can be formed from the normalized crown-level array, $\mathbf{l}_{cz,norm}$, and crown characteristics as:

274
$$\mathbf{l}_{pz} = \mathbf{l}_{cz}, norm \times \begin{bmatrix} \left(LAI_{c}^{(1)} \times A_{c}^{(1)} \times 0.0001SD^{(1)} \right) \\ \vdots \\ \left(LAI_{c}^{(Nsp)} \times A_{c}^{(Nsp)} \times 0.0001SD^{(Nsp)} \right) \end{bmatrix}$$
(5)

Where $l_{cz,norm}$ provided the vertical distribution of leaf area from the ground to the crown top and integrates to 1.

A similar scaling approach is to be used independently to derive the plot-level, whole 277 canopy \mathbf{b}_{pz} (if needed and not assumed = 0) and then $\mathbf{a}_{pz} = \mathbf{l}_{pz} + \mathbf{b}_{pz}$. It is expected that \mathbf{a}_{pz} 278 and \mathbf{l}_{pz} formed in this bottom-up approach by combining and scaling the individual species-279 specific profiles may diverge to some degree from directly observed full plot-level mean vertical 280 profiles of leaf area distribution. This is because the sources of error associated with the species-281 specific estimates of plant and leaf area densities are different than those in the direct plot-level 282 lidar measurements. Upon implementation, modelers should consider whether species-specific 283 crown-level or whole-plot-level data are more important to preserve or could make slight 284 modifications to each to optimize overall convergence with all scales of observations. 285

286 2.2 Multi-scale application – flux drivers

The purpose of any hydrodynamic module within a land surface model is to predict the 287 sensible heat, latent heat, and carbon fluxes of the land surface, incorporating the effects of tree 288 hydrodynamics on stomatal conductance. As such, any hydrodynamic module resolves a coupled 289 set of functions for each surface flux. Each of the functions for surface fluxes requires multiple 290 291 variables as drivers. These variables, specifically, the vertical profiles of net radiation, R_{nz} , air temperature, T_{cz} , humidity, q_{cz} , and wind speed U_{cz} , provide the values of the canopy 292 environment that dictate the evaporative demand. Most models include a virtual entity for 293 canopy air, embedded as a sub-grid-scale value (or set of values) within the lowest resolved 294 atmospheric grid cell. The conditions in the canopy air are diagnosed (or prognosed, in models 295 that resolve a fully coupled numerical scheme that link the surface fluxes and canopy air 296

conditions) as a function of the canopy characteristics (leaf density, roughness length) and the
 conditions in the open air above the canopy. Canopy air may be represented by a single bulk set
 of conditions or be vertically detailed.

Here, we discuss hydrodynamic models which resolve multiple species-specific, 300 vertically resolved crowns, representative of a mixed canopy. Each of the species-specific 301 crowns include a canopy air column with potentially different conditions. All the vertical canopy 302 columns interact with a single, common, site-level atmospheric column. However, it is not 303 obvious which canopy characteristics and at what scale best capture the relationship between the 304 conditions in the open air above the canopy and the values of the vertical profiles of the flux 305 drivers in the canopy air. For example, when resolving the specific-specific, crown-level R_{nz} at 306 midday in the checkerboard mixed forest (Figure 2), the multi-plot approach which represents 307 separate species-specific canopies will provide more realistic light attenuation profiles for each 308 309 species than the mixed-plot approach, which will result in excessive shading over all trees of species B. However, wind speed, temperature, and humidity profiles are mixed inside the canopy 310 at a length scale that is related to the eddy correlation scale and is typically larger than the size of 311 individual tree crowns. Therefore, for T_{az} , humidity, q_{cz} , and wind speed U_{cz} , the mixed-plot and 312 averaged-plot approaches will provide more realistic profiles than the multi-plot approach. 313

We propose a mixed approach, where vertical profiles of flux drivers are derived from 314 canopy representations of different scales. Specifically, radiation should be characterized at the 315 species-specific crown scale, i.e., $R_{nz}^{(sp)} = f(l_{cz}^{(sp)})$. Some unresolved issues for radiative 316 transfer formulation are the reflection and long-wave emissions from the surrounding trees, 317 which in a mixed forest do not have to originate from crowns of the same species and shading 318 from near-by crowns when the sun is far from zenith (e.g., early in the morning). There are many 319 different approaches and degrees of complexity to model within-canopy radiation conditions [see 320 review of models participating in the RAMI intercoparisson project, *Widlowski et al.*, 2015; 321 Widlowski et al., 2011]. Some models, such as ACTS [Ni-Meister et al., 2010; Yang et al., 2010], 322 323 are particularly suitable for hydrodynamic applications as they utilize a crown-based clumped approach and apply a crown-to-canopy scaling that utilizes crown diameter and stand density. 324

Flux drivers that are controlled by atmospheric surface layer mixing at typical length scales larger than a single tree crown (e.g, temperature, humidity, wind speed, and turbulent mixing) are characterized by the mean, whole-site canopy, e.g., $T_{cz}^{(sp)} = f(l_{pz})$. Wind speed and turbulence mixing profiles through a canopy of prescribed vegetation density can be determined using 1st or 2nd order closure approaches [*Massman*, 1997; *Massman and Weil*, 1999]. One approach to predict within-canopy temperature and humidity utilizes the vorticity-thickness length scale [*Harman and Finnigan*, 2007; 2008].

332 Scaling of fluxes from crowns to forests

Any hydrodynamic model must represent a discrete virtual individual tree crown. Consequently, fluxes from the leaves to the air, whether prescribed per leaf area or per ground area, must be scaled over the leaf area of the crown to represent the fluxes that are supported by an individual stem. To illustrate that, imagine a tree with a stem diameter of 30 cm supporting a crown with a diameter of 10 m, and a given transpiration rate per leaf area. Compare this tree to a tree with a stem of the same diameter and same $LAI_c^{(sp)}$, but growing in a sparser canopy, such that the crown area is 1.5 times larger than that of the first tree. As the crown area, and with it the

- total leaf area that are supported by the stem are larger, the sap flux (water per unit active xylem
- area) should be 1.5x larger, assuming xylem conductivity did not change. To preserve the
- conservation of mass through transport from the stem to the leaves and the air, hydrodynamic
- models must explicitly represent the crown by resolving flux from the leaves per leaf area, $F_{leaf}^{(sp)}$
- [kg m⁻² leaf s⁻¹], and multiply by the total leaf area of the crown $\left(LAI_c^{(sp)} \times A_c^{(sp)}\right)$ to get the
- total flow from a single crown, $F_{tree}^{(sp)}$ [kg s⁻¹].
- The same flow (with time lag due to changes of water storage in the crown) will be transported through the conductive area of a single stem and consumed from the soil by the corresponding root system of that single virtual tree. This tree-level flow can be represented as a flux per ground area under the crown, $F_c^{(sp)}$, where:

350
$$F_{c}^{(sp)} = \frac{F_{tree}^{(sp)}}{A_{c}^{(sp)}} = F_{leaf}^{(sp)} \times LAI_{c}^{(sp)}.$$
 (6)

F_c^(sp) will have the units of [x/s/m² projected ground area under the crown], where x corresponds to the units of the flux in question, i.e., Joules for energy fluxes, kg for mass fluxes, etc. This representation of transpiration in terms of $F_c^{(sp)}$ is convenient since $LAI_c^{(sp)}$ and $A_c^{(sp)}$ must already be known in order to calculate flux drivers that are specified at the crown scale (e.g., radiation). In its most general form, scaling from each species-specific flux to the whole-plot flux

to the atmosphere per unit ground area, F_p , is done as:

357
$$F_p = \sum_{sp} \Big(F_{tree}^{(sp)} \times 0.0001 SD^{(sp)} \Big).$$
(7)

358 When representing the tree-level flow in terms of $F_c^{(sp)}$, equation 7 becomes:

359
$$F_p = \sum_{sp} \Big(F_c^{(sp)} \times A_c^{(sp)} \times 0.0001 SD^{(sp)} \Big).$$
(7a)

Alternatively, from equation 2, the total leaf area of the crown could also be represented as $LAI_p^{(sp)}/0.0001SD^{(sp)}$, and $F_{tree}^{(sp)}$ could be represented as:

$$F_{tree}^{(sp)} = F_{leaf}^{(sp)} \times \frac{LAI_p^{(sp)}}{0.0001SD^{(sp)}}.$$
(8)

We note that although equation 8 does not require knowledge of $LAI_c^{(sp)}$ or $A_c^{(sp)}$, these

364 characteristics must be known in order to calculate crown-scale flux drivers. Thus, using either

365 equation 7a or equation 8 requires knowledge of the same canopy characteristics.

366 5 Conclusions

362

- 367 Canopy construction for hydrodynamic models requires species-specific and whole-plot
- 368 characteristics of the crowns and canopy. These include crown-level species-specific
- 369 characteristics of $LAI_c^{(sp)}$, $A_c^{(sp)}$, and $SD^{(Nsp)}$ (scalars), as well as the vertical profiles $\mathbf{a}_{cz}^{(sp)}$, $\mathbf{b}_{cz}^{(sp)}$
- and $\mathbf{l}_{cz}^{(sp)}$ (vectors), and the species-specific plot-level $LAI_p^{(sp)}$. From these characteristics the
- 371 whole plot LAI_p , \mathbf{a}_{pz} , and \mathbf{l}_{pz} can be calculated, though these may also be independently
- 372 observed. These crown, species, and whole-canopy characteristics provide a consistent multi-
- 373 scale description of the canopy and provide all the required canopy characteristics needed to
- 374 resolve and scale canopy fluxes.

375

376 Acknowledgments

We thank the support of the Ameriflux management project through core site US-UMB.

378

379 **References**

- Atkins, J. W., G. Bohrer, R. T. Fahey, B. S. Hardiman, T. H. Morin, A. E. L. Stovall, N.
 Zimmerman, and C. M. Gough (2018), Quantifying vegetation and canopy structural
 complexity from terrestrial LiDAR data using the FORESTR R package, *Methods in Ecology and Evolution*, 9(10), 2057-2066, https://doi.org/10.1111/2041-210X.13061.
- and Evolution, 9(10), 2057-2066, <u>https://doi.org/10.1111/2041-210X.13061</u>.
 Bohrer, G., H. Mourad, T. A. Laursen, D. Drewry, R. Avissar, D. Poggi, R. Oren, and G. G.
- Katul (2005), Finite-element tree crown hydrodynamics model (FETCH) using porous media
 flow within branching elements a new representation of tree hydrodynamics, *Water*
- 387 *Resources Research*, *41*(11), W11404, <u>https://doi.org/10.1029/2005WR004181</u>.
- Bonan, G. B., E. G. Patton, J. J. Finnigan, D. D. Baldocchi, and I. N. Harman (2021), Moving
 beyond the incorrect but useful paradigm: reevaluating big-leaf and multilayer plant canopies
 to model biosphere-atmosphere fluxes a review, *Agricultural and Forest Meteorology*, *306*,
 108435, <u>https://doi.org/10.1016/j.agrformet.2021.108435</u>.
- Bonan, G. B., E. G. Patton, I. N. Harman, K. W. Oleson, J. J. Finnigan, Y. Lu, and E. A.
 Burakowski (2018), Modeling canopy-induced turbulence in the Earth system: a unified
 parameterization of turbulent exchange within plant canopies and the roughness sublayer
 (CLM-ml v0), *Geoscientific Model Development*, 11(4), 1467-1496,
- 396 https://doi.org/10.5194/gmd-11-1467-2018.
- Cai, X., W. J. Riley, Q. Zhu, J. Tang, Z. Zeng, G. Bisht, and J. T. Randerson (2019), Improving
 representation of deforestation effects on evapotranspiration in the E3SM land model,
 Journal of Advances in Modeling Earth Systems, 11(8), 2412-2427, 10.1029/2018ms001551.
- Christoffersen, B. O., et al. (2016), Linking hydraulic traits to tropical forest function in a size structured and trait-driven model (TFS v.1-Hydro), *Geoscientific Model Development*, 9(11),
- 402 4227-4255, <u>https://doi.org/10.5194/gmd-9-4227-2016</u>.
- Dai, Y., R. E. Dickinson, and Y.-P. Wang (2004), A two-big-leaf model for canopy temperature,
 photosynthesis, and stomatal conductance, *Journal of Climate*, *17*(12), 2281-2299,
 https://doi.org/10.1175/1520-0442(2004)017<2281:atmfct>2.0.co;2.
- Detto, M., H. C. Muller-Landau, J. Mascaro, and G. P. Asner (2013), Hydrological networks and
 associated topographic variation as templates for the spatial organization of tropical forest
 vegetation, *PLoS One*, 8(10), e76296, https://doi.org/10.1371/journal.pone.0076296.
- Farquhar, G. D., D. A. Walker, and C. B. Osmond (1989), Models of integrated photosynthesis
 of cells and leaves, *Philosophical Transactions of the Royal Society of London. B, Biological Sciences*, 323(1216), 357-367, https://doi.org/10.1098/rstb.1989.0016.
- 412 Fisher, R. A., et al. (2018), Vegetation demographics in Earth System Models: A review of
- 413 progress and priorities, *Global Change Biology*, 24(1), 35-54,
- 414 <u>https://doi.org/10.1111/gcb.13910</u>.

- Garrity, S. R., K. Meyer, K. D. Maurer, B. S. Hardiman, and G. Bohrer (2012), Estimating plotlevel tree structure in a deciduous forest by combining allometric equations, spatial wavelet
 analysis and airborne lidar, *Remote Sensing Letters*, 3(5), 443–451,
- 418 <u>https://doi.org/10.1080/01431161.2011.618814</u>.
- Harman, I. N., and J. J. Finnigan (2007), A simple unified theory for flow in the canopy and
 roughness sublayer, *Boundary-Layer Meteorology*, *123*(2), 339-363, 10.1007/s10546-0069145-6.
- Harman, I. N., and J. J. Finnigan (2008), Scalar concentration profiles in the canopy and
 roughness sublayer, *Boundary-Layer Meteorology*, *129*(3), 323-351, 10.1007/s10546-0089328-4.
- Ju, Y., and G. Bohrer (2022), Classification of wetland begetation based on NDVI time series
 from the HLS dataset, *Remote Sensing Letters*, *14*, 2107, <u>https://doi.org/10.3390/rs14092107</u>.
- Kennedy, D., S. Swenson, K. W. Oleson, D. M. Lawrence, R. Fisher, A. C. L. d. Costa, and P.
 Gentine (2019), Implementing plant hydraulics in the Community Land Model, version 5, *Journal of Advances in Modeling Earth Systems*, *11*(2), 485-513, 10.1029/2018MS001500.
- Koven, C. D., et al. (2020), Benchmarking and parameter sensitivity of physiological and
 vegetation dynamics using the Functionally Assembled Terrestrial Ecosystem Simulator
- 431 Vegetation dynamics using the Functionally Assembled Terrestrial Ecosystem Simulato
 432 (FATES) at Barro Colorado Island, Panama, *Biogeosciences*, *17*(11), 3017-3044,
 433 https://doi.org/10.5194/bg-17-3017-2020.
- Lawrence, D. M., et al. (2019), The Community Land Model Version 5: Description of new
 features, benchmarking, and impact of forcing uncertainty, *Journal of Advances in Modeling Earth Systems*, 11(12), 4245-4287, https://doi.org/10.1029/2018MS001583.
- Lefsky, M. A., W. B. Cohen, G. G. Parker, and D. J. Harding (2002), Lidar remote sensing for
 ecosystem studies, *Bioscience*, 52(1), 19-30, <u>https://doi.org/10.1641/0006-</u>
 3568(2002)052[0019:LRSFES]2.0.CO;2.
- Li, L., Z.-L. Yang, A. M. Matheny, H. Zheng, S. C. Swenson, D. M. Lawrence, M. Barlage, B.
 Yan, N. G. McDowell, and L. R. Leung (2021), Representation of plant hydraulics in the
 Noah-MP land surface model: model development and multiscale evaluation, *Journal of*
- 443 Advances in Modeling Earth Systems, 13(4), e2020MS002214,
 444 https://doi.org/10.1029/2020MS002214.
- Liang, J., G. Wang, D. M. Ricciuto, L. Gu, P. J. Hanson, J. D. Wood, and M. A. Mayes (2019),
 Evaluating the E3SM land model version 0 (ELMv0) at a temperate forest site using flux and
 soil water measurements, *Geosci. Model Dev.*, *12*(4), 1601-1612,
- 448 https://doi.org/10.5194/gmd-12-1601-2019.
- Loveland, T. R., and A. S. Belward (1997), The IGBP-DIS global 1km land cover data set,
 DISCover: First results, *International Journal of Remote Sensing*, 18(15), 3289-3295,
 <u>https://doi.org/10.1080/014311697217099</u>.
- Massman, W. J. (1997), An analytical one-dimensional model of momentum transfer by
 vegetation of arbitrary structure, *Boundary-Layer Meteorology*, 83(3), 407-421.
- Massman, W. J., and J. C. Weil (1999), An analytical one-dimensional second-order closure
 model of turbulence statistics and the Lagrangian time scale within and above plant canopies
 of arbitrary structure, *Boundary-Layer Meteorology*, *91*(1), 81-107,
 10.1023/A:1001810204560.
- 458 Matheny, A. M., G. Bohrer, C. S. Vogel, T. H. Morin, L. He, G. Mirfenderesgi, K. V. R. Schäfer,
- 459 C. M. Gough, V. Y. Ivanov, and P. S. Curtis (2014), Species-specific transpiration responses

- to intermediate disturbance in a northern hardwood forest, *Journal of Geophysical Research-Biogeosciences*, *119*, 2292-2311, 10.1002/2014JG002804.
- Ni-Meister, W., W. Yang, and N. Y. Kiang (2010), A clumped-foliage canopy radiative transfer
 model for a global dynamic terrestrial ecosystem model. I: Theory, *Agricultural and Forest Meteorology*, *150*(7), 881-894, <u>https://doi.org/10.1016/j.agrformet.2010.02.009</u>.
- Schweiger, A. K., J. Cavender-Bares, P. A. Townsend, S. E. Hobbie, M. D. Madritch, R. Wang,
 D. Tilman, and J. A. Gamon (2018), Plant spectral diversity integrates functional and
 phylogenetic components of biodiversity and predicts ecosystem function, *Nature Ecology & Evolution*, 2(6), 976-982, https://doi.org/10.1038/s41559-018-0551-1.
- Sellers, P. J., J. A. Berry, G. J. Collatz, C. B. Field, and F. G. Hall (1992), Canopy reflectance,
 photosynthesis, and transpiration. III. A reanalysis using improved leaf models and a new
 canopy integration scheme, *Remote Sensing of Environment*, 42(3), 187-216,
 https://doi.org/10.1016/0034-4257(92)90102-P.
- 473 Serbin, S. P., A. Singh, B. E. McNeil, C. C. Kingdon, and P. A. Townsend (2014), Spectroscopic
 474 determination of leaf morphological and biochemical traits for northern temperate and boreal
 475 tree species, *Ecological Applications*, 24(7), 1651-1669, https://doi.org/10.1890/13-2110.1.
- Simard, M., N. Pinto, J. B. Fisher, and A. Baccini (2011), Mapping forest canopy height globally
 with spaceborne lidar, *Journal of Geophysical Research: Biogeosciences*, *116*(G4), G04021,
 https://doi.org/10.1029/2011JG001708.
- Song, J., G. R. Miller, A. T. Cahill, L. M. T. Aparecido, and G. W. Moore (2020), Modeling land
 surface processes over a mountainous rainforest in Costa Rica using CLM4.5 and CLM5, *Geoscientific Model Development*, *13*(11), 5147-5173, <u>https://doi.org/10.5194/gmd-13-5147-</u>
 2020.
- Sothe, C., M. Dalponte, C. M. d. Almeida, M. B. Schimalski, C. L. Lima, V. Liesenberg, G. T.
 Miyoshi, and A. M. G. Tommaselli (2019), Tree species classification in a highly diverse
 subtropical forest integrating UAV-based photogrammetric point cloud and hyperspectral
 data, *Remote Sensing*, 11(11), 1338, https://doi.org/10.3390/rs11111338.
- Widlowski, J., et al. (2015), The fourth phase of the radiative transfer model intercomparison
 (RAMI) exercise: Actual canopy scenarios and conformity testing, *Remote Sensing of Environment*, 169, 418-437, https://doi.org/10.1016/j.rse.2015.08.016.
- Widlowski, J., et al. (2011), RAMI4PILPS: An intercomparison of formulations for the
 partitioning of solar radiation in land surface models, *Journal of Geophysical Research: Biogeosciences*, *116*(G2), G02019, https://doi.org/10.1029/2010JG001511.
- Xu, X., D. Medvigy, J. S. Powers, J. M. Becknell, and K. Guan (2016), Diversity in plant
 hydraulic traits explains seasonal and inter-annual variations of vegetation dynamics in
 seasonally dry tropical forests, *New Phytologist*, 212(1), 80-95,
 https://doi.org/10.1111/nph.14009.
- Yang, W., W. Ni-Meister, N. Y. Kiang, P. R. Moorcroft, A. H. Strahler, and A. Oliphant (2010),
 A clumped-foliage canopy radiative transfer model for a Global Dynamic Terrestrial
 Ecosystem Model II: Comparison to measurements, *Agricultural and Forest Meteorology*,
- 500 *150*(7), 895-907, <u>https://doi.org/10.1016/j.agrformet.2010.02.008</u>.
- Yu, X., J. Hyyppä, P. Litkey, H. Kaartinen, M. Vastaranta, and M. Holopainen (2017), Single sensor solution to tree species classification using multispectral airborne laser scanning,
 Remote Sensing, 9(2), 108, https://doi.org/10.3390/rs9020108.
- 504