

The Greenhouse Effect – Re-examination of the Impact of an Increase in Carbon Dioxide in the Atmosphere

Trevor Underwood¹

¹Affiliation not available

November 30, 2022


Abstract

Examination of the radiation budget at the surface of the Earth shows that there are three factors affecting the surface temperature; the amount of solar radiation absorbed by the atmosphere and by the surface respectively, and the amount of leakage of infrared radiation emitted from the surface directly into space. If there were no leakage, the upwelling infrared radiation from the Earth's surface would be equal to the incoming solar radiation absorbed by the atmosphere plus twice the solar radiation absorbed by the surface. This results from the summation of a sequence of equal upward and downward re-emissions of infrared radiation absorbed by the atmosphere following the initial absorption of solar radiation. At current levels of solar absorption, this would result in total upwelling radiation of approximately 398.6 W/m², or a maximum surface temperature of 16.4°C. Allowing for leakage of infrared radiation through the atmospheric window, the resulting emission from the Earth's surface is reduced to around 396 W/m², corresponding to the current average global surface temperature of around 15.9°C. Absorption of solar and infrared radiation by greenhouse gases is determined by the absorption bands for the respective gases and their concentrations. Absorption of incoming solar radiation is largely by water vapor and ozone, and an increase in absorption would reduce not increase the surface temperature. Moreover, it is probable that all emitted infrared radiation that can be absorbed by greenhouse gases, primarily water vapor, with a small contribution from carbon dioxide and ozone, is already fully absorbed, and the leakage of around 5.5 % corresponds to the part of the infrared red spectrum that is not absorbed by greenhouse gases. The carbon dioxide absorption bands, which represent a very small percentage of the infrared spectrum, are most likely fully saturated. In these circumstances, increased concentrations of greenhouse gases, and carbon dioxide in particular, will have no effect on the emitted radiation. The surface temperature is probably at the thermodynamic limit for the current luminosity of the sun. Satellite based measurements since 1979 suggest that any global warming over the past 150 years may be due to an increase in total solar irradiance, which we are still a decade or two from being able to confirm.

The Greenhouse Effect – Re-examination of the Impact of an Increase in Carbon Dioxide in the Atmosphere[†]

[†] Extract from Underwood (2018). *The Greenhouse Effect – re-examination of the impact of an increase in carbon dioxide in the atmosphere. Unpublished.*

Examination of the radiation budget at the surface of the Earth shows that there are five primary factors affecting the surface temperature; the amount of solar radiation absorbed by the atmosphere and by the surface respectively, the amount of heat emitted from the surface in the form of thermals and evaporation, and the proportion of infrared radiation emitted from the surface directly into space. The Greenhouse Effect equations are solved by calculating the downwelling flux from the atmosphere and substituting this in the equation for the radiative balance at the Earth's surface.



The global annual mean earth's energy budget for 2000–2005 (W m^{-2}). The broad arrows indicate the schematic flow of energy in proportion to their importance. Adapted from Trenberth et al. (2009) (Fig. 1, Trenberth and Fasullo 2012).

Solution of the Greenhouse Effect equations

The Greenhouse Effect equations can be solved for the infrared flux emitted by the Earth's surface, E_T , and consequently for the temperature at the surface, by calculating the downwelling

energy flux from the atmosphere, E_D , and substituting this in the equation for energy flux balance at the Earth's surface, $E_S = E_{TH} + E_{EV} + E_T - E_D$, where E_S is the solar energy absorbed by the surface, E_{TH} is the loss of energy from the surface through thermals, and E_{EV} is the loss of energy from the surface through evaporation.

The downwelling energy flux from the atmosphere, $E_D = E_{TH} + E_{EV} + E_{OA}$, where E_{TH} is the reversal of the loss of energy from the surface through thermals, E_{EV} is the reversal of the loss of energy from the surface through evaporation, and E_{OA} is the sum of the part of the outgoing terrestrial infrared red radiation absorbed by the atmosphere that is re-emitted downward towards the Earth's surface, in multiple absorptions and re-emissions by the atmosphere and the Earth's surface. $E_{OA} = R_a + R_a f/2 + R_a f^2/4 + \dots + R_a f^n/2^n = R_a + R_a r$, or $E_{OA} = (1 + r)R_a$, where $R_a = (E_A + fE_S)/2$ is the initial downward infrared flux from the atmosphere; f is the proportion of terrestrial infrared radiation absorbed by the atmosphere; and $r = f/2 + f^2/4 + \dots + f^n/2^n$.

This assumes that the heat resulting from thermals E_{TH} , and evaporation E_{EV} , is returned to the surface as this does not affect the quantum state of greenhouse gases, and that part of the outgoing terrestrial infrared radiation is absorbed by the atmosphere and, as with the solar radiation absorbed by the atmosphere, changes the quantum state of the greenhouse gas molecules. On re-emission, half of the absorbed radiation is radiated downward towards the Earth's surface, to be absorbed and re-emitted, and half radiated upward into the atmosphere. This is confirmed by Trenberth and Fasullo (2012)'s diagram of energy fluxes as the downwelling flux of 333 W/m^2 cannot be due to re-emitted radiation alone as this would result in a similar upward flux, which greatly exceeds the outgoing flux at the TOA.

$r = f/2 + f^2/4 + \dots + f^n/2^n$ is a self-similar geometric series that can be summed by multiplying by the common ratio and subtracting. Multiplying by $f/2$ gives $rf/2 = f^2/4 + \dots + f^n/2^n$, and subtracting from $r = f/2 + f^2/4 + \dots + f^n/2^n$, gives $r - rf/2 = f/2$, or $r = f/(2 - f)$. Recognizing that $E_E = (1 - f)E_T$, where E_E is the total infrared radiation emitted from the Earth's surface that escapes directly into space, $E_E = E_T - f E_T$, so $f E_T = E_T - E_E$, or $f = (E_T - E_E)/E_T$. Substituting global average values from the corrected Trenberth and Fasullo (2012) data in $f = (E_T - E_E)/E_T$, gives $f = (396 - 22)/396 = 0.9444$. With $f = 0.9444$, $r = f/(2 - f) = 0.9444/1.0556 = 0.89$, which is effectively reached after seven iterations: $r = 0.9444/2 + 0.9444^2/4 + 0.9444^3/8 + 0.9444^4/16 + 0.9444^5/32 + \dots + f^n/2^n$. Substituting $r = f/(2 - f)$ in $E_{OA} = (1 + r)R_a$, gives $E_{OA} = [1 + f/(2 - f)]R_a = [(2 - f + f)/(2 - f)]R_a$, so $E_{OA} = 2R_a/(2 - f)$, and substituting $R_a = (E_A + fE_S)/2$ in $E_{OA} = 2R_a/(2 - f)$, gives the outgoing infrared radiation emitted by the atmosphere,

$$E_{OA} = (E_A + fE_S)/(2 - f).$$

Substituting global average values from the corrected Trenberth and Fasullo (2012) data in $E_{OA} = (E_A + fE_S)/(2 - f)$, gives $E_{OA} = (78.2 + 0.9444 \times 160.2)/(2 - 0.9444) = (78.2 + 151.3)/1.0556$, so the global average outgoing infrared radiation emitted by the atmosphere, $E_{OA} = (78.2 +$

$151.3/1.0556 = 229.5/1.0556 = 217.4 \text{ W/m}^2$. As an equal amount of infrared radiation is emitted upward, this compares with the corrected Trenberth and Fasullo (2012) estimate of 216.4 W/m^2 emitted by the atmosphere upward into space.

Substituting $E_{OA} = (E_A + fE_S)/(2 - f)$ in $E_D = E_{TH} + E_{EV} + E_{OA}$, gives the downwelling energy flux from the atmosphere,

$$E_D = E_{TH} + E_{EV} + (E_A + fE_S)/(2 - f).$$

Substituting global average values from the corrected Trenberth and Fasullo (2012) data in $E_D = E_{TH} + E_{EV} + E_{OA}$, gives the total downwelling energy flux from the atmosphere $E_D = 17 + 80 + 217.4 = 314.4 \text{ W/m}^2$, compared with Trenberth and Fasullo (2012)'s estimated residual of 333 W/m^2 and Wild et al. (2015)'s estimation of 342 W/m^2 . This equation shows that the amount of the downwelling radiation, and consequently the infrared radiation from the surface, is highly dependent on the absorption of solar radiation by the atmosphere, and the existence of thermals and evaporation, in addition to the absorption of solar radiation by the Earth's surface.

Substituting $E_D = E_{TH} + E_{EV} + (E_A + fE_S)/(2 - f)$ into the equation for the energy flux balance at Earth's surface, $E_S = E_{TH} + E_{EV} + E_T - E_D$, and expressing in terms of E_T provides a solution of the Greenhouse Effect equations in terms of the radiation emitted by the Earth's surface. At energy flux equilibrium at the Earth's surface, $E_S = E_{TH} + E_{EV} + E_T - E_{TH} - E_{EV} - (E_A + fE_S)/(2 - f)$, or the radiation emitted by the Earth's surface,

$$E_T = E_S + (E_A + fE_S)/(2 - f),$$

which is independent of E_{TH} and E_{EV} .

Substituting global average values from the corrected Trenberth and Fasullo (2012) data, $E_T = 160.2 + (78.2 + 0.9444 \times 160.2)/(2 - 0.9444)$, or $E_T = 160.2 + (78.2 + 151.3)/1.0556$, or $E_T = 160.2 + 217.4$, or the radiation emitted by the Earth's surface, $E_T = 160.2 + 217.4 = 377.6 \text{ W/m}^2$, compared with Trenberth and Fasullo (2012)'s value of 396 W/m^2 and Wild et al. (2015)'s estimation of 398 W/m^2 .

Substituting $E_T = 377.6 \text{ W/m}^2$ in Planck's formulation based on quantum mechanics of the Stefan-Boltzmann Law, $E = \sigma T^4$, where $\sigma = 2\pi^5 k^4/15c^2 h^3 = 5.67 \times 10^{-8} \text{ Wm}^{-2} \text{ K}^{-4}$, or $T^4 = E/\sigma = 377.6 \times 10^8/5.67 = 66.60 \times 10^8$, and the average global surface temperature of the Earth, $T = 285.67^\circ\text{K} = 285.67 - 273.15 = 12.52^\circ\text{C}$, compared with 15.94°C based on Trenberth and Fasullo (2012)'s estimate for E_T of 396 W/m^2 .

It should be remembered that this is an estimate of the Greenhouse Effect, the effect of absorption of greenhouse gases in the atmosphere on the surface temperature, not a complete estimate of the surface temperature. In the absence of an atmosphere, but allowing for an average reflection (albedo) from the Earth's surface of 6.8% ($= 23.1/341.3$), the temperature of

the surface of the Earth would be given by $T^4 = E/\sigma = 340.3 \times (1 - 0.068) \times 10^8/5.67 = 55.936507 \times 10^8$, so $T = 273.48^\circ\text{K} = 273.48 - 273.15 = 0.3^\circ\text{C}$.

If $f = 1$, $E_T = E_S + (E_A + fE_S)/(2 - f)$ becomes $E_T = 2E_S + E_A$, independent of E_{TH} and E_{EV} . This states that with complete absorption by the atmosphere, the infrared flux emitted by the Earth's surface equals twice the solar radiation absorbed by the surface plus the solar radiation absorbed by the atmosphere, showing that the cycle of emissions and re-radiations effectively doubles the downward flux from the atmosphere. Substituting global average values from the corrected Trenberth and Fasullo (2012) data in $E_T = 2E_S + E_A$, gives $E_T = 2 \times 160.2 + 78.2 = 398.6 \text{ W/m}^2$. Applying the Stefan-Boltzmann Law, $T^4 = E/\sigma = 398.6 \times 10^8/5.67 = 70.30 \times 10^8$, this corresponds to a maximum surface temperature, $T = 289.56^\circ\text{K} = 289.56 - 273.15 = 16.41^\circ\text{C}$.

Substituting formulae for the fluxes in the Greenhouse Effect equation, $E_T = E_S + (E_A + fE_S)/(2 - f)$, gives $\sigma T^4 = (1 - \alpha_p - \alpha_a)S_0/4 + [\alpha_a S_0/4 + f(1 - \alpha_p - \alpha_a)S_0/4]/(2 - f)$, or $\sigma T^4 = [(2 - f)(1 - \alpha_p - \alpha_a)S_0/4 + \alpha_a S_0/4 + f(1 - \alpha_p - \alpha_a)S_0/4]/(2 - f)$, or $T^4 = [(2 - f)(1 - \alpha_p - \alpha_a) + \alpha_a + f(1 - \alpha_p - \alpha_a)]S_0/4(2 - f)\sigma$, or $T^4 = [2 - 2\alpha_p - 2\alpha_a - f(1 - \alpha_p - \alpha_a) + \alpha_a + f(1 - \alpha_p - \alpha_a)]S_0/4(2 - f)\sigma$, or the average global surface temperature of the Earth T is given by

$$T^4 = (2 - 2\alpha_p - \alpha_a)S_0/4(2 - f)\sigma$$

where α_p is the proportion of incoming solar radiation reflected back into space by the clouds and Earth's surface (the albedo); α_a is the proportion of incoming solar radiation absorbed by the atmosphere; and S_0 is the solar flux at the top of the atmosphere. From this it can be seen that the surface temperature is increased by a reduction in solar reflexivity or a reduction in solar atmospheric absorptivity as well as by an increase in the atmospheric infrared absorption coefficient; and, of course, is also increased by an increase in solar irradiance.

An increase in the infrared absorption coefficient, f , involves increasing the internal energy of the greenhouse gas molecule by changing its quantum state, which can only occur at particular wavelengths known as absorption bands. However, as seen above, f is currently around 0.9444, which suggests that at the current surface temperature all of the radiation within the emission bands is fully absorbed, and that the remaining 5.56 percent of the infrared emission represents radiation with wavelengths within the atmospheric window. If this is true, there can be no further increase in f , and no increase in the surface temperature with an increase in carbon dioxide, and any increase in surface temperature will depend on an increase in solar irradiance. After losing a critical instrument when NASA's Glory satellite failed to achieve orbit in 2011, it will now be another 20 years before we have enough data to be sure.