

# Managing type 1 diabetes during the COVID-19 pandemic.

Ines CHARRADA<sup>1</sup>, Cyrine Bennasrallah<sup>1</sup>, Manel Ben Fredj<sup>1</sup>, Nadia Slimene<sup>1</sup>, Wafa Dhouib<sup>1</sup>, Hela Abroug<sup>1</sup>, Imen Zemni<sup>1</sup>, Meriam Oumayma Beji<sup>1</sup>, Chawki Loussaief<sup>1</sup>, Ines Bouanène<sup>1</sup>, and Asma Belguith Sriha<sup>1</sup>

<sup>1</sup>Faculté de medecine de Monastir

September 30, 2020

## Abstract

We report 2 imported cases of COVID 19 in patients with type 1 diabetes. The viral shedding duration was 18 and 22 days in the two cases. Both patients had maintained correct blood glucose levels without ketoacidosis decompensation with regular medical follow-up by telephone.

## Hosted file

article COVID19.pdf available at <https://authorea.com/users/363163/articles/484028-managing-type-1-diabetes-during-the-covid-19-pandemic>

## Hosted file

table 1.pdf available at <https://authorea.com/users/363163/articles/484028-managing-type-1-diabetes-during-the-covid-19-pandemic>

## Hosted file

Table 2.pdf available at <https://authorea.com/users/363163/articles/484028-managing-type-1-diabetes-during-the-covid-19-pandemic>