Dislocation Of Both Interphalangeal Joints In The Same Finger – Clinical Image

Alexei Buruian¹, Tiago Roseiro¹, Claudia Vale¹, Andre Carvalho¹, Emanuel Seiça¹, Fabio Fernandes¹, Manuel Gameiro¹, and Antonio Mendes¹

¹Hospital Distrital da Figueira da Foz

November 4, 2020

Abstract

Although isolated proximal interphalangeal (PIP) and distal interphalangeal (DIP) dislocations are commonly seen in the emergency room, simultaneous dislocation is a rare occurrence.

Keywords: Orthopaedics, double, interphalangeal, dislocation

Key Clinical Message

Excelent results may be obtained after closed reduction, followed by a short period of immobilization and early finger motion in patients with simultaneous dislocation of both interphalangeal joints of a finger.

A 54-year-old female patient presented to the emergency room after a fall on an outstretched hand. On physical examination

Author Contributions:

Alexei Buruian: Article elaboration, review and submission

Tiago Roseiro, Claudia Vale, Andre Carvalho, Emanuel Seiça, Fabio Fernandes: drafting and critical review of the manuscript.

Manuel Gameiro: Senior orthopaedic surgeon responsible for patient treatment

Antonio Mendes: Orthopaedic surgeon tutor involved in article elaboration and review.

All authors: read and approved the final manuscript.

Acknowledgements: Carlos Pereira, head of Orthopaedics department at Hospital Distrital da Figueira da Foz.

This manuscript has not been published and is not under consideration for publication elsewhere.

We have no conflicts of interest or funding to disclose.

Our institution does not require ethical approval for case reports.

Written informed consent has been obtained.

References:

Ahmed Abdelaal, Tomos Edwards, and Sambandam Anand, Simultaneous dislocation of both the proximal and distal interphalangeal joints of a little finger. BMJ Case Rep. 2016; 2016; bcr2015213914.







