

# Identifying Healthcare Professionals' Fear of COVID-19 and Their Attitudes towards Being Vaccinated

Eda Ergin<sup>1</sup> and Semih AKKOYUN<sup>2</sup>

<sup>1</sup>Affiliation not available

<sup>2</sup>Balikesir Universitesi

June 14, 2021

## Abstract

**Objective:** This study was conducted to determine healthcare professionals' fear of COVID-19 in and their attitudes towards being vaccinated against COVID-19. **Method:** The sample of this descriptive study was composed of 302 healthcare professionals working in a university research and training hospital. The data were collected with the help of an online survey method created with the help of Google Forms using the "Form for Determining Healthcare Professionals' Attitudes towards Being Vaccinated" and "The Fear of COVID-19 Scale" between December 28, 2020 and January 13, 2021. **Results:** The mean total Fear of COVID-19 Scale score of the healthcare professionals participating in the study was found to be  $17.20 \pm 6.015$ . A significant relationship was found between the Fear of COVID-19 Scale total score and participating healthcare professionals' gender, status of education, status of flu vaccination and status of being diagnosed with COVID-19 ( $p < 0.005$ ). **Conclusion:** According to research results, it was found that healthcare professionals experienced moderate fear, almost half of them were undecided about the COVID-19 vaccine, their thoughts about vaccination were positive and they were mostly concerned about the side effects of the vaccine.

## ABSTRACT

**Objective:** This study was conducted to determine healthcare professionals' fear of COVID-19 in and their attitudes towards being vaccinated against COVID-19.

**Method:** The sample of this descriptive study was composed of 302 healthcare professionals working in a university research and training hospital. The data were collected with the help of an online survey method created with the help of Google Forms using the "Form for Determining Healthcare Professionals' Attitudes towards Being Vaccinated" and "The Fear of COVID-19 Scale" between December 28, 2020 and January 13, 2021.

**Results:** The mean total Fear of COVID-19 Scale score of the healthcare professionals participating in the study was found to be  $17.20 \pm 6.015$ . A significant relationship was found between the Fear of COVID-19 Scale total score and participating healthcare professionals' gender, status of education, status of flu vaccination and status of being diagnosed with COVID-19 ( $p < 0.005$ ).

**Conclusion:** According to research results, it was found that healthcare professionals experienced moderate fear, almost half of them were undecided about the COVID-19 vaccine, their thoughts about vaccination were positive and they were mostly concerned about the side effects of the vaccine.

**Keywords:** COVID-19, Vaccine, Healthcare Professional

**What's known**

The psychological and well-being of healthcare professionals are seriously affected by the increased workload due to the COVID-19 epidemic, having to undertake new duties in addition to their regular responsibilities' and the anxiety of their relatives or themselves being infected by the disease

Vaccine hesitancy and misinformation present substantial obstacles to achieving sufficient coverage and community immunity in many countries.

### **What's new**

This was the first study to evaluate healthcare professionals' fear of COVID-19 and their attitudes towards vaccination before the COVID-19 vaccination program started in Turkey

It was determined that almost half of the healthcare professionals were undecided regarding the coronavirus vaccine, they had positive views about vaccination and they were mostly concerned about the side effects of the vaccine.

Healthcare professionals experienced moderate fear and that fear of COVID-19 was significantly higher in female healthcare professionals, in healthcare professionals who had the influenza vaccine, and in healthcare professionals who were not diagnosed with COVID-19.

## **INTRODUCTION**

Coronavirus disease 2019 (COVID-19) first originated on December 31, 2019 in the city of Wuhan, located in the Hubei province of China. It was declared a global pandemic by the World Health Organization (WHO) on March 11, 2020 due to its rapid spread from the first case and its global effect on the whole world <sup>1</sup>. It is known that healthcare professionals were previously affected by Severe Acute Respiratory Syndrome (SARS) and Middle East Respiratory Syndrome (MERS) epidemics <sup>2</sup>. The psychological and well-being of healthcare professionals are seriously affected by the increased workload due to the COVID-19 epidemic, having to undertake new duties in addition to their regular responsibilities' and the anxiety of their relatives or themselves being infected by the disease<sup>3,4</sup>.

Reports sent to WHO declared that 22,073 healthcare professionals from 52 countries tested positive for COVID-19 as of 8 April 2020<sup>5</sup>. A meta-analysis study determined that 10% of 119,216 COVID-19 positive cases were healthcare professionals with a mortality rate of 0.3% <sup>6</sup>. In Turkey, 29,865 healthcare professionals were reported to be positive for COVID-19 and 52 died as a result of COVID-19 as of September 2, 2020, <sup>7</sup>. Lengthy working hours, exposure to psychological and physical violence and lack of protective equipment are known to make healthcare professionals vulnerable to infection by suppressing their immune systems <sup>1</sup>. Previous studies demonstrated that vaccines saved more lives compared to studies conducted in the field of health. The history of modern vaccination starts with the invention of smallpox vaccine in 1789<sup>8</sup>. Although work for vaccination and treatment for COVID-19 disease continues, an effective method has not been found yet. The targeted substances in vaccine studies were determined in relation to the following criteria: the vaccine will be used in people older than 60 years of age and with chronic diseases, the vaccine will not develop undesired immune responses and it will be suitable for storage<sup>9</sup>.

A study conducted with an overseas sample found that healthcare professionals would experience the same reservations against COVID-19 vaccine when it became available just like the reservations they experienced against seasonal influenza vaccine <sup>10</sup>. In Turkey, the COVID-19 vaccination program started on January 13, 2021 with the CoronaVac inactive vaccine produced in China by the company named Sinovac. Healthcare professionals, being in the high risk group, were included in the vaccination program in the first phase<sup>11</sup>. The literature offers limited studies on healthcare professionals' attitudes towards the COVID-19 vaccine. Therefore, this study aimed to determine healthcare professionals' fear of COVID-19 and their attitudes towards vaccination before the COVID-19 vaccination program started in Turkey.

## **METHOD**

This descriptive research was conducted on healthcare professionals over 18 to determine healthcare professionals' fear of COVID-19 and their attitudes towards vaccination in a university hospital between the

dates of December 28 2020- January 13, 2021, at a period when COVID-19 vaccination was not yet being implemented in Turkey. The questionnaire form prepared by using Google Forms and delivered online to participants was used to collect data. The reporting of this study followed the STROBE checklist.

The study universe was composed of 858 healthcare professionals working in the university hospital. Two of the 304 healthcare professionals who accepted to participate in the study were excluded because they did not meet the research criteria and hence, the study sample consisted of 302 healthcare professionals. Sample size was calculated as 305 persons with the help of Epi-Info Statcalc Program with 50% expected prevalence, 97% confidence limit, 5% deviation, however, only 302 persons were reached.

*The inclusion criteria were as follows:* (i) to be 18 years old or above, (ii) voluntarily participating in research, (iii) to be living in Turkey during the COVID-19 pandemic, and (iv) using a smartphone

*The exclusion criteria were as follows:* (i) having any psychological problems and (ii) refusing to participate in the research.

## Data Collection Tools

“Form for Determining Healthcare Professionals’ Attitudes towards Being Vaccinated” and “The Fear of COVID-19 Scale” were used to collect research data.

### Form for Determining Healthcare Professionals’ Attitudes towards Being Vaccinated

The questionnaire consists of a total of 16 multiple-choice questions asking about demographic factors including age, gender, marital status and educational status of healthcare professionals and the views of healthcare professionals on coronavirus.

### The Fear of COVID-19 Scale

The scale developed by Ahorsu et al. (2020) was adapted to Turkish by Bakioğlu et al. (2020). the scale consists of one dimension and 7 items<sup>12</sup>. There is no reverse item in the scale. Each statement in the scale has a Likert-type scoring ranging from 1(strongly disagree) to 5 (strongly agree). The total score obtained from all items of the scale reflects the level of fear of coronavirus (COVID-19) experienced by the individual. The scores that can be obtained from the scale range between 7 and 35. A high score indicates “high” level of COVID-19 fear. The Cronbach alpha internal consistency coefficient of the scale was found to be .88<sup>12</sup>. In this study, the Cronbach alpha internal consistency coefficient was found to be .87.

## Ethical Aspect of the Research

Written permission was obtained to conduct the study from the institution (93559075-900- E.10989) where the study was conducted and from İzmir Bakırçay University ethics committee on 11.12.2020 (E-93802310-050.01.04-10007). An application was made to the Ministry of Health General Directorate of Health Services Scientific Research Platform and the approval of the Ministry of Health (2020-12-30T01\_16\_45) was obtained. Before the participants filled in the data forms, they were provided with information about the study on the first page of the online link and they were asked to check the statement “I agree to participate in the study” if they volunteered to participate in the study.

## Data Analysis

SPSS (Statistical Package for Social Science) 21.0 package program was used to analyze the data. Numerical and percentage distribution were employed in data analysis. Since the data did not conform to normal distribution, Kruskal-Wallis Test was used and mean standard deviation values and median values were provided for comparison with independent variables.

## Findings

Of the healthcare professionals included in the study, 59.9% (n: 181) were female, 60.6% (n: 183) were single, 36.1% (n: 109) were high school graduates, 42.4% (n: 128) had a seniority of 4-10 years, 72.5% (n: 219) were working as nurses, 87.7% (n: 265) were not vaccinated against influenza, 91.9% (n: 275) were not vaccinated

against influenza the previous year and 55% reported that they or one of their relatives had COVID-19. The total mean Fear of COVID-19 Scale score was found to be  $17.20 \pm 6.015$  (Table 1).

A significant correlation was found between participating healthcare professionals' total Fear of COVID-19 Scale score and their gender, education status, their flu vaccination status and their COVID diagnosis ( $p < 0.005$ ). No significant relationship was found between the Fear of COVID-19 Scale total score and marital status, year of employment, seniority and previous year's flu vaccination status ( $p > 0.005$ ).

Analysis of participating healthcare professionals' views about the coronavirus vaccine showed that 47% (n: 142) opted for the following options "The discovery of the coronavirus vaccine reduces my fears" and "I may be willing to administer the coronavirus vaccine" while 43.4% (n: 131) expressed that they were undecided (Table 2).

Comparison of healthcare professionals' views on the coronavirus vaccine with the Fear of COVID-19 Scale mean scores pointed to a significant difference between the Fear of COVID-19 Scale mean scores and the following statement: "The discovery of the coronavirus vaccine alleviates my fears" and "I may be willing to administer the coronavirus vaccine" ( $p < 0.05$ ) (Table 2).

## DISCUSSION

This study examined healthcare professionals' attitudes towards vaccination and their fears in regards to COVID-19 in the period before the COVID-19 vaccination program was available in Turkey. It was found that healthcare professionals experienced moderate levels of COVID-19 fear and that fear of COVID-19 was significantly higher in women. This finding is consistent with the results of various studies reporting that women have higher levels of anxiety and risk perception<sup>10,13-15</sup>. In early May 2020, more than 150,000 healthcare professionals were infected with COVID-19, and more than 1,400 people died worldwide<sup>16</sup>. This is believed to be the most important cause of COVID-19 fear among healthcare professionals.

Influenza and COVID-19 have similar symptoms and more severe infections, complications or fatal outcomes are recorded when an individual has influenza and COVID-19 at the same time<sup>17</sup>. This study found a significant relationship between the fear of COVID-19 among healthcare professionals and the status of being vaccinated with influenza vaccine and it was concluded that those who received the influenza vaccine had higher COVID-19 fear. Another study determined that the disease progressed more severely in cases where an individual is exposed to COVID-19 virus at the same time he/she contracts the influenza virus. The same study emphasized that the immunization rate against influenza virus is 55-60% in individuals who have been vaccinated and underlined the importance of influenza vaccination in order not to be infected with both coronavirus and influenza virus at the same time<sup>18</sup>.

In their compilation of 2017 in regards to vaccination rates in health care professionals in Turkey, Özışık et al. reported that the rate of influenza vaccine varied between 12.3%-35.3%<sup>19</sup>. The increasing demand for influenza vaccine is likely due to the increased awareness of respiratory viral disease<sup>20</sup>. Identifying the importance of COVID-19 vaccination and its relationship with the influenza vaccine is believed to shed light on future vaccination campaigns. A significant relationship was found between the educational status of the healthcare professionals participating in the study and the Fear of COVID-19 Scale total score. A study conducted by Mertens et al. found that the fear of COVID-19 decreased when the education level increased<sup>10</sup>.

Understanding health professionals' reservations about vaccination has important implications for public health administrations during outbreaks. Due to the uncertainty of vaccine features such as efficacy, side effects and duration of immunity, 43.4% of healthcare professionals in this study reported that they were undecided about getting the COVID-19 vaccine. Low vaccine acceptance rates were reported in other countries as well (53.7% in Italy and 58.9% in France) which supports this finding<sup>21,22</sup>. Low COVID-19 vaccine acceptance rates (52.0%) were reported among healthcare professionals in Malta<sup>20</sup>. Two studies based on the early stages of the outbreak (February and March) among nurses in Hong Kong reported low COVID-19 acceptance rates (40.0% and 63.0%)<sup>15,23</sup>. Contrary to the findings in this study, a study examining the

views of physicians, nurses and normal individuals found that 78% of physicians, 61% of nurses and 75% of individuals stated that they wanted to be vaccinated against COVID-19 and it was determined that 60% of physicians, 55% of nurses and 70% of individuals in the general population wanted their children to be vaccinated<sup>24</sup>. It was concluded that healthcare professionals' reservations about vaccination was related to lack of sufficient information about the side effects of the vaccine and that increased the level of anxiety in 68.2% of healthcare professionals. Healthcare authorities can organize a vaccination program for healthcare professionals to vaccinate them against COVID-19. Future research is also needed to explore potential strategies for raising awareness and changing attitudes towards vaccination.

## Conclusion

Based on the findings of this research, it was concluded that healthcare professionals experienced moderate fear and that fear of COVID-19 was significantly higher in female healthcare professionals, in healthcare professionals who had the influenza vaccine, and in healthcare professionals who were not diagnosed with COVID-19. It was determined that almost half of the healthcare professionals were undecided regarding the coronavirus vaccine, they had positive views about vaccination and they were mostly concerned about the side effects of the vaccine.

To prevent the prevalence of COVID-19 vaccine hesitancy, it is suggested to build trust in COVID-19 vaccination among healthcare professionals and the public via timely and clear messages advocating for the safety and efficacy of COVID-19 vaccines through trusted channels.

## Limitations

There are some limitations to be considered for the present study. First of all, this study was conducted with healthcare professionals' working in one university hospital during the pandemic at a period when COVID-19 vaccination was not yet being implemented in Turkey. Therefore, the results from the present study cannot be generalized for other healthcare professionals'. Second, data collection forms are filled online, and we did not have the chance to control the data collection process.

## REFERENCES

1. WHO. Coronavirus disease Situation Report-200. 2020;(July). [https://bestpractice.bmj.com/topics/en-us/3000168/pdf/3000168/Coronavirus disease 2019 %28COVID-19%29.pdf](https://bestpractice.bmj.com/topics/en-us/3000168/pdf/3000168/Coronavirus%20disease%202019%20COVID-19%29.pdf).
2. Pala SÇ, Metintaş S. Covid-19 pandemisinde sağlık çalışanları. *Estud halk sağlığı Derg* . 2020;5:156-168.
3. David K, Meng KL, Sin EC, et al. Risk perception and impact of severe acute respiratory syndrome (SARS) on work and personal lives of healthcare workers in Singapore: What can we learn? *Med Care* . 2005;43(7):676-682. doi:10.1097/01.mlr.0000167181.36730.cc
4. Maunder RG, Lancee WJ, Rourke S, et al. Factors associated with the psychological impact of severe acute respiratory syndrome on nurses and other hospital workers in Toronto. *Psychosom Med* . 2004;66(6):938-942. doi:10.1097/01.psy.0000145673.84698.18
5. WHO. Coronavirus disease (COVID-19) Pandemic. World Health Organizations.
6. Sahu AK, Amrithanand VT, Mathew R, Aggarwal P, Nayer J, Bhoi S. COVID-19 in health care workers – A systematic review and meta-analysis. *Am J Emerg Med* . 2020;38(9):1727-1731. doi:10.1016/j.ajem.2020.05.113
7. Medimagazin. Republic of Turkey Ministry of Health Report. <https://www.medimagazin.com.tr/guncel/genel/tr-fahrettin-koca-ilk-kez-sayi-verdi-29-bin865-saglik-calisanlari-enfekte-11-681-91089.html>. Published 2020. Accessed September 3, 2021.
8. Akdeniz M, Kavucuk E. Aşılama ve Aşıların Tarihçesi. *Klin Tıp Aile Hekim* . 2016;8(2):11-28. <https://dergipark.org.tr/tr/pub/ktah/issue/45376/487032>.

9. Chen WH, Strych U, Hotez PJ, Bottazzi ME. The SARS-CoV-2 Vaccine Pipeline: an Overview. *Curr Trop Med Reports* . 2020;7(2):61-64. doi:10.1007/s40475-020-00201-6
10. Mertens G, Gerritsen L, Duijndam S, Saleminck E, Engelhard IM. Fear of the coronavirus (COVID-19): Predictors in an online study conducted in March 2020. *J Anxiety Disord* . 2020;74(June):102258. doi:10.1016/j.janxdis.2020.102258
11. Republic of Turkey Ministry of Health. REPUBLIC OF TURKEY MINISTRY OF HEALTH COVID-19 VACCINATION INFORMATION PLATFORM. <https://covid19asi.saglik.gov.tr/>. Published 2021. Accessed January 15, 2021.
12. Bakioglu F, Korkmaz O, Ercan H. Fear of COVID-19 and Positivity: Mediating Role of Intolerance of Uncertainty, Depression, Anxiety, and Stress. *Int J Ment Health Addict* . 2020. doi:10.1007/s11469-020-00331-y
13. Kong X, Zheng K, Tang M, et al. Prevalence and Factors Associated with Depression and Anxiety of Hospitalized Patients with COVID-19. *medRxiv* . 2020;21(1):1-9. doi:10.1101/2020.03.24.20043075
14. Kurt O, Oguzoncul AF. Levels of anxiety and depression related to COVID-19 among physicians: An online cross-sectional study from Turkey. *Ann Clin Anal Med* . 2020;11(suppl 3):1-6. doi:10.4328/acam.20206
15. Wang K, Wong ELY, Ho KF, et al. Intention of nurses to accept coronavirus disease 2019 vaccination and change of intention to accept seasonal influenza vaccination during the coronavirus disease 2019 pandemic: A cross-sectional survey. *Vaccine* . 2020;38(45):7049-7056. doi:10.1016/j.vaccine.2020.09.021
16. Bandyopadhyay S, Baticulon RE, Kadhum M, et al. Infection and mortality of healthcare workers worldwide from COVID-19: A systematic review. *BMJ Glob Heal* . 2020;5(12):1-11. doi:10.1136/bmjgh-2020-003097
17. Hashemi SA, Safamanesh S, Ghafouri M, et al. Co-infection with COVID-19 and influenza A virus in two died patients with acute respiratory syndrome, Bojnurd, Iran. *J Med Virol* . 2020;92(11):2319-2321. doi:10.1002/jmv.26014
18. Grech V, Borg M. Influenza vaccination in the COVID-19 era. *Early Hum Dev* . 2020;148. doi:10.1016/j.earlhumdev.2020.105116
19. Ozisik L, Tanriover MD, Altinel S, Unal S. Vaccinating healthcare workers: Level of implementation, barriers and proposal for evidence-based policies in Turkey. *Hum Vaccines Immunother* . 2017;13(5):1198-1206. doi:10.1080/21645515.2016.1269992
20. Grech V, Gauci C. Vaccine hesitancy in the University of Malta Faculties of Health Sciences, Dentistry and Medicine vis-à-vis influenza and novel COVID-19 vaccination. *Early Hum Dev* . 2020;(xxxx):105258. doi:10.1016/j.earlhumdev.2020.105258
21. La Vecchia C, Negri E, Alicandro G, Scarpino V. Attitudes towards influenza vaccine and a potential COVID-19 vaccine in Italy and differences across occupational groups, September 2020. *Med Lav* . 2020;111(6):445-448. doi:10.23749/mdl.v111i6.10813
22. Lazarus J V., Ratzan SC, Palayew A, et al. A global survey of potential acceptance of a COVID-19 vaccine. *Nat Med* . 2020;27(February):225-228. doi:10.1038/s41591-020-1124-9
23. Wang J, Jing R, Lai X, et al. Acceptance of covid-19 vaccination during the covid-19 pandemic in china. *Vaccines* . 2020;8(3):1-14. doi:10.3390/vaccines8030482
24. Dror AA, Eisenbach N, Taiber S, et al. Vaccine hesitancy: the next challenge in the fight against COVID-19. *Eur J Epidemiol* . 2020;35(8):775-779. doi:10.1007/s10654-020-00671-y

**Hosted file**

Table.docx available at <https://authorea.com/users/419787/articles/526162-identifying-healthcare-professionals-fear-of-covid-19-and-their-attitudes-towards-being-vaccinated>