

Correlated evolution of distinct signals associated with increased social selection in female white-shouldered fairywrens

John Jones¹, Karan Odom², Ian Hoppe³, Doka Nason⁴, Serena Ketaloya⁴, and Jordan Karubian¹

¹Tulane University

²University of Maryland at College Park

³University of Nebraska-Lincoln

⁴NA

September 24, 2021

Abstract

Conspicuous female signals have recently received substantial scientific attention, but the degree to which female-specific selection drives their evolution remains unclear. Species that express female-specific phenotypic variation among populations represent a useful opportunity to address this knowledge gap. White-shouldered fairywrens (*Malurus alboscapulatus*) are tropical songbirds with a well-resolved phylogeny where female, but not male, coloration varies allopatrically across subspecies. We explored how four distinct signaling modalities, each putatively associated with increased social selection, are expressed in two populations that vary in competitive pressure on females. Females in a derived subspecies (*M. a. moretoni*) have evolved more ornamented plumage, more complex vocalizations, and shorter tails (a signal of social dominance) relative to an ancestral subspecies (*M. a. lorentzi*) with drab females. Moreover, in response to simulated territorial intrusions broadcasting female song, female *M. a. moretoni* are more aggressive and more coordinated with their mate in both movement and vocalizations. These results suggest that correlated phenotypic shifts in female color, morphology, song complexity, and behavior may have occurred in response to changes in social selection, consistent with the idea that female-specific selection has driven the evolution of these signals.

Hosted file

Main text body - EcoEvo 8August.docx available at <https://authorea.com/users/423893/articles/538490-correlated-evolution-of-distinct-signals-associated-with-increased-social-selection-in-female-white-shouldered-fairywrens>