Hypoxemia indicative of Rendu Osler disease during SARS Cov-2 pandemic

Herveat RAMANANDAFY¹, Princy Parfait Andriamahenina ¹, Harison Michel Tiaray ¹, Anjara Mihaja Nandimbiniaina ¹, Angela Zamelina Razafindrasoa ¹, Sonia Marcelle Razafimpihanina ¹, Diamondra ombanjanahary Andriarimanga ¹, Solohery Jean Noel Ratsimbazafy ¹, Jocelyn Robert Rakotomizao ¹, Joëlson Lovaniaina Rakotoson ¹, and Hanta Marie Danielle Vololontiana²

¹University of Madagascar

²University of Antananarivo Faculty of Medicine

November 10, 2021

Abstract

Rendu Osler's disease is a genetic disease characterized by mucocutaneous and visceral telangiectasias. Rendu Osler's disease was discovered during hypoxemia during an outbreak of SARS-Cov2. This was a 36-year-old woman with exertional dyspnea and severe hypoxemia revealing pulmonary arteriovenous malformations on chest CT scan.

Hosted file

manuscrit revised.docx available at <https://authorea.com/users/445462/articles/544974-hypoxemia-indicative-of-rendu-osler-disease-during-sars-cov-2-pandemic>