

INTERLEUKIN 6 AS A MARKER OF SEVERE BACTERIAL INFECTION IN CHILDREN WITH SICKLE CELL DISEASE AND FEVER

Elena Rincon-Lopez¹, Maria Navarro Gómez¹, Teresa Hernández-Sampelayo Matos¹, David Aguilera-Alonso¹, Eva Dueñas-Moreno², Jesús Saavedra-Lozano¹, Begoña Santiago-García¹, Maria Del Mar Santos², Marina García-Morin¹, Cristina Beléndez¹, Jorge Lorente-Romero², and ELENA CELA¹

¹Hospital General Universitario Gregorio Marañón

²Hospital General Universitario Gregorio Marañón

March 30, 2022

Abstract

Introduction Etiological diagnosis of fever in sickle cell disease (SCD) children is often challenging. **Objective:** to analyze the pattern of inflammatory biomarkers in SCD febrile children and controls, in order to determine predictors of severe bacterial infection (SBI). **Methods** Prospective, case-control study of febrile and steady-state SCD children carried out during 3 years. Clinical characteristics and laboratory parameters, including 10 serum proinflammatory cytokines (IL-1 β , IL-2, IL-4, IL-6, IL-8, IL-10, IL-12p70, IL-17a, IFN- γ and TNF- α) and comparisons among study subgroups were analyzed. **Results** A total of 137 patients (78 cases and 59 controls) were included in the study; 78.5% males, median age 4.1 (1.7-7.5) years. Four cases were diagnosed with SBI, 41 viral infection (VI) and 33 no proven infection (NPI). IL-6 was significantly higher in patients with SBI than in patients with VI or NPI (163 vs 0.7 vs 0.7 pg/ml, $p < 0.001$), and undetectable in all controls. The rest of the cytokines analyzed did not show any significant difference. The optimal cut-off value of IL-6 for the diagnosis of SBI was 125 pg/mL, with high PPV and NPV (PPV of 100% for a prevalence of 5, 10 and 15% and NPV of 98.7%, 97.3% and 95.8% for those prevalences, respectively). **Conclusion** We found that IL-6 (optimal cut-off value of 125 pg/ml) was a very good marker for SBI in this cohort of febrile SCD children, with high PPV and NPV. Therefore, IL-6 may be useful, alone or combined with other biomarkers, to guide the management of these patients.

INTERLEUKIN 6 AS A MARKER OF SEVERE BACTERIAL INFECTION IN CHILDREN WITH SICKLE CELL DISEASE AND FEVER

Elena María Rincón-López^{1 2}, MD; María Luisa Navarro Gómez^{1 3}, MD, PhD; Teresa Hernández-Sampelayo Matos^{1 3}, MD, PhD; David Aguilera-Alonso¹, Eva Dueñas Moreno¹; MD; Jesús Saavedra-Lozano^{1 3} MD, PhD; Begoña Santiago García¹, MD, PhD; María del Mar Santos Sebastián¹, MD; Marina García Morín⁴, MD; Cristina Beléndez Bieler⁴, MD; Jorge Lorente Romero⁵, MD; Elena Cela de Julián^{3 4}, MD, PhD on behalf of F-DREP Study Group*.

*A list of other contributors of the F-DREP Study Group and their affiliations is provided in the Collaborators section.

Affiliations: ¹Pediatric Infectious Diseases Unit, Hospital General Universitario Gregorio Marañón, Instituto de Investigación Sanitaria Gregorio Marañón, Madrid, Spain. ²PhD Program in Medicine, Universidad Complutense de Madrid, Spain. ³Universidad Complutense de Madrid, Spain. ⁴Pediatric Hematology and

Oncology Unit, Hospital General Universitario Gregorio Marañón, Madrid, Spain. 5Pediatric Emergency Unit, Hospital General Universitario Gregorio Marañón, Madrid, Spain.

Address correspondence to: Elena M. Rincón-López. Servicio de Pediatría, Sección Enfermedades Infecciosas, Hospital Materno-Infantil Gregorio Marañón, c/ O'Donnell, 48-50, 28009 Madrid, Spain. Phone number: 0034 650060285. E-mail: elenarinconlopez@hotmail.com

Word count: Abstract: 250 words. Main text: 2738 words.

Figures and tables: 2 figures and 4 tables. 1 supplemental table.

Short title: IL-6 in sickle cell disease.

Key words: Sickle cell disease, infection, fever, cytokines, interleukin-6.

Data availability: The data that support the findings of this study are available on request from the corresponding author. The data are not publicly available due to privacy or ethical restrictions.

Abbreviations:

SCD	Sickle cell disease
SBI	Severe bacterial infection
VOC	Vaso-occlusive crisis
HGUGM	Hospital General Universitario Gregorio Marañón
CRP	C-reactive protein
Multiplex-PCR	Multiplex polymerase chain reaction
RSV	Respiratory syncytial virus
IL	Interleukin
IFN	Interferon
TNF	Tumor necrosis factor
UTI	Urinary tract infection
VI	Viral infection
NPI	No proven infection
CVC	Central venous catheter
ACS	Acute chest syndrome
PICU	Pediatric intensive care unit
IQR	Interquartile range
ROC	Receiver operating characteristic
AUC	Area under the ROC curve
PPV	Positive predictive value
NPV	Negative predictive value
PR	Prevalence rate
CI	Confidence interval

ABSTRACT

Introduction

Etiological diagnosis of fever in sickle cell disease (SCD) children is often challenging. Objective: to analyze the pattern of inflammatory biomarkers in SCD febrile children and controls, in order to determine predictors of severe bacterial infection (SBI).

Methods

Prospective, case-control study of febrile and steady-state SCD children carried out during 3 years. Clinical characteristics and laboratory parameters, including 10 serum proinflammatory cytokines (IL-1 β , IL-2, IL-

4, IL-6, IL-8, IL-10, IL-12p70, IL-17a, IFN- γ and TNF- α) and comparisons among study subgroups were analyzed.

Results

A total of 137 patients (78 cases and 59 controls) were included in the study; 78.5% males, median age 4.1 (1.7-7.5) years. Four cases were diagnosed with SBI, 41 viral infection (VI) and 33 no proven infection (NPI). IL-6 was significantly higher in patients with SBI than in patients with VI or NPI (163 vs 0.7 vs 0.7 pg/ml, $p < 0.001$), and undetectable in all controls. The rest of the cytokines analyzed did not show any significant difference. The optimal cut-off value of IL-6 for the diagnosis of SBI was 125 pg/mL, with high PPV and NPV (PPV of 100% for a prevalence of 5, 10 and 15% and NPV of 98.7%, 97.3% and 95.8% for those prevalences, respectively).

Conclusion

We found that IL-6 (optimal cut-off value of 125 pg/ml) was a very good marker for SBI in this cohort of febrile SCD children, with high PPV and NPV. Therefore, IL-6 may be useful, alone or combined with other biomarkers, to guide the management of these patients.

INTRODUCTION

Febrile episodes are common in children with sickle cell disease (SCD), often with challenging etiological diagnosis. These patients are at risk of severe bacterial infections (SBI), although they have become less frequent in high-income countries in recent years¹⁻⁵. Therefore, it would be important to find biomarkers that could help distinguish the etiology of these febrile episodes early in order to guide antimicrobial therapy. Several biomarkers [white blood count (WBC), neutrophils, C-reactive protein (CRP) and procalcitonin, among others] have been studied in SCD patients with bacteremia and other bacterial infections with variable results^{1,6-9}. Serum cytokines have also been studied in steady-state SCD patients and those with vaso-occlusive crisis (VOC)¹⁰⁻¹⁵, but they have not been well studied in SCD patients with an infection.

The aim of this study was to analyze the pattern of inflammatory biomarkers, including pro-inflammatory cytokines, in SCD children with fever of different etiologies, in order to find useful parameters to discriminate the patients at risk of SBI. Since SBI has become less frequent in well-controlled SCD children in recent years, this could help to guide the management of these patients, reducing unnecessary antibiotic treatments and hospital admissions.

MATERIAL AND METHODS

Study design

We conducted a prospective, case-control study with patients younger than 18 years with SCD from June 2015 to June 2018. Children with SCD and fever were eligible to participate in the study as cases and asymptomatic steady-state SCD children were included as controls. Exclusion criteria of cases and controls included age older than 18 years, hematopoietic stem cell transplantation, incomplete diagnostic tests and patients whose parents or legal guardians did not sign the informed consent form for the study. Controls were also excluded if they had had a febrile episode in the previous month. The study received approval from the Institutional Review Board.

Study setting

This study was conducted at the Hospital General Universitario Gregorio Marañón (HGUGM) in Madrid, Spain, a tertiary hospital with 120 pediatric beds and around 7000 pediatric inpatient admissions per year. HGUGM is a national reference center for children with SCD in Spain, with about 300 SCD pediatric patients upon follow-up.

Study protocol and definitions

Patients included in the study as cases were those with an axillary temperature of 38°C or more, documented at home or within the first 24 hours of hospitalization. Controls were asymptomatic steady-state SCD patients recruited during their follow-up attendance in the Pediatric Hematology clinic. Patients recruited as cases could also be included in the study as controls if at least one month had passed since their last fever episode. The following samples were collected in all cases: blood tests (including complete blood count, biochemistry, CRP and procalcitonin), blood cultures and nasopharyngeal samples for viral detection by a multiplex polymerase chain reaction (multiplex-PCR) assay [including influenza A, B and C, respiratory syncytial virus (RSV), human metapneumovirus, adenovirus, human bocavirus, rhinovirus, human parainfluenza virus and coronavirus]. Other diagnostic tests and the patients' management were performed under the criteria of the treating physicians. In all cases and controls a blood sample was collected and stored at the HGUGM Biobank for subsequent cytokine analysis. This analysis was performed by the DIAplex Human Th1 / Th2 / Inflammation kit (bioNova científica S. L., Spain), a multiplexed fluorescent bead-based immunoassay for the quantification of multiple human cytokines in serum and culture supernatants by flow cytometry, including the following: interleukin (IL)-1 β , IL-2, IL-4, IL-6, IL-8, IL-10, IL-12p70, IL-17a, interferon- γ (IFN- γ) and tumor necrosis factor- α (TNF- α). The limit of detection for the different cytokines was: IL-1 β 3.5 pg/mL, IL-2 12.4 pg/mL, IL-4 4.3 pg/mL, IL-6 1.4 pg/mL, IL-8 1.3 pg/mL, IL-10 1.7 pg/mL, IL-12p70 3.4 pg/mL, IL-17a 8.7 pg/mL, TNF α 9.8 pg/mL and IFN- γ 0.8 pg/mL. For statistical analysis, when a single value was non-detectable, it was considered to be half the limit of detection of that cytokine.

SBI was defined as bacteremia, meningitis, pneumonia, osteomyelitis, urinary tract infection (UTI) or any other severe infection with the identification of a plausible microorganism in a normally sterile site (only confirmed SBI were included in this group). For the diagnosis of UTI, pyuria in urine analysis was required in addition to the positive urine culture. Viral infection (VI) was defined as a positive result in the multiplex-PCR from respiratory samples. Patients with a bacterial infection and a viral detection in the respiratory sample at the same time were classified in the group of bacterial-viral coinfections and excluded from the subanalyses. Patients with no proven infection (NPI) were those not included in the previous groups.

Variables evaluated

Data were collected in a database specifically designed for the study. Data related to cases and controls included baseline characteristics such as age, gender, SCD type, diagnostic method, parents' continent of origin, immunization status (including vaccination against *Haemophilus influenzae* type b, *Streptococcus pneumoniae* and *Neisseria meningitidis* serogroups A, B, C, W, Y), penicillin prophylaxis, treatment with hydroxyurea, other treatments, long-term central venous catheter (CVC), splenectomy and history of previous hospital admissions. Clinical data during the episode (only in cases) included duration of fever before admission, maximum axillary temperature during the episode, respiratory symptoms, hemodynamic instability (defined as hypotension or persistent tachycardia requiring fluid resuscitation or inotropic drugs) and hypoxemia (< 92%). Laboratory parameters included hemoglobin, WBC, neutrophils and platelets on admission, initial and maximum CRP and procalcitonin during the episode, blood cultures, as well as other cultures and viral tests. Other recorded data were inpatient/outpatient management, antimicrobial treatment, need for antibiotic change because of unfavorable clinical course, diagnosis of VOC or acute chest syndrome (ACS), duration of fever and of hospitalization, pediatric intensive care unit (PICU) admission and final outcome (hospital discharge or exitus). Cytokine levels in all cases and controls were also evaluated.

Statistical analysis

Qualitative variables were expressed as frequencies or rates. Quantitative variables were reported as means (standard deviation) or medians [interquartile range (IQR)], according to their distribution. The χ test was used to compare categorical variables; Yates correction was applied to any contingency table and Fisher's exact test was used when at least one cell had a value <5. T-test and ANOVA were used to compare the means of normally distributed variables whereas Mann-Whitney U test and Kruskal-Wallis test were performed to compare the medians of variables not normally distributed. Receiver operating characteristics (ROC) curves and area under the ROC curve (AUC) were calculated for significant parameters. The optimal cut-off point of each biomarker was selected using the Youden Index, based on the maximum sum of sensitivity and

specificity. Sensitivity, specificity, positive predictive values (VPP) and negative predictive values (NPV) were calculated according to various possible prevalence rates (PR) of SBI. A p value <0.05 was considered significant. Statistical analysis was performed using IBM SPSS Statistics software version 22.0 (IBM Corp., Armonk, N.Y., USA).

RESULTS

A total of 137 patients were included in the study: 78 cases and 59 controls (Figure 1). Median age of the patients was 4.1 (1.7-7.5) years for cases and 5.2 (2.1-9.3) for controls ($p = 0.193$), with a higher percentage of males in the group of cases (78.5% vs 62.1%, $p = 0.035$). The majority of the children were born in Spain and had been diagnosed by newborn screening, while their parents were predominantly from Africa and South/Central America. Most patients were appropriately immunized and receiving penicillin prophylaxis, while approximately half of them were on hydroxyurea treatment. Baseline characteristics of cases and controls are summarized in Table 1.

Four out of 78 cases (5.1%) were diagnosed with SBI: two of them were catheter-related bacteremia (caused by *Staphylococcus aureus* and *Enterobacter cloacae*), one *Streptococcus pneumoniae* (serotype 9N) bacteremic pneumonia and one *Escherichia coli* UTI. Another patient was also diagnosed with *Escherichia coli* UTI, but she had a viral detection in a respiratory sample at the same time (influenza B) and, therefore, she was considered a bacterial-viral coinfection and excluded from the subanalyses. A virus was detected by multiplex-PCR in 41/78 (52.6%) of the respiratory samples. The most frequently detected virus was influenza (A or B) in 23.3% of the cases, followed by rhinovirus in 20.9%, adenovirus in 14% and enterovirus in 11.6%. When comparing the baseline characteristics among cases with SBI, VI and the rest of patients with NPI (Supplemental Table 1), no statistically significant differences were found, except for the presence of CVC and hypertransfusional regimen, which were more frequent in the SBI group (50% vs 9.8% vs 36.4%, $p = 0.011$ and 50% vs 4.9% vs 15.2%, $p = 0.018$, respectively).

Data on clinical presentation, laboratory parameters and outcome during the febrile episodes in all cases, and comparisons among groups, are summarized in Table 2. Most patients were hospitalized (81%) and received at least one dose of empiric antibiotic treatment (96.2%). Three patients (3.8%) required PICU admission and no patient died. Patients with SBI had more often hemodynamic instability compared to patients with VI or NPI (25% vs 2.4% vs 0, respectively; $p = 0.012$) and significantly higher inflammatory parameters (neutrophils, CRP and procalcitonin). They required more frequently a change in the antibiotic therapy (75% vs 7.5% vs 15.6%; $p = 0.001$) and longer hospital admissions (7.5 vs 3 vs 5 days; $p = 0.013$). Upper respiratory symptoms were more frequent in patients with VI compared to patients with SBI or NPI (80.5% vs 50% vs 42.4%, respectively; $p = 0.003$).

Analysis of proinflammatory cytokine levels and other biomarkers

IL-6 was significantly higher in patients with SBI than in those with VI or NPI [median (IQR) pg/ml: 163 (70.4-459.5) vs 0.7 (0.7-0.7) vs 0.7 (0.7-0.7), respectively; $p < 0.001$], being undetectable in all controls. To note, the patient with the highest level of IL-6 was the case of confirmed bacterial pneumonia (733 pg/ml). However, in the patient with the bacterial-viral coinfection, IL-6 was undetectable. Figure 2 represents the values of IL-6 and other significant biomarkers (neutrophils, CRP and PCT) of the study subgroups, including the optimal cut-off point for each parameter calculated by the Youden index. IL-12p70 was not detected in any of the cases or controls. The rest of the cytokines analyzed (IL-1 β , IL-2, IL-4, IL-8, IL-10, IL-17a, TNF α , IFN- γ) did not show any significant difference between cases and controls or among different groups of cases. No differences were found in any of the cytokines analyzed between the cases diagnosed with VOC and the rest of the patients. None of the cytokine levels significantly correlated with possible confounders as the age or the number of previous admissions.

The AUC of IL-6 to discriminate confirmed SBI was 0.87 (95% CI 0.77-0.93), similarly to the other significant parameters [neutrophils 0.87 (0.78-0.94); CRP 0.89 (0.81-0.96); procalcitonin 0.84 (0.73-0.91)]. The optimal cut-off value of IL-6 for the diagnosis of SBI was 125 pg/mL. Data of sensibility, specificity, PPV and NPV (the last two according to various possible PR of SBI) of IL-6 and other significant biomarkers are represented

in Table 4. IL-6 had very high values of PPV and NPV, NPV values being similar to other biomarkers, with significantly higher PPV values (PPV of 100% for a PR of 5, 10 and 15%; NPV of 98.7%, 97.3% and 95.8% for a PR of 5, 10 and 15%, respectively).

DISCUSSION

In this prospective, case-control study we compared the clinical characteristics and laboratory parameters, including 10 serum proinflammatory cytokines, in SCD children with fever of different etiologies and asymptomatic steady-state controls. We found that IL-6 was a very good biomarker for the diagnosis of SBI in these patients, with similar NPV to other biomarkers and better PPV. To our knowledge, this is the first study describing IL-6 as a marker of SBI in febrile SCD children.

Most of the children included in this study had been diagnosed with SCD by newborn screening, had a complete vaccination status and were receiving penicillin prophylaxis. In the group of SCD patients with fever, we found a low proportion of confirmed SBI (5.1%), similarly to other studies performed in high-income countries^{1,3–5}. However, in more than a half of the cases (52.6%) a virus was detected by multiplex-PCR from respiratory samples, as it has been observed in recent pediatric studies^{16,17}. In the remaining 42.8% of cases an infectious etiology was not confirmed. The only statistically significant differences among the baseline characteristics of study subgroups were a higher percentage of patients with CVC and on hypertransfusional regimen in the group of SBI, suggesting a poorer clinical baseline condition of these patients and also highlighting the presence of long-term CVC as a predisposing factor for bacteremia⁶.

Despite the low proportion of confirmed SBI in this cohort, almost all patients received at least one dose of antibiotic (96.2%), in agreement to current guidelines^{18–20}. This high use of antibiotics emphasizes the importance of defining laboratory parameters that could early discriminate the etiology of fever in these patients, in order to optimize the antimicrobial prescription. Children with SBI presented more frequently with hemodynamic instability and had higher inflammatory parameters (neutrophils, CRP and procalcitonin) than the rest of the patients, similarly to what was described in other studies^{1,5–9}. We also found that children with SBI had significantly higher IL-6 levels when compared with those with VI, NPI and controls, with the highest value observed in the only patient with confirmed pneumococcal pneumonia. The optimal cut-off value of IL-6 for the diagnosis of SBI was 125 pg/ml, with a very high PPV (significantly better than other biomarkers) and NPV applying to different PR. In a setting with a PR of SBI of 5% in children with SCD (average PR in high-income countries^{1,4,5}), the PPV would be 100% and NPV 98.7%. Therefore, we consider that IL-6 could be a very useful marker for the diagnosis of SCD patients with confirmed SBI.

IL-6 is an important factor in systemic and local immune response. It can be secreted by almost all stromal cells and cells of the immune system at an early stage of inflammation (with rapid elevation in 1-3 hours), and further promote the synthesis and release of CRP by the liver^{21,22}. IL-6 has been previously described as a marker of different types of SBI (alone or in combination with other biomarkers)^{23,24}, such as early-onset neonatal sepsis^{25,26}, sepsis and septic shock^{21,23,27}, bacteremia in patients with febrile neutropenia^{28,29} or pneumococcal pneumonia^{30–33}. The early elevation of IL-6 that occurs in the presence of SBI is an important advantage of this cytokine over other biomarkers.

In SCD patients, it has previously been described that IL-6 and other proinflammatory cytokines are elevated in asymptomatic steady-state patients when compared to healthy individuals^{10–13}. However, in our study all controls (steady-state SCD patients) had undetectable levels of IL-6. A possible explanation for this could be that almost all children included in this study were receiving vitamin D supplements. According to the study performed by Adegoke et al., children with SCD had a maintained vitamin D deficiency with a high level of proinflammatory cytokines (IL-6, IL-8 and IL-18), which normalizes after vitamin D supplementation³⁴. Other previous studies also found higher levels of IL-6 in patients with VOC^{14,15}, a finding that we did not observe in our study.

The strengths of this study were its prospective design, the homogeneity of the patients and that it was carried out in a reference center for patients with SCD in Spain; therefore, this cohort may be quite representative of the pediatric population with SCD in high-income countries. However, it has also several limitations.

Most importantly, the sample size was relatively small, with few cases of confirmed bacterial infection, and the categorization in several study groups might have decreased the statistical power. Blood samples were collected regardless the previous duration of fever and serial samples were not collected, so it was not possible to evaluate the trend over time of the different biomarkers during the febrile episode. Finally, only confirmed SBI were included in the study to avoid bias in the analysis, but maybe some not-confirmed bacterial infections (e. g. pneumonia) were included in the NPI group.

Conclusion

In this prospective study, we found that IL-6 (with an optimal cut-off of 125 pg/ml) was a good marker of SBI with a high PPV and NPV. Therefore, given its rapid elevation, IL-6 may be useful (alone or in combination with other biomarkers) to early discriminate SCD children at risk of SBI, in order to guide their management. In settings with a low PR of SBI, its use could help to reduce unnecessary antibiotic treatments and hospital admissions. These findings should be confirmed in larger cohorts and multicenter studies.

ACKNOWLEDGEMENTS

We thank HGUGM nurses for their help in collecting samples and the Spanish HIV HGM Biobank for storing and processing blood samples for cytokine analysis. We also thank Ana María Rincón-López, official translator, for her careful review of the manuscript.

COLLABORATORS

Collaborators of F-DREP Study Group:

- Carmen Garrido Colino. Pediatric Hematology and Oncology Unit, Hospital General Universitario Gregorio Marañón, Madrid, Spain.
- Jorge Huerta Aragonés. Pediatric Hematology and Oncology Unit, Hospital General Universitario Gregorio Marañón, Madrid, Spain.
- Cristina Mata Fernández. Pediatric Hematology and Oncology Unit, Hospital General Universitario Gregorio Marañón, Madrid, Spain.
- Concepción Míguez Navarro. Pediatric Emergency Unit, Hospital General Universitario Gregorio Marañón, Madrid, Spain.
- Andrea Mora Capín. Pediatric Emergency Unit, Hospital General Universitario Gregorio Marañón, Madrid, Spain.
- Rafael Marañón Pardillo. Pediatric Emergency Unit, Hospital General Universitario Gregorio Marañón, Madrid, Spain.
- Arístides Rivas García. Pediatric Emergency Unit, Hospital General Universitario Gregorio Marañón, Madrid, Spain.
- Paula Vázquez López. Pediatric Emergency Unit, Hospital General Universitario Gregorio Marañón, Madrid, Spain.
- Jose Luis Jiménez Fuentes. Immunology Section, Laboratorio InmunoBiología Molecular, Instituto Investigación Sanitaria Gregorio Marañón, Hospital General Universitario Gregorio Marañón, Spanish HIV HGM BioBank, Madrid, Spain.
- Maria Ángeles Muñoz Fernández. Immunology Section, Laboratorio InmunoBiología Molecular, Instituto Investigación Sanitaria Gregorio Marañón, Hospital General Universitario Gregorio Marañón, Spanish HIV HGM BioBank, Madrid, Spain.
- Rosario Zamarro Arranz. Health educator of parents and children with sickle cell disease. Nurse in Hospital General Universitario Gregorio Marañón, Madrid, Spain.

FUNDING: This work was supported by an unrestricted ESPID (European Society for Pediatric Infectious Diseases) Grant Award. The sponsor did not participate in the study design or interpretation of the results.

CONFLICT OF INTEREST: The authors have no conflicts of interest or financial relationships relevant to this article to disclose.

REFERENCES

1. Morrissey BJ, Bycroft TP, Almossawi O, Wilkey OB, Daniels JG. Incidence and Predictors of Bacterial infection in Febrile Children with Sick Cell Disease. *Hemoglobin*. 2015;39(5):316–9.
2. Cannas G, Merazga S, Viro E. Sick cell disease and infections in high- and low-income countries. *Mediterr J Hematol Infect Dis*. 2019;11(1):1–9.
3. Narang S, Fernandez ID, Chin N, Lerner N, Weinberg GA. Bacteremia in children with sickle hemoglobinopathies. *J Pediatr Hematol Oncol*. 2012; 34:13–16.
4. Bansil NH, Kim TY, Tieu L BB. Incidence of Serious Bacterial Infections in Febrile Children With Sick Cell Disease. *Clin Pediatr (Phila)*. 2013;52(7):661–6.
5. Rincón-López EM, Navarro Gómez ML, Hernández-Sampelayo Matos T, Saavedra-Lozano J, Aguilar de la Red Y, Hernández Rupérez B, et al. Low-risk factors for severe bacterial infection and acute chest syndrome in children with sickle cell disease. *Pediatr Blood Cancer*. 2019;66(6):1–7.
6. Chang TP, Kriengsoontorkij W, Chan LS, Wang VJ. Predictors for Bacteremia in Febrile Sick Cell Disease Children in the Post-7-Valent Pneumococcal Conjugate Vaccine Era. *J Pediatr Hematol Oncol*. 2013;35:377–382.
7. Elenga N, Placide L, Cuadro-alvarez E, Long L, Razafindrakoto SH, Mrcic Y, et al. Does Procalcitonin Predict Bacterial Infection in Febrile Children with Sick Cell Disease? *Indian J Pediatr*. 2019;86(1):95–96.
8. Savlov D, Beck CE, DeGroot J, Odame I, Friedman JN. Predictors of Bacteremia Among Children With Sick Cell Disease Presenting With Fever. *J Pediatr Hematol Oncol*. 2014;36(5):384–8.
9. West DC, Andrada E, Azari R, Rangaswami AA, Kuppermann N. Predictors of Bacteremia in Febrile Children With Sick Cell Disease. *J Pediatr Hematol Oncol*. 2002;24(4):279–83.
10. Olenski Gilli SC, Pericole FV, Benites BD, Sippert EÂ, Castilho LM, Addas-Carvalho M, et al. Cytokine polymorphisms in sickle cell disease and the relationship with cytokine expression. *Exp Hematol*. 2016;44(7):583–9.
11. Hibbert JM, Hsu LL, Bhathena SJ, Irune I, Sarfo B, Creary MS, et al. *Exp Biol Med (Maywood)*. 2005;230(1):68–74.
12. Taylor SC, Shacks SJ, Mitchell RA, Banks A. Serum Interleukin-6 Levels in the Steady State of Sick Cell Disease. *J Interf Cytokine Res*. 1995;15(12):1061–4.
13. Taylor SC, Shacks SJ, Qu Z, Wiley P. Type 2 Cytokine Serum Levels in Healthy Sick Cell Disease Patients. *J Natl Med Assoc*. 1997;89(11):753–7.
14. Pathare A, Al Kindi S, Alnaqdy AA, Daar S, Knox-Macaulay H, Dennison D. Cytokine profile of sickle cell disease in Oman. *Am J Hematol*. 2004;77(4):323–8.
15. Sarray S, Saleh LR, Lisa Saldanha F, Al-Habboubi HH, Mahdi N, Almawi WY. Serum IL-6, IL-10, and TNF α levels in pediatric sickle cell disease patients during vasoocclusive crisis and steady state condition. *Cytokine*. 2015;72(1):43–7.
16. Ploton MC, Sommet J, Koehl B, Gaschignard J, Holvoet L, Mariani-Kurkdjian P, et al. Respiratory pathogens and acute chest syndrome in children with sickle cell disease. *Arch Dis Child*. 2020;105(9):891–5.
17. Srinivasan A, Wang WC, Gaur A, Smith T, Gu Z, Kang G, et al. Prospective Evaluation for Respiratory Pathogens in Children With Sick Cell Disease and Acute Respiratory Illness. *Pediatr Blood Cancer*. 2014;61:507–11.
18. Cela, E, Ruiz, A, Cervera Á. Enfermedad de células falciformes. Guía de práctica clínica de la Sociedad Española de Hematología y Oncología Pediátricas. CeGe, editor. 2019. Available from:

<http://www.sehop.org/wp-content/uploads/2019/03/Guía-SEHOP-Falciforme-2019.pdf>

19. Buchanan G. Evidence-Based Management of Sickle Cell Disease. NIH Expert panel report 2014. Available from: https://www.nhlbi.nih.gov/sites/default/files/media/docs/sickle-cell-disease-report_020816.0.pdf
20. Noronha SA, Sadreameli SC, Strouse JJ. Management of Sickle Cell Disease in Children. *South Med J*. 2016;109(9):495–502.
21. Shao WX, Yu DJ, Zhang WY, Wang XJ. Clinical significance of interleukin-6 in the diagnosis of sepsis and discriminating sepsis induced by gram-negative bacteria. *Pediatr Infect Dis J*. 2018;37(8):801–5.
22. Hunter CA, Jones SA. IL-6 as a keystone cytokine in health and disease. *Nat Immunol*. 2015;16(5):448–57.
23. Slaats J, ten Oever J, van de Veerdonk FL, Netea MG. IL-1 β /IL-6/CRP and IL-18/ferritin: Distinct Inflammatory Programs in Infections. *PLoS Pathog*. 2016;12(12):1–13.
24. Holub M, Lawrence DA, Andersen N, Davidová A, Beran O, Marešová V, et al. Cytokines and chemokines as biomarkers of community-acquired bacterial infection. *Mediators Inflamm*. 2013;2013:190145.
25. Cernada M, Badía N, Modesto V, Alonso R, Mejías A, Golombek S, et al. Cord blood interleukin-6 as a predictor of early-onset neonatal sepsis. *Acta Paediatr Int J Paediatr*. 2012;101(5):203–7.
26. Tasci Y, Dilbaz B, Uzmez Onal B, Caliskan E, Dilbaz S, Doganci L, et al. The value of cord blood interleukin-6 levels for predicting chorioamnionitis, funisitis and neonatal infection in term premature rupture of membranes. *Eur J Obstet Gynecol Reprod Biol*. 2006;128(1–2):34–9.
27. Waage A, Brandtzaeg P, Halstensen A, Kierulf P, Espevik T. The complex pattern of cytokines in serum from patients with meningococcal septic shock. Association between interleukin 6, interleukin 1, and fatal outcome. *J Exp Med*. 1989;169(1):333–8.
28. Bal ZŞ, Özdemir NK, Şen S, Karapınar DY, Azarsız E, Aydemir Ş, et al. Diagnostic Accuracy of Interleukin-6, Interleukin-8, and Interleukin-10 for Predicting Bacteremia in Children with Febrile Neutropenia. *Turkish J Hematol*. 2017;34(3):254–7.
29. Diepold M, Noellke P, Duffner U, Kontny U, Berner R. Performance of Interleukin-6 and Interleukin-8 serum levels in pediatric oncology patients with neutropenia and fever for the assessment of low-risk. *BMC Infect Dis*. 2008;8:2–8.
30. Nascimento-Carvalho EC, Vasconcellos ÂG, Clarêncio J, Andrade D, Barral A, Barral-Netto M, et al. Evolution of cytokines/chemokines in cases with community-acquired pneumonia and distinct etiologies. *Pediatr Pulmonol*. 2020;55(1):169–76.
31. Vasconcellos ÂG, Clarêncio J, Andrade D, Cardoso MRA, Barral A, Nascimento-Carvalho CM. Systemic cytokines and chemokines on admission of children hospitalized with community-acquired pneumonia. *Cytokine*. 2018;107:1–8.
32. Fuchs A, Gotta V, Decker ML, Szinnai G, Baumann P, Bonhoeffer J, et al. Cytokine kinetic profiles in children with acute lower respiratory tract infection: a post hoc descriptive analysis from a randomized control trial. *Clin Microbiol Infect*. 2018;24(12):1341.
33. Calbo E, Alsina M, Rodríguez-Carballeira M, Lite J, Garau J. Systemic expression of cytokine production in patients with severe pneumococcal pneumonia: Effects of treatment with a β -lactam versus a fluoroquinolone. *Antimicrob Agents Chemother*. 2008;52(7):2395–402.
34. Adegoke SA, Smith OS, Adekile AD, Figueiredo MS. Relationship between serum 25-hydroxyvitamin D and inflammatory cytokines in paediatric sickle cell disease. *Cytokine*. 2017;96:87–93.

LEGENDS

- Figure 1. Flow diagram.

- Figure 2. Biomarkers' values in the study subgroups and optimal cut-off points.
- Table 1. Baseline characteristics of patients.
- Table 2. Characteristics of cases during the febrile episode and differences among study subgroups.
- Table 3. Cytokines values and comparisons among study subgroups.
- Table 4. Diagnostic tests of IL-6 and other significant biomarkers for the prediction of SBI.
- Supplemental table 1. Baseline characteristics of cases and comparisons among study subgroups.

Figure 1. Flow diagram





Figure 2. Biomarkers' values in the study subgroups and optimal cut-off points



Hosted file

TABLE 1. Baseline characteristics of patients.docx available at <https://authorea.com/users/469331/articles/562775-interleukin-6-as-a-marker-of-severe-bacterial-infection-in-children-with-sickle-cell-disease-and-fever>

Hosted file

TABLE 2. Characteristics of cases during the febrile episode and differences among study subgroups.docx available at <https://authorea.com/users/469331/articles/562775-interleukin-6-as-a-marker-of-severe-bacterial-infection-in-children-with-sickle-cell-disease-and-fever>

Hosted file

TABLE 3. Cytokines values and comparisons among study subgroups.docx available at <https://authorea.com/users/469331/articles/562775-interleukin-6-as-a-marker-of-severe-bacterial-infection-in-children-with-sickle-cell-disease-and-fever>

Hosted file

TABLE 4. Diagnostic tests of IL-6 and other significant biomarkers for the prediction of SBI.docx available at <https://authorea.com/users/469331/articles/562775-interleukin-6-as-a-marker-of-severe-bacterial-infection-in-children-with-sickle-cell-disease-and-fever>