

# Bronchiolitis Before and After the SARS-CoV2 Pandemic: Twelve Years of Experience in a Spanish Paediatric Hospital

Fuensanta Guerrero-del-Cueto<sup>1</sup>, J. M. Ramos Fernandez<sup>1</sup>, Isabel Leiva-Gea<sup>1</sup>, Elena Reina-Moreno<sup>2</sup>, Ana Ortiz-Ortigosa<sup>1</sup>, Begoña Carazo-Gallego<sup>1</sup>, Ana Cordon<sup>1</sup>, and Esmeralda Núñez-Cuadros<sup>1</sup>

<sup>1</sup>Hospital Regional Universitario de Malaga

<sup>2</sup>Universidad de Malaga Facultad de Medicina

January 16, 2023

## Abstract



*Introduction:* Acute bronchiolitis (AB) is the main cause of hospitalization in children under two years of age, with a regular winter seasonality, mostly due to the respiratory syncytial virus (RSV). *Objectives:* To describe the epidemiology of bronchiolitis hospitalizations in our centre in the last twelve years, and analyse the changes in clinical characteristics, microbiology, and adverse outcomes during the SARS-CoV2 pandemic. *Methods:* Ambispective observational study including patients admitted for bronchiolitis between April 2010 and December 2021 in a Spanish tertiary paediatric hospital. Relevant demographic, clinical, microbiological, and adverse outcome variables were collected in an anonymized database. The pandemic period (April 2020 to December 2021) was compared to 2010-2015 seasons using appropriate statistical tests. *Results:* There were 2138 bronchiolitis admissions, with a mean of 195.6 per year between 2010-2019 and a 2–4-month peak between November and March. In the winter of 2020, there was a 94.4% reduction of bronchiolitis hospitalizations, with only eleven cases admitted in the first year of the pandemic. Bronchiolitis cases increased during the summer of 2021 in a six-month long peak, reaching a total of 171 cases. Length of stay was significantly shorter during the pandemic, but no differences were found in clinical and microbiological characteristics or other adverse outcomes. *Conclusions:* The SARS-CoV2 pandemic has modified the seasonality of bronchiolitis hospitalizations, with a dramatic decrease in cases during the winter of 2020-2021, and an extemporaneous summer-autumn peak in 2021 with longer duration but similar patient characteristics.

## Hosted file

Bronchiolitis\_R2\_clean.docx available at <https://authorea.com/users/490452/articles/573824-bronchiolitis-before-and-after-the-sars-cov2-pandemic-twelve-years-of-experience-in-a-spanish-paediatric-hospital>

## Hosted file

Paediatric Pulmonology Bronchiolitis without figures and tables.docx available at <https://authorea.com/users/490452/articles/573824-bronchiolitis-before-and-after-the-sars-cov2-pandemic-twelve-years-of-experience-in-a-spanish-paediatric-hospital>



## Hosted file

Image legends.docx available at <https://authorea.com/users/490452/articles/573824-bronchiolitis-before-and-after-the-sars-cov2-pandemic-twelve-years-of-experience-in-a-spanish-paediatric-hospital>

## Hosted file

Tables.docx available at <https://authorea.com/users/490452/articles/573824-bronchiolitis-before-and-after-the-sars-cov2-pandemic-twelve-years-of-experience-in-a-spanish-paediatric-hospital>