

Left and right ventricular functional changes after transapical, transfemoral, transcatheter, or surgical aortic valve replacement: A single-center observational study

Assami Rösner¹, Didrik Kjønaas², Siri Malm¹, and Henrik Schirmer³

¹Universitetssykehuset Nord-Norge Harstad

²Universitetssykehuset Nord-Norge Kreftavdelingen

³Akershus Universitetssykehus Indremedisinsk avdeling

July 7, 2022

Abstract

Background: Decreasing right ventricular (RV) and left ventricular (LV) function after surgical or transcatheter aortic valve replacement (SAVR or TAVR, respectively) is an important risk factor for morbidity and mortality. Although transapical (TA)-TAVR is an independent risk factor for post-procedural mortality, limited knowledge is available regarding long-term changes in RV and LV function. The study aimed to evaluate LV and RV performance following four different AVR procedures, including TA-, transfemoral (TF)-TAVR, and SAVR with and without coronary artery bypass grafting ($\pm$ CABG). **Methods:** Patients with severe AS were consecutively included and assigned to TA-TAVR, TF-TAVR, or SAVR $\pm$ CABG groups. A total of 130 patients underwent preoperative conventional and strain-rate-imaging echocardiography, with similar controls in the period between 6 and 12 months after the procedure. **Results:** After AVR, NYHA classes III and IV were reduced from 105 (81%) to 6 (5%) patients. While most of the systolic and diastolic functional parameters indicated improved LV function in the TF-TAVR and both SAVR groups, LV function did not significantly change after TA-TAVR. The right ventricular functional parameters were unchanged or even improved equally after TA-TAVR and TF-TAVR, while they were significantly reduced after SAVR. The Cardiac Index (CI) improved significantly after TF-TAVR from 2.3 ± 0.7 to 2.6 ± 0.7 , while staying unchanged after TA-TAVR and SAVR $\pm$ CABG. **Conclusion:** This study demonstrated significant changes in LV and RV systolic and diastolic function with functional improvement or deterioration depending on the type of aortic valve replacement. The most significant improvement in CI was observed after TF-TAVR, which is the least invasive procedure.

Left and right ventricular functional changes aftertransapical,transfemoral, transcatheter, or surgical aortic valve replacement: A single-center observational study

Assami Rösner MD, PhD^{1,2}, Didrik Kjønaas MD, PhD³, Siri Malm MD, PhD⁴, Henrik Schirmer MD, PhD^{5,6}

¹ Department of Cardiology, Division of Cardiothoracic and Respiratory Medicine, University Hospital of North Norway, Norway

² Institute of Clinical Medicine, UiT, The Arctic University of Norway, Tromsø, Norway

³ Department of General and Gastrointestinal Surgery, Division of Surgery, Oncology and Women's Health, University Hospital of North Norway, Norway

⁴ Department of Internal Medicine, Division of Internal Medicine, University Hospital of North Norway, Harstad

⁵ Department of Cardiology, Akershus University Hospital, Lørenskog, Norway

⁶ Institute of Clinical Medicine, Cardiovascular Research Group, Campus Ahus, University of Oslo, Norway

Corresponding Author: Assami Rösner, MD, PhD

Department of Cardiology, Division of Cardiothoracic and Respiratory Medicine,

University Hospital of North Norway, 9019 Tromsø, Norway

Tel: +47 77627347

e-mail: assami.rosner@unn.no

List of abbreviations:

AS: aortic stenosis

AVR: aortic valve replacement

CABG: Coronary artery bypass grafting

CI: cardiac index

CO: cardiac output

DT: E wave deceleration time

E: E wave velocity

e': early diastolic tissue Doppler velocity

ET: ejection time

FAC: fractional area change

HV-SFF: hepatic vein systolic filling fraction

IVRT: isovolumic relaxation time

LV: left ventricular

LVOT: left ventricular outflow tract

MAPSE: mitral annulus plane systolic excursion

MV: mitral valve

PSV: peak systolic velocity by tissue Doppler

PV-SFF: pulmonary vein systolic filling fraction

RV: right ventricular

SAVR: surgical aortic replacement

SV: stroke volume

TA: transapical

TAPSE: tricuspid annulus plane systolic excursion

TAVR: Transcatheter aortic valve replacement

TF: transfemoral

TR_{peak}: tricuspid regurgitation systolic peak gradient

TV: tricuspid valve

Declarations

Ethics approval and consent to participate

The study was approved by the Regional Ethical Committee of North Norway (REK Nord 397/2010). Written informed consent was obtained from the patients for publication of their individual details and accompanying images in this manuscript. The consent form is held by the author's institution and is available for review by the Editor-in-Chief.

Availability of data and materials

The data are stored as a research file at the University Hospital in North Norway with limited access to dedicated researchers.

Competing interests

The authors declare that they have no competing interests.

Funding

This work was funded and supported by the Regional Health Authorities of North Norway (HN-ID 6884/SFP1078-12) and (HN SFP 1172-14)

Authors' contributions

AR conceived, designed the study, handled the funding, acquired data, performed statistical analysis and wrote the manuscript. SM acquired data. DK performed strain-analysis. AR performed statistical analysis. HS and SM made critical revision of the manuscript for important intellectual content.

Acknowledgements

We would like to thank Editage (www.editage.com) for English language editing.

Abstract

Background : Decreasing right ventricular (RV) and left ventricular (LV) function after surgical or transcatheter aortic valve replacement (SAVR or TAVR, respectively) is an important risk factor for morbidity and mortality. Although transapical (TA)-TAVR is an independent risk factor for post-procedural mortality, limited knowledge is available regarding long-term changes in RV and LV function. The study aimed to evaluate LV and RV performance following four different AVR procedures, including TA-, transfemoral (TF)-TAVR, and SAVR with and without coronary artery bypass grafting ($\pm$ CABG).

Methods: Patients with severe AS were consecutively included and assigned to TA-TAVR, TF-TAVR, or SAVR $\pm$ CABG groups. A total of 130 patients underwent preoperative conventional and strain-rate-imaging echocardiography, with similar controls in the period between 6 and 12 months after the procedure.

Results: After AVR, NYHA classes III and IV were reduced from 105 (81%) to 6 (5%) patients. While most of the systolic and diastolic functional parameters indicated improved LV function in the TF-TAVR and both SAVR groups, LV function did not significantly change after TA-TAVR. The right ventricular functional parameters were unchanged or even improved equally after TA-TAVR and TF-TAVR, while they were significantly reduced after SAVR. The Cardiac Index (CI) improved significantly after TF-TAVR from 2.3 ± 0.7 to 2.6 ± 0.7 , while staying unchanged after TA-TAVR and SAVR $\pm$ CABG.

Conclusion: This study demonstrated significant changes in LV and RV systolic and diastolic function with functional improvement or deterioration depending on the type of aortic valve replacement. The most significant improvement in CI was observed after TF-TAVR, which is the least invasive procedure.

Keywords : aortic stenosis; interventional cardiology; surgical aortic valve replacement; transcatheter aortic valve replacement; transapical; transfemoral

Background

Transcatheter aortic valve replacement (TAVR) is an established alternative to surgical aortic replacement (SAVR) in elderly patients with severe symptomatic aortic stenosis (AS)(1). Transapical (TA) TAVR is a substantial risk factor for increased post-procedural mortality(2). Additionally, decreased right ventricular (RV) and left ventricular (LV) functions after SAVR or TAVR have been indicated as important risk factors for morbidity and mortality(3, 4). Although, LV and RV systolic and diastolic functional changes after transfemoral (TF) TAVR and SAVR have been described in previous studies(5-7), the influence of moderately invasive procedures, such as TA-TAVR, on LV and RV function has been scarcely investigated, rendering controversial results(8, 9).

In AS, the LV responds to chronic pressure overload by developing myocardial hypertrophy, fibrosis, and global systolic and diastolic dysfunctions. Several studies have shown the reversibility of hypertrophy and improvement of systolic function, as well as decreased filling pressures after both SAVR and TAVR(10-15). In AS with preserved ejection fraction (EF), myocardial strain and peak systolic velocity (PSV) by tissue Doppler are highly sensitive markers of LV functional improvement after aortic valve replacement (AVR)(14-17). In contrast, the right ventricular (RV) function and geometry are less affected by severe AS. Thus, despite reduced systolic pulmonary artery pressure, RV function has not been shown to improve, but rather to deteriorate after SAVR. Following open cardiac surgery, both RV geometry and function deteriorate, while the TAVR procedure does not seem to affect RV function(5, 7, 18). The present study aimed to investigate late postoperative changes (6–12 months) in LV and RV function after TAVR and SAVR, with a focus on potential differences between TA and TF access for TAVR, compared to SAVR with and without coronary artery bypass grafting (CABG). We performed conventional systolic and diastolic functional echocardiographic parameters from LV and RV tissue Doppler imaging (TDI) and speckle tracking echocardiography (STE) with strain-rate imaging to detect changes after AVR. Since TA-TAVR is more invasive than TF-TAVR, we hypothesized that TA-TAVR might show less LV improvement and decreased RV function. We also investigated whether CABG was an additional factor that improved or deteriorated the LV or RV function.

Methods

Study design, setting, and population

This single-center study was performed at the University Hospital of North Norway, Tromsø. Between February 2010 and June 2013, 175 consecutive patients with severe symptomatic AS eligible for either TAVR or SAVR were included in the study. The indication for TAVR or SAVR was based on a decision made by a multidisciplinary heart team, determined by the patient's suitability for either method, technical feasibility, the risk for open-heart surgery, age, comorbidities, and mental status. Patients with an inability to provide informed consent, a life expectancy of fewer than 12 months, and low motivation for interventional treatment were excluded from the study. All study participants were invited to undergo a preoperative clinical assessment and echocardiography with post-procedural control echocardiography at 6 (range 5–7) and 12 (range 11–13) months. All patients who returned to, at least one of the clinical control visits were included in the study. The study population was divided into four groups; TA- TAVR. TF-TAVR and SAVR with and without CABG. Clinical characteristics, mortality, and perioperative complications were obtained from the patients' electronic journals.

The study was approved by the Regional Ethical Committee of North Norway (REK Nord 397/2010) and all patients provided written informed consent.

Echocardiography

All patients underwent preoperative transthoracic echocardiographic (TTE) evaluation using an iE33 scanner (S5-1 probe, Philips Medical Systems, Andover, MA, USA) in the left lateral decubital position. Conventional 2-dimensional greyscale images were obtained in the parasternal long and short axes, as well as in the apical four-, two-, and three-chamber views. The LVEF was derived from the standard biplane Simpson mode. The apical four- and two-chamber views were used to calculate left atrial volumes at end-systole. Mitral annular plane systolic excursion (MAPSE) and peak systolic velocity (PSV) by tissue Doppler were measured in the septal and lateral mitral rings in the apical four-chamber view, reported as the average

of these. Intraventricular septum thickness and LV mass in diastole were measured on M-mode images of the parasternal long-axis view. Diastolic LV function was assessed by E/A ratio, E/e' ratio, E deceleration time (DT), E-Wave velocity, the mean of septal and lateral wall tissue Doppler velocities (e'), the systolic filling fraction of the pulmonary veins (PV-SFF), the left atrial volume at end-systole, the peak gradient over the tricuspid regurgitation (TR_{peak}) and the isovolumetric relaxation time (IVRT). The degree of AS was expressed as the mean gradient estimated from the Doppler flow across the aortic valve and indexed aortic valve opening area, derived by the continuity equation. The LV stroke volume (SV), cardiac output (CO), and cardiac index (CI) were derived from the left ventricular outflow tract (LVOT) diameter and the LVOT velocity time integral. To assess RV geometry, the longitudinal diameter between the RV apex and center of the tricuspid valve, largest ventricular transverse diameter, RV diastolic area, and RV systolic area were measured in a 4Ch view optimized for the RV ventricle. For RV systolic function, tricuspid annular systolic excursion (TAPSE), PSV of the free RV lateral wall, and fractional area change (FAC) were measured. RV diastolic function was measured using the tricuspid inflow parameters RV E/A ratio, E deceleration time (RV E DT), tricuspid E velocity, tissue Doppler RV peak early diastolic velocity (RV e'), RV E/e', and SFF of the hepatic veins (HV-SFF).

Since a quantitative method for technical reasons was not possible in all patients, we performed a multiparametric, semiquantitative evaluation of mitral regurgitation and aortic regurgitation, as recommended in the guidelines(19).

Strain analyses

Strain and SR analyses were performed using the speckle-tracking software VVI 7 (Siemens, Mountain View, CA, USA). Longitudinal LV strain was obtained by analyzing the LV in the apical four-, two-, and three-chamber views, and circumferential LV strain was obtained by analyzing the midventricular short axis view. Peak systolic strain values were defined as the peak values between aortic valve opening and closure. The timing for strain analysis was derived from Doppler measurements of the aortic and mitral valves. The start of the cardiac cycle was defined as the peak R on the ECG.

Regional strain curves with artifacts due to reverberation, air artifacts, missing segments, or insufficient tracking were discarded based on subjective visual assessment. The peak global longitudinal and circumferential strains and strain rate (SR) were calculated from a global endocardial curve. In patients with atrial fibrillation, the strain from three cycles, if available, was obtained and then averaged.

The TAVR procedure

All TAVR procedures were performed under general anesthesia with either TF or TA access, using either the self-expanding Medtronic CoreValve (Medtronic Inc., Minneapolis, Minnesota, USA) or the Edwards SAPIEN balloon-expandable valve (Edwards Lifesciences, Irvine, California, USA). TF access was the preferred modality, but TA access was used in the presence of highly calcified and tortuous pelvic vessels, given the acceptable LVEF and respiratory function. Other access sites were excluded from this study. The valve was implanted during rapid pacing (180 beats/min). The valve function and degree of valvular leakage were evaluated using TTE before discharge.

As this study was performed in the early stage of TAVR, SAVR was performed when the patients were younger than 75 years, the Euro Score indicated low surgical risk, or if CABG was required for sufficient revascularization.

Statistical analysis

All statistical analyses were performed using SPSS 26 (SPSS Inc., Chicago, IL, USA). Since none of the measurements at the 6 months compared to the 12 months follow-up were significantly different, we compared the baseline measurements with the averaged measurements of both follow-up studies using a paired t-test. For patient characteristics and the comparison of pre-to post-procedural findings, one-way ANOVA was performed with Bonferroni post-hoc analysis. Categorical variables were compared using the chi-square test with separate subgroup analyses. P -values <0.05 were regarded as significant.

Reproducibility

To evaluate inter- and intraobserver variability for longitudinal and circumferential strains, 30 patients were randomly selected for repeated strain analysis by the first and second investigators. The variability was calculated as the level of agreement.

Results

Patients

Of the 175 initially included patients, 20 died during the procedure or in the first 6 months following surgery, whereas 8/37 (22%), 9/56 (16%), 3/38 (8%), and 0/25 (0%) patients underwent TA-TAVR, TF-TAVR, SAVR, and SAVR+CABG, respectively. An additional 23 patients did not meet either of the follow-up controls, mostly because of the reduced general condition combined with long travel distances across northern Norway. The echocardiographic data of the first visit were lost for one patient, and for one patient, the postoperative imaging results were excluded due to low imaging quality, leaving 130 participants for the final analysis of pre- and at least one of two postoperative clinical and echocardiographic controls in the period from 6 to 12 months.

Of the included patients with at least one postoperative control, 21 died between six months and two years after aortic valve replacement.

Patient characteristics of the four patient groups are listed in Table 1. The table shows that coronary artery disease (CAD) was present in nearly two-thirds of the patients, while 71/130 (46%) patients were revascularized in connection with AVR investigation or intervention.

The patients undergoing TAVR were significantly older and had higher Euro scores. The TA-TAVI group was mainly male, whereas female participants were dominant in the TF group. Only one out of 11 patients with recent myocardial infarction underwent SAVR, and one out of 27 patients with previous CABG was elected for SAVR. As expected, COPD was more frequent in the TAVR group than in the SAVR group. The number of patients with persistent atrial fibrillation or postoperative need for ventricular pacing did not differ significantly between the groups.

LV systolic and diastolic function

The parameters directly related to a successful AVR were similarly reduced after all procedures, including reduction of the transvalvular gradient, shortening of the ejection time, and reduction of myocardial mass.

Table 2 and Figure 1 demonstrate the indicators of systolic and diastolic LV function, overall cardiac performance, and their respective changes in the four groups. The NT-proBNP and NYHA class levels indicated a higher severity of pre-interventional heart failure in TAVR patients, followed by a significant reduction in NT-proBNP and NYHA class improvement in 94% of all patients. Indicators of overall cardiac performance, such as SVI and CI, increased only in the TF-TAVR group. LV volumes, as a marker of general LV function, decreased in all groups except the TA-TAVR group.

Longitudinal systolic functional parameters, such as MAPSE, PSV, longitudinal strain, and SR, increased significantly in the TF-TAVR and SAVR groups. However, the TA-TAVR group did not show significant changes in any longitudinal or circumferential functional parameters, except for improved mitral PSV. The circumferential strain and SR increased significantly only after TF-TAVR and SAVR + CABG. In summary, none of the functional parameters indicated a deterioration of LV systolic function after the procedure, while improvement in both longitudinal and circumferential parameters was most marked and consistent in the TF-TAVR group.

The diastolic LV parameters are shown in Figure 1 and Table 3. LV E/e' , TR_{peak} and IVRT indicated similarly reduced filling pressures after all types of intervention, while SFF of the pulmonary veins decreased and E/A ratio increased after SAVR but remained unchanged after TAVR. Interestingly, a reduction of E/e' was occurring mainly due to increasing e' at unchanged or even higher E wave velocity. However, differences

between the groups regarding pre-to post-procedural changes were not significant, except for the E/A ratio, which increased after SAVR only, indicating higher filling pressures after open surgery. MV e' (Table 3) increased after TAVR and SAVR, reflecting an improvement of LV relaxation properties.

RV systolic and diastolic function

The RV geometrical and functional systolic parameters are shown in Figure 2 and Table 4. After SAVR, the RV was shortened and dilated (higher RV area), and both the TAPSE and RV PSV were significantly reduced. RV function after SAVR deteriorated compared to TA-TAVR and TF-TAVR, in which the RV geometrical and systolic functional properties improved.

After SAVR, all RV diastolic parameters indicated deterioration of diastolic properties with higher filling pressures (Figure 3), which was significantly different from the unchanged or slightly improved diastolic RV properties after both TAVR procedures.

Results of the inter- and intra-observer variability are shown in Table 5.

Discussion

The present study demonstrated that after 6–12 months, LV systolic and diastolic properties improved significantly in all AVR procedures. Thus, the PSV MR, longitudinal strain, and circumferential strain increased significantly after both transcatheter and surgical procedures. Diastolic Doppler-based parameters and NT-proBNP levels indicated a reduction in LV filling pressures, which coincided with substantially improved symptoms. However, following surgical procedures such as SAVR and SAVR with CABG, RV systolic and diastolic function worsened, while TA-TAVR and TF-TAVR showed similarly unchanged indicators of RV systolic and diastolic function. There were no significant differences in pre-to post-procedural changes between SAVR with and without CABG.

LV systolic function

In this non-randomized study, patients were consecutively included as they were assigned to different procedures. Therefore, the TA and TF-TAVR groups were significantly older and had lower preoperative LV systolic deformations. NT-proBNP and diastolic parameters indicated higher preprocedural filling pressures in the TAVR group than in the SAVR group. However, the long-term effect of afterload reduction similarly improved the long- and short-axis deformations after all procedures. This was reflected by the post-procedural improvement in longitudinal strain, SR, PSV, and circumferential strain. In contrast to strain and SR, EF was less sensitive to significant changes in any of the procedures, but the mean values (except for TA-TAVR) indicated a tendency towards improvement. Changes in LV function after SAVR and TAVR have been the focus of several studies(17, 20-26). A review by Garg et al. in 2017 summarized these changes with uniformly increasing longitudinal and circumferential strain, regional function, and twist, whereas EF was reported as either higher or unchanged(22).

TA access for TAVR has emerged as a significant risk factor for early post-procedural mortality(2, 27, 28). The following question is whether the increased mortality in TA-TAVR could be partly due to the changes in the left ventricular functional properties. Previous echocardiographic and cardiac magnetic resonance studies have indicated reduced regional apical, radial, and longitudinal strains, and in some studies, neither global strain nor EF improved after TA-TAVR(24, 29). These findings were confirmed in the present study, in which TA-TAVR was the only procedure without significant improvement in longitudinal or circumferential strain and SR. However, none of these parameters showed deterioration of LV function.

As untreated AS comprises severe systolic and diastolic dysfunction, improvement in LV function seems to be an important factor for positive outcomes, and a lack of LV functional improvement might be connected with a higher risk of TA-TAVR.

LV diastolic function

As previously described in the same AS population(30), diastolic dysfunction consists of both impaired

relaxation and increased end-diastolic pressure due to increased ventricular and atrial filling and myocardial stiffness following fibrosis. Signs of increased stiffness due to high filling pressures are expected to reverse as a result of afterload reduction(31), whereas stiff myocardium due to macro- or micro-scarring is expected to render irreversibly high filling pressures(32). Accordingly, successful AVR led to signs of decreased filling pressures; thus, peak pulmonary artery pressure and MV E/e' decreased, and IVRT increased similarly after all types of procedures. However, in contrast to TAVR, after SAVR, the E/A ratio increased, and the SFF of the pulmonary veins decreased significantly. One explanation might be that E and A wave velocities and pulmonary venous flow are more dependent on ventricular elasticity rather than filling pressure. Based on this observation, we hypothesize that surgical procedures may lead to micro-scarring, especially after combined procedures, while LV elasticity after TAVR, even with TA access, remains unchanged.

Impaired relaxation seems to be reversible to some degree when stenosis is removed and ventricular pressure decreases, while age-related reduced relaxation remains. The best indicator reflecting improved relaxation was e' , which increased after all types of procedures, while DT, as a second relaxation indicator, improved significantly only after TF-TAVR. For TA-TAVR, only IVRT was significantly changed, but the small group size left the measured effect sizes generally underpowered. However, interpreting trends and effect size, the diastolic properties before and after intervention were similar in both TAVR groups, with substantially reduced filling pressures and improvement of early relaxation, while elasticity or compliance of the LV seemed to be unchanged.

RV systolic and diastolic function

In contrast to the improvement of LV function, RV systolic function is consistently reported to be reduced after SAVR, but improved or unchanged following TAVR(5, 7). In our study, the RV longitudinal diameter decreased in both SAVR and TAVR; as previously described, the transverse diameters increased only after SAVR. Okada et al. reported a reduction in these diameters after both TA-TAVR and TF-TAVR, whereas our data showed a slight reduction in both diameters and the RV area only in the TF-TAVR group, while the area after TA-TAVR remained unchanged. Reduction in the RV area can be regarded as a sign of overall improved RV function.

Open surgery was followed by a reduction in RV systolic function, while RV systolic parameters remained unchanged after TAVR. The reduction in TAPSE and RV PSV after open surgery is a known phenomenon and appears also to be present in minimally invasive procedures(33). MR studies have reported reduced RV EF and new RV scars after SAVR(34, 35). In addition to decreasing systolic RV function, in the present study, we observed a decrease in RV diastolic function after SAVR. Increasing E/e' and decreasing DT and SFF of the hepatic veins indicated increased RV filling pressures and/or loss of RV elasticity. Decreasing e' of the RV lateral wall resulted in reduced relaxation properties.

Our findings on systolic and diastolic RV function support previous observations of the negative impact of SAVR, in which the most plausible cause seems to be micro-scarring due to cardioplegia and extra-corporal circulation(34). TA-TAVR with a small pericardial incision does not interfere with RV structures and does not seem to affect RV configuration and function.

Overall performance and clinical improvement

The overall cardiac performance indicated by CI and SVi increased only significantly after TF-TAVR, while it remained unchanged after the other procedures. Since LV function improved after SAVR, the unchanged overall cardiac performance could be related to reduced RV function. In TA-TAVR, RV function remained unchanged, while the improvement in LV systolic and diastolic function was less pronounced. TA access is an independent risk factor for early post-procedural mortality(2). Thus, our data reflect a bias towards survivors. The early post-procedural function was not investigated in this study, but it has been previously shown to impact outcomes(3, 4).

The subjective reported improvement after AVR was convincing: 81% of 130 participants presented pre-procedural NYHA III-IV versus 5% of patients post-AVR. The NT-proBNP level, an indicator of diastolic

filling pressure, decreased in all the groups. Even though NT-proBNP levels were higher in the pre-operative TAVR patients, not just due to age, the reduction of NT-proBNP was most pronounced in the TF-TAVR patients. Our data indicate that this minimally invasive procedure results in the best overall cardiac performance after 6–12 months, whereas the reversibility of heart failure symptoms seems to be more dependent on successful afterload reduction. However, postoperative improvements in RV and LV function and reversibility of NT-proBNP after TAVR and SAVR have an impact on outcome and exercise capacity(4, 36).

Studie's limitations

This study has several limitations. First, owing to the high mortality in the TA-TAVR group, only 28 of the initially 37 included patients were available for follow-up examinations. This left our analyses underpowered due to small effect sizes. However, 24 and 28 patients in the small group were sufficient to show clinically relevant large effect sizes. Smaller changes, without statistical significance, could indicate the direction of change or differences between the SAVR and SAVR +CABG groups. Second, the study does not reflect the TAVR population to date, as it was conducted between 2010 and 2013 when the first regular TAVR at our hospital started in 2008. The patient population undergoing TAVR has profoundly changed, and TA-TAVR has become a rare procedure. Thus, a similar study on TA-TAVR with sufficient patient numbers would be difficult to perform to date. Third, this study was not randomized, leading to intervention groups comprising different age groups with differing symptoms, cardiac functions, and risk profiles. In the elderly age group with more severe AS, the LV and RV functions were significantly poorer, and the post-procedural functional recovery might have been impaired due to this. However, our data aimed to show clinically relevant post-procedural changes in comparison to the preoperative state. The advantage of less invasive procedures could be demonstrated since TF-TAVR showed the most significant improvement in overall cardiac performance even in a group with higher age and more myocardial scarring.

Conclusion

Six to twelve months after AVR, heart failure symptoms were reduced in most of the patients, with no difference between treatment groups, LV systolic function and indicators of LV filling pressures improved after all types of intervention. However, we found significant differences in several post-procedural LV and RV functional parameters depending on the type of intervention. The most significant improvement in CI after TF-TAVR indicated the best overall cardiac performance after the least-invasive procedure.

References

1. Vahanian A, Alfieri O, Al-Attar N, Antunes M, Bax J, Cormier B, et al. Transcatheter valve implantation for patients with aortic stenosis: a position statement from the European Association of Cardio-Thoracic Surgery (EACTS) and the European Society of Cardiology (ESC), in collaboration with the European Association of Percutaneous Cardiovascular Interventions (EAPCI). *Eur Heart J*. 2008;29:1463-70.
2. Kjonas D, Dahle G, Schirmer H, Malm S, Eidet J, Aaberge L, et al. Predictors of early mortality after transcatheter aortic valve implantation. *Open Heart*. 2019;6:e000936.
3. Haddad F, Couture P, Tousignant C, Denault AY. The right ventricle in cardiac surgery, a perioperative perspective: II. Pathophysiology, clinical importance, and management. *Anesth Analg*. 2009;108:422-33.
4. Sato K, Kumar A, Jones BM, Mick SL, Krishnaswamy A, Grimm RA, et al. Reversibility of Cardiac Function Predicts Outcome After Transcatheter Aortic Valve Replacement in Patients With Severe Aortic Stenosis. *J Am Heart Assoc*. 2017;6.
5. Cao Y, Singh V, Wang A, Zhang L, He T, Su H, et al. Meta-analysis of right ventricular function in patients with aortic stenosis after transfemoral aortic valve replacement or surgical aortic valve replacement. *Ther Adv Chronic Dis*. 2020;11:2040622320933775.
6. Mehdipour G, Chen S, Chatterjee S, Torkian P, Ben-Yehuda O, Leon MB, et al. Cardiac structural changes after transcatheter aortic valve replacement: systematic review and meta-analysis of cardiovascular magnetic resonance studies. *J Cardiovasc Magn Reson*. 2020;22:41.

7. Ren B, Spitzer E, Geleijnse ML, Zijlstra F, de Jaegere PPT, Van Mieghem NM, et al. Right ventricular systolic function in patients undergoing transcatheter aortic valve implantation: A systematic review and meta-analysis. *Int J Cardiol.* 2018;257:40-5.
8. Okada DR, Rahmouni HW, Herrmann HC, Bavaria JE, Forfia PR, Han Y. Assessment of right ventricular function by transthoracic echocardiography following aortic valve replacement. *Echocardiography.* 2014;31:552-7.
9. Quick S, Speiser U, Pfluecke C, Youssef A, Sveric K, Strasser R, et al. Aortic stenosis: right and left ventricular function in the early postprocedural phase. Comparison between transcatheter and surgical aortic valve implantation. *Acta Cardiol.* 2013;68:583-9.
10. Duncan AE, Sarwar S, Kateby Kashy B, Sonny A, Sale S, Alfirevic A, et al. Early Left and Right Ventricular Response to Aortic Valve Replacement. *Anesth Analg.* 2017;124:406-18.
11. Forsberg LM, Tamas E, Vanky F, Nielsen NE, Engvall J, Nylander E. Left and right ventricular function in aortic stenosis patients 8 weeks post-transcatheter aortic valve implantation or surgical aortic valve replacement. *Eur J Echocardiogr.* 2011;12:603-11.
12. Keyl C, Schneider J, Beyersdorf F, Ruile P, Siepe M, Pioch K, et al. Right ventricular function after aortic valve replacement: a pilot study comparing surgical and transcatheter procedures using 3D echocardiography. *Eur J Cardiothorac Surg.* 2016;49:966-71.
13. Reskovic Luksic V, Postolache A, Martinez C, Dulgheru R, Ilardi F, Tridetti J, et al. Global and regional myocardial function and outcomes after transcatheter aortic valve implantation for aortic stenosis and preserved ejection fraction. *J Cardiovasc Med (Hagerstown).* 2020;21:238-45.
14. Schattke S, Baldenhofer G, Prauka I, Zhang K, Laule M, Stangl V, et al. Acute regional improvement of myocardial function after interventional transfemoral aortic valve replacement in aortic stenosis: a speckle tracking echocardiography study. *Cardiovasc Ultrasound.* 2012;10:15.
15. Zhang K, Sheu R, Zimmerman NM, Alfirevic A, Sale S, Gillinov AM, et al. A Comparison of Global Longitudinal, Circumferential, and Radial Strain to Predict Outcomes After Cardiac Surgery. *J Cardiothorac Vasc Anesth.* 2019;33:1315-22.
16. Koschutnik M, Dannenberg V, Nitsche C, Dona C, Siller-Matula JM, Winter MP, et al. Right ventricular function and outcome in patients undergoing transcatheter aortic valve replacement. *Eur Heart J Cardiovasc Imaging.* 2021;22:1295-303.
17. Nieh CC, Teo AY, Soo WM, Lee GK, Singh D, Poh KK. Improvement in left ventricular function assessed by tissue Doppler imaging after aortic valve replacement for severe aortic stenosis. *Singapore Med J.* 2015;56:672-6.
18. Zhao Y, Lindqvist P, Nilsson J, Holmgren A, Naslund U, Henein MY. Trans-catheter aortic valve implantation—early recovery of left and preservation of right ventricular function. *Interact Cardiovasc Thorac Surg.* 2011;12:35-9.
19. Vahanian A, Beyersdorf F, Praz F, Milojevic M, Baldus S, Bauersachs J, et al. 2021 ESC/EACTS Guidelines for the management of valvular heart disease. *Eur J Cardiothorac Surg.* 2021;60:727-800.
20. Dimitriadis Z, Scholtz S, Ensminger S, Wiemer M, Fischbach T, Scholtz W, et al. Left ventricular adaptation after TAVI evaluated by conventional and speckle-tracking echocardiography. *Int J Cardiol.* 2017;228:633-7.
21. Forsberg LM, Tamas E, Vanky F, Engvall J, Nylander E. Differences in recovery of left and right ventricular function following aortic valve interventions: a longitudinal echocardiographic study in patients undergoing surgical, transapical or transfemoral aortic valve implantation. *Catheter Cardiovasc Interv.* 2013;82:1004-14.

22. Garg V, Ho JK, Vorobiof G. Changes in myocardial deformation after transcatheter and surgical aortic valve replacement. *Echocardiography*. 2017;34:603-13.
23. Kim SJ, Samad Z, Bloomfield GS, Douglas PS. A critical review of hemodynamic changes and left ventricular remodeling after surgical aortic valve replacement and percutaneous aortic valve replacement. *Am Heart J*. 2014;168:150-9 e1-7.
24. Meyer CG, Frick M, Lotfi S, Altiok E, Koos R, Kirschfink A, et al. Regional left ventricular function after transapical vs. transfemoral transcatheter aortic valve implantation analysed by cardiac magnetic resonance feature tracking. *Eur Heart J Cardiovasc Imaging*. 2014;15:1168-76.
25. Sandstede JJ, Beer M, Hofmann S, Lipke C, Harre K, Pabst T, et al. Changes in left and right ventricular cardiac function after valve replacement for aortic stenosis determined by cine MR imaging. *J Magn Reson Imaging*. 2000;12:240-6.
26. Vollema EM, Sugimoto T, Shen M, Tastet L, Ng ACT, Abou R, et al. Association of Left Ventricular Global Longitudinal Strain With Asymptomatic Severe Aortic Stenosis: Natural Course and Prognostic Value. *JAMA Cardiol*. 2018;3:839-47.
27. Halkin A, Steinvil A, Witberg G, Barsheshet A, Barkagan M, Assali A, et al. Mortality prediction following transcatheter aortic valve replacement: A quantitative comparison of risk scores derived from populations treated with either surgical or percutaneous aortic valve replacement. The Israeli TAVR Registry Risk Model Accuracy Assessment (IRRMA) study. *Int J Cardiol*. 2016;215:227-31.
28. Iung B, Laouenan C, Himbert D, Eltchaninoff H, Chevreul K, Donzeau-Gouge P, et al. Predictive factors of early mortality after transcatheter aortic valve implantation: individual risk assessment using a simple score. *Heart*. 2014;100:1016-23.
29. Al-Hijji MA, Zack CJ, Nkomo VT, Pislaru SV, Pelikka PA, Reeder GS, et al. Left ventricular remodeling and function after transapical versus transfemoral transcatheter aortic valve replacement. *Catheter Cardiovasc Interv*. 2019;94:738-44.
30. Caglayan HA, Kjonas D, Malm S, Schirmer H, Rosner A. Echocardiographic assessment of diastolic dysfunction in elderly patients with severe aortic stenosis before and after aortic valve replacement. *Cardiovasc Ultrasound*. 2021;19:32.
31. Leite S, Rodrigues S, Tavares-Silva M, Oliveira-Pinto J, Alaa M, Abdellatif M, et al. Afterload-induced diastolic dysfunction contributes to high filling pressures in experimental heart failure with preserved ejection fraction. *Am J Physiol Heart Circ Physiol*. 2015;309:H1648-54.
32. van Heerebeek L, Hamdani N, Falcao-Pires I, Leite-Moreira AF, Begieneman MP, Bronzwaer JG, et al. Low myocardial protein kinase G activity in heart failure with preserved ejection fraction. *Circulation*. 2012;126:830-9.
33. Schwartz LA, Rozenbaum Z, Ghantous E, Kramarz J, Biner S, Ghermezi M, et al. Impact of Right Ventricular Dysfunction and Tricuspid Regurgitation on Outcomes in Patients Undergoing Transcatheter Aortic Valve Replacement. *J Am Soc Echocardiogr*. 2017;30:36-46.
34. Musa TA, Uddin A, Fairbairn TA, Dobson LE, Steadman CD, Kidambi A, et al. Right ventricular function following surgical aortic valve replacement and transcatheter aortic valve implantation: A cardiovascular MR study. *Int J Cardiol*. 2016;223:639-44.
35. Kempny A, Diller GP, Kaleschke G, Orwat S, Funke A, Schmidt R, et al. Impact of transcatheter aortic valve implantation or surgical aortic valve replacement on right ventricular function. *Heart*. 2012;98:1299-304.
36. Sato K, Kumar A, Krishnaswamy A, Mick S, Desai MY, Griffin BP, et al. B-type natriuretic peptide is associated with remodeling and exercise capacity after transcatheter aortic valve replacement for aortic stenosis. *Clin Cardiol*. 2019;42:270-6.

Table 1: Patient characteristics

<i>n</i> total
Male
Age (y)
BMI (kg/m ²)
PCI pre TAVR
CAD
Angina
Previous CABG
New MI (90 days)
Perifer Vascular Disease
Cerebrovascular Disease
COPD
Cancer
Hypertension
Diabetes
Smoking
Cholesterol (mmol/l)
GFR pre (ml/min/1.73m ²)
GFR post (ml/min/1.73m ²)
Heart failure <2 weeks
CK MB post
Atrial fibrillation/flutter preop
Atrial ibrillation/flutter postop
Preoperative ventr pacing or LBBB
New postoperative ventricular pacing or LBBB
LogEuroScore ()
Significant difference towards * TA-TAVR; + TF-TAVR; ++ SAVR. TAVR: Transcatheter aortic valve replacement; TA: tr

Table 2: Clinical and LV systolic functional parameters by conventional echocardiography and strain rate imaging

	Procedure	n	Pre Mean ±SD	Post Mean ±SD	<i>p</i> -value
NT-proBNP (pg/ml)	TA-TAVR	22	4306±7433	2001±2561	0.117
	TF-TAVR	41	5030±8464	1856±3389	0.004
	SAVR	35	1092±1085	761±1064	0.094
	SAVR+CABG	17	1151±1479	712±787	0.078
NYHA class III-IV (n/%)	TA-TAVR	27	24 (89%)	1 (4%)	<0.001
	TF-TAVR	41	39 (95%)	3 (7%)	<0.001
	SAVR	36	25 (69%)	1 (3%)	<0.001
	SAVR+CABG	24	17 (71%)	1 (4%)	<0.001
EF Simpson Biplane (%)	TA-TAVR	27	52 ±13	50 ±9	0.390
	TF-TAVR	43	52 ±15	55 ±9	0.120
	SAVR	36	56 ±12	60 ±9	0.054
	SAVR+CABG	24	55 ±9	58 ±9	0.179

LV volume diastole (ml)	TA-TAVR	27	139 ±60	123 ±32	0.181
	TF-TAVR	43	121 ±52	101 ±40	<0.001
	SAVR	36	121 ±32	99 ±27	<0.001
	SAVR+CABG	24	121 ±42	94 ±28	<0.001
PSV mitral ring (cm/s)	TA-TAVR	27	5.3 ±1.3	5.8 ±1.2	0.022
	TF-TAVR	43	5.4 ±1.2	6.5 ±1.4	<0.001
	SAVR	34	6.5±1.4	7.8 ±2.8	0.011
	SAVR+CABG	24	6.2 ±1.1	6.9 ±2.5	0.089
Longitudinal Strain ET (%)	TA-TAVR	26	-8.2 ±2.2	-8.5 ±2.8	0.221
	TF-TAVR	42	-9.4 ±2.3	-9.9 ±2.6	<0.001
	SAVR	34	-11.0 ±2.3	-11.8 ±2.5	<0.001
	SAVR+CABG	24	-9.3 ±2.6	-9.3 ±3.2	0.817
Circumferential Strain ET (%)	TA-TAVR	18	-13.7 ±3.4	-14.5 ±5.0	0.257
	TF-TAVR	26	-13.1 ±3.5	-14.1 ±3.7	<0.001
	SAVR	23	-15.0±3.2	-15.3 ±3.6	0.353
	SAVR+CABG	15	-14.4 ±4.5	-14.5 ±4.7	0.737
Longitudinal SR ET (1/s)	TA-TAVR	26	0.52 ±0.12	0.53 ±0.14	0.373
	TF-TAVR	42	-0.57 ±0.12	-0.60 ±0.14	0.011
	SAVR	34	-0.67 ±0.11	-0.71 ±0.13	<0.001
	SAVR+CABG	24	-0.61 ±0.12	-0.64 ±0.14	0.029
Circumferential SR ET (1/s)	TA-TAVR	18	0.82 ±0.25	0.89 ±0.41	0.179
	TF-TAVR	26	-0.75 ±0.19	-0.84 ±0.22	<0.001
	SAVR	23	-0.88 ±0.16	-0.89 ±0.22	0.570
	SAVR+CABG	15	-0.88 ±0.33	-0.97 ±0.40	0.036

No significantly differing pre-to post AVR changes between groups. TAVR: Transcatheter aortic valve replacement; TA: transapical; TF: transfemoral; SAVR: surgical aortic valve replacement; CABG: coronary artery bypass graft; NT-proBNP: N-terminal pro-Brain Natriuretic Peptid; NYHA: New York Heart Association; EF: ejection fraction; LV: left ventricular; PSV: peak systolic velocity; ET: ejection time; SR: strain rate	No significantly differing pre-to post AVR changes between groups. TAVR: Transcatheter aortic valve replacement; TA: transapical; TF: transfemoral; SAVR: surgical aortic valve replacement; CABG: coronary artery bypass graft; NT-proBNP: N-terminal pro-Brain Natriuretic Peptid; NYHA: New York Heart Association; EF: ejection fraction; LV: left ventricular; PSV: peak systolic velocity; ET: ejection time; SR: strain rate	No significantly differing pre-to post AVR changes between groups. TAVR: Transcatheter aortic valve replacement; TA: transapical; TF: transfemoral; SAVR: surgical aortic valve replacement; CABG: coronary artery bypass graft; NT-proBNP: N-terminal pro-Brain Natriuretic Peptid; NYHA: New York Heart Association; EF: ejection fraction; LV: left ventricular; PSV: peak systolic velocity; ET: ejection time; SR: strain rate	No significantly differing pre-to post AVR changes between groups. TAVR: Transcatheter aortic valve replacement; TA: transapical; TF: transfemoral; SAVR: surgical aortic valve replacement; CABG: coronary artery bypass graft; NT-proBNP: N-terminal pro-Brain Natriuretic Peptid; NYHA: New York Heart Association; EF: ejection fraction; LV: left ventricular; PSV: peak systolic velocity; ET: ejection time; SR: strain rate	No significantly differing pre-to post AVR changes between groups. TAVR: Transcatheter aortic valve replacement; TA: transapical; TF: transfemoral; SAVR: surgical aortic valve replacement; CABG: coronary artery bypass graft; NT-proBNP: N-terminal pro-Brain Natriuretic Peptid; NYHA: New York Heart Association; EF: ejection fraction; LV: left ventricular; PSV: peak systolic velocity; ET: ejection time; SR: strain rate	No significantly differing pre-to post AVR changes between groups. TAVR: Transcatheter aortic valve replacement; TA: transapical; TF: transfemoral; SAVR: surgical aortic valve replacement; CABG: coronary artery bypass graft; NT-proBNP: N-terminal pro-Brain Natriuretic Peptid; NYHA: New York Heart Association; EF: ejection fraction; LV: left ventricular; PSV: peak systolic velocity; ET: ejection time; SR: strain rate
---	---	---	---	---	---

Table 3: LV diastolic functional parameters before and after different procedures of aortic valve replacement

	Procedure	n	Pre Mean $\pm$ SD	Post Mean $\pm$ SD	<i>p</i> -value
E-Wave Velocity (cm/s)	TA-TAVR	27	101 $\pm$ 38	93 $\pm$ 40	0.306
	TF-TAVR	43	98 $\pm$ 26	93 $\pm$ 30	0.399
	SAVR	36	90 $\pm$ 22	93 $\pm$ 25	0.452
	SAVR+CABG	24	97 $\pm$ 30	101 $\pm$ 30	0.345
E/A ()	TA-TAVR	17	1.2 $\pm$ 0.7	0.8 $\pm$ 0.6	0.147

MV e' (cm/s)	TF-TAVR	29	1.0 ±0.7	0.9 ±0.2	0.149
	SAVR	29	0.7 ±0.2	0.9 ±0.2	0.013
	SAVR+CABG	22	0.87 ±0.24	0.93 ±0.32	0.350
	TA-TAVR	27	5.7 ±1.6	5.9 ±1.2	0.552
Pulmonary veins: SFF (%)	TF-TAVR	43	5.4 ±1.6	6.0 ±1.6	0.006
	SAVR	34	6.2 ±3.8	7.2 ±1.7	0.108
	SAVR+CABG	24	5.5 ±1.6	6.9 ±2.5	0.024
	TA-TAVR	25	42 ±22	44 ±15	0.695
LA volume index (ml)	TF-TAVR	38	50 ±15	50 ±13	0.796
	SAVR	35	55 ±16	52 ±14	0.075
	SAVR+CABG	24	59 ±9	52 ±12	0.003
	TA-TAVR	27	58 ±20	60 ±39	0.797
IVRT (ms)	TF-TAVR	42	55 ±28	51 ±22	0.259
	SAVR	35	44 ±15	42 ±12	0.300
	SAVR+CABG	24	45 ±16	42 ±15	0.236
	TA-TAVR	27	64±30	87 ±35	0.036
	TF-TAVR	43	71 ±60	88 ±47	0.123
	SAVR	36	72 ±34	82 ±26	0.068
	SAVR+CABG	24	72 ±38	103 ±46	0.024

Significant difference towards *	Significant difference towards *	Significant difference towards *	Significant difference towards *	Significant difference towards *	Significant difference towards *
TA-TAVR; + TF-TAVR; ++ SAVR. TAVR:	TA-TAVR; + TF-TAVR; ++ SAVR. TAVR:	TA-TAVR; + TF-TAVR; ++ SAVR. TAVR:	TA-TAVR; + TF-TAVR; ++ SAVR. TAVR:	TA-TAVR; + TF-TAVR; ++ SAVR. TAVR:	TA-TAVR; + TF-TAVR; ++ SAVR. TAVR:
Transcatheter aortic valve replacement; LV: left ventricular; TA: transapical; TF: transfemoral; SAVR: surgical aortic valve replacement; CABG: coronary artery bypass graft; SFF: systolic filling fraction; LA: left atrium; IVRT: isovolumic relaxation time	Transcatheter aortic valve replacement; LV: left ventricular; TA: transapical; TF: transfemoral; SAVR: surgical aortic valve replacement; CABG: coronary artery bypass graft; SFF: systolic filling fraction; LA: left atrium; IVRT: isovolumic relaxation time	Transcatheter aortic valve replacement; LV: left ventricular; TA: transapical; TF: transfemoral; SAVR: surgical aortic valve replacement; CABG: coronary artery bypass graft; SFF: systolic filling fraction; LA: left atrium; IVRT: isovolumic relaxation time	Transcatheter aortic valve replacement; LV: left ventricular; TA: transapical; TF: transfemoral; SAVR: surgical aortic valve replacement; CABG: coronary artery bypass graft; SFF: systolic filling fraction; LA: left atrium; IVRT: isovolumic relaxation time	Transcatheter aortic valve replacement; LV: left ventricular; TA: transapical; TF: transfemoral; SAVR: surgical aortic valve replacement; CABG: coronary artery bypass graft; SFF: systolic filling fraction; LA: left atrium; IVRT: isovolumic relaxation time	Transcatheter aortic valve replacement; LV: left ventricular; TA: transapical; TF: transfemoral; SAVR: surgical aortic valve replacement; CABG: coronary artery bypass graft; SFF: systolic filling fraction; LA: left atrium; IVRT: isovolumic relaxation time

Table 4: RV geometry and functional parameters

	Procedure	n	Pre Mean $\pm$ SD	Post Mean $\pm$ SD	p-value
RVOT VTI (cm)	TA-TAVR	24	14.5 $\pm$ 3.9	15.1 $\pm$ 9.3	0.776
	TF-TAVR	39	14.8 $\pm$ 4.6	16.4 $\pm$ 9.1	0.281
	SAVR	36	16.1 $\pm$ 6.1	16.8 $\pm$ 7.2	0.653
	SAVR+CABG	23	17.3 $\pm$ 10.0	14.9 $\pm$ 2.8	0.248
RV FAC (%)	TA-TAVR	27	30 $\pm$ 14	35 $\pm$ 9	0.033
	TF-TAVR	43	32 $\pm$ 15	33 $\pm$ 11	0.684
	SAVR	36	34.8 $\pm$ 13.7	34.1 $\pm$ 13.7	0.841
	SAVR+CABG	22	35.6 $\pm$ 18.0	37.8 $\pm$ 9.8	0.494
TV E (cm/s)	TA-TAVR	27	45 $\pm$ 12	48 $\pm$ 10	0.406
	TF-TAVR	41	44 $\pm$ 12	46 $\pm$ 19	0.627
	SAVR	36	45.5 $\pm$ 9.1	48.1 $\pm$ 9.4	0.158
	SAVR+CABG	24	40.4 $\pm$ 9.2	51.5 $\pm$ 10.5	<0.001
TV A (cm/s)	TA-TAVR	17	41 $\pm$ 6	42 $\pm$ 6	0.382
	TF-TAVR	27	44 $\pm$ 9	42 $\pm$ 11	0.267
	SAVR	29	44.0 $\pm$ 12.3	41.1 $\pm$ 7.9	0.230

No	SAVR+CABG	22	44.2 ±12.2	41.7 ±9.8	0.411
significantly	significantly	No	significantly	significantly	No
differing pre-to	differing pre-to	significantly	differing pre-to	differing pre-to	significantly
post AVR	post AVR	differing pre-to	post AVR	post AVR	differing pre-to
changes	changes	post AVR	changes	changes	post AVR
between	between	changes	between	between	changes
groups.	groups.	between	groups.	groups.	between
TAVR:	TAVR:	groups.	TAVR:	TAVR:	groups.
Transcatheter	Transcatheter	TAVR:	Transcatheter	Transcatheter	TAVR:
aortic valve	aortic valve	Transcatheter	aortic valve	aortic valve	Transcatheter
replacement;	replacement;	aortic valve	replacement;	replacement;	aortic valve
TA:	TA:	replacement;	TA:	TA:	replacement;
transapical;	transapical;	TA:	transapical;	transapical;	TA:
TF:	TF:	transapical;	TF:	TF:	transapical;
transfemoral;	transfemoral;	TF:	transfemoral;	transfemoral;	TF:
SAVR:	SAVR:	transfemoral;	SAVR:	SAVR:	transfemoral;
surgical aortic	surgical aortic	SAVR:	surgical aortic	surgical aortic	SAVR:
valve	valve	surgical aortic	valve	valve	surgical aortic
replacement;	replacement;	valve	replacement;	replacement;	valve
CABG:	CABG:	replacement;	CABG:	CABG:	replacement;
coronary	coronary	CABG:	coronary	coronary	CABG:
artery bypass	artery bypass	coronary	artery bypass	artery bypass	coronary
graft; RVOT	graft; RVOT	artery bypass	graft; RVOT	graft; RVOT	artery bypass
VTI: right	VTI: right	graft; RVOT	VTI: right	VTI: right	graft; RVOT
ventricular	ventricular	VTI: right	ventricular	ventricular	VTI: right
outflow tract	outflow tract	ventricular	outflow tract	outflow tract	ventricular
velocity time	velocity time	outflow tract	velocity time	velocity time	outflow tract
integral; RV	integral; RV	velocity time	integral; RV	integral; RV	velocity time
FAC: right	FAC: right	integral; RV	FAC: right	FAC: right	integral; RV
ventricular	ventricular	FAC: right	ventricular	ventricular	FAC: right
fractional area	fractional area	ventricular	fractional area	fractional area	ventricular
change; TV:	change; TV:	fractional area	change; TV:	change; TV:	fractional area
tricuspid valve	tricuspid valve	change; TV:	tricuspid valve	tricuspid valve	change; TV:
		tricuspid valve			tricuspid valve

Table 5: Bland Altman Limits of Agreement for myocardial peak systolic strain- and SR measurements

	Mean	SD
Intra-observer variability		
Longitudinal Strain ET (%)	1.8	±3.0
Circumferential Strain ET (%)	1.4	±2.8
Longitudinal SR ET (1/s)	0.08	±0.19
Circumferential SR ET (1/s)	0.07	±0.21
Inter-observer variability		
Longitudinal Strain ET (%)	2.0	±3.9
Circumferential Strain ET (%)	2.2	±3.2
Longitudinal SR ET (1/s)	0.11	±0.21
Circumferential SR ET (1/s)	0.13	±0.24
ET: during ejection time; SR: strain rate	ET: during ejection time; SR: strain rate	ET: during ejection time; SR: strain

Figure legends

Figure 1: Left ventricular systolic and diastolic functional parameters pre and post aortic valve replacement. Comparison of the different treatment groups.

TAVR, transcatheter aortic valve replacement; SAVR, surgical aortic valve replacement; CABG, coronary artery bypass graft; MAPSE, mitral annular plane systolic excursion.

Figure 2: Right ventricular geometrical and systolic functional parameters pre- and post-aortic valve replacement. Comparison of the different treatment groups.

TAVR, transcatheter aortic valve replacement; SAVR, surgical aortic valve replacement; CABG, coronary artery bypass graft; TAPSE, tricuspid annular plane systolic excursion.

Figure 3: Right ventricular diastolic parameters pre- and post-aortic valve replacement. Comparison of the different treatment groups.

TAVR, transcatheter aortic valve replacement; SAVR, surgical aortic valve replacement; CABG, coronary artery bypass graft; TDI, tissue Doppler imaging





