

Geometry of solutions of the geometric curve flows in space

Shiping Zhong¹, Zehui Zhao¹, and Xinjie Wan¹

¹School of Mathematics and Computer Sciences Gannan Normal University Ganzhou
341000 PR China

November 4, 2022

Abstract

In this study, we aim at investigating the geometry of surfaces corresponding to the geometry of solutions of the geometric curve flows in Euclidean 3-space $\mathbb{R}^3$ considering the Frenet frame. In particular, we express some geometric properties and some characterizations of u -parameter curves and t -parameter curves of some trajectory surfaces including the Hasimoto surface, shortening trajectory surface, minimal trajectory surface, τ -normal trajectory surface in $\mathbb{R}^3$.

ARTICLE TYPE

Geometry of solutions of the geometric curve flows in space[†]

Shiping Zhong* | Zehui Zhao | Xinjie Wan

¹School of Mathematics and Computer Sciences, Gannan Normal University, Ganzhou, 341000, P.R. China

Correspondence

*Corresponding author Email: zhongshiping@gnnu.edu.cn

Summary

In this study, we aim at investigating the geometry of surfaces corresponding to the geometry of solutions of the geometric curve flows in Euclidean 3-space $\mathbb{R}^3$ considering the Frenet frame. In particular, we express some geometric properties and some characterizations of u -parameter curves and t -parameter curves of some trajectory surfaces including the Hasimoto surface, shortening trajectory surface, minimal trajectory surface, $\sqrt{\tau}$ -normal trajectory surface in $\mathbb{R}^3$.

KEYWORDS:

The geometric curve flows in space, Hasimoto surface, Shortening trajectory surface, Minimal trajectory surface, $\sqrt{\tau}$ -normal trajectory surface

1 | INTRODUCTION

This article investigates geometry of Surfaces defined as trajectories of evolving geometric curves flow in Euclidean 3-space $\mathbb{R}^3$. In recent years, the theory of surfaces with the connection of the geometric curve flows in different spaces and integrable non-linear equations is a subject of research attention (see e.g. ^{1,2,3,4,5,6,7,8,9,10,11,12,13,14,15,16}) and finding geometric curves flows which generate various types of surfaces may expand our understanding of their geometric and topological properties, in order to a deep understanding of the physical world (see e.g. ^{17,18,19,20}).

Hasimoto¹⁰ in 1972 showed the connection between the binormal flow and the nonlinear Schrödinger equation. Surfaces generated by the binormal flow, referred to as Hasimoto surfaces, have been previously considered in ^{1,12}. For Hashimoto surfaces, A lot of researches were done using the different frames (such as the Frenet frame¹, the Bishop frame¹³, the Darboux frame^{7,9}, the quasi-frame¹¹, the modified orthogonal frame⁸, the hybrid frame¹⁶ and so on) in Euclidean 3-space¹, Minkowski 3-space^{5,14,15}, Galilean 3-space⁶ and pseudo-Galilean 3-space². Trajectory surfaces have been studied for the special case of inextensible flows in²¹, curves flows of elastic rods in²², and the curve shortening flow in^{23,24,25}. Recently, J. Minarčík and M. Beneš²⁶ introduced minimal surface generating flow for space curves of non-vanishing torsion in Euclidean 3-space $\mathbb{R}^3$, and analyzed some properties of space curves evolved by the minimal surface generating flow.

The aim of this paper is to study the geometry of trajectory surfaces corresponding to the geometry of solutions of the geometric curve flows in Euclidean 3-space. The paper is organized as follows. Section 2 contains an introduction to geometric curves flow and recalls the notion of trajectory surfaces, including the Hasimoto surface, shortening trajectory surface, minimal trajectory surface, $\sqrt{\tau}$ -normal trajectory surface in $\mathbb{R}^3$. In Section 3, we investigate the geometric properties of some trajectory surfaces via the Frenet frame. Also, the characterization of the u -parameter curves and t -parameter curves of the trajectory surfaces is examined. In section 4, we draw the main conclusions, as well as some questions that we postpone for the future.

[†]This is an example for title footnote.

2 | TRAJECTORY SURFACES FOR THE GEOMETRIC CURVE FLOWS IN SPACE

In this section, we define the necessary notation for the parametric space curves in motion, recall the governing equations for a general geometric flow of curves in $\mathbb{R}^{3,27}$ and introduce the notion of trajectory surfaces generated by this general motion law, and finally we obtain the Gaussian curvature, mean curvature, principal curvatures, the geodesic torsion, geodesic curvature, and normal curvature of the trajectory surfaces.

Assume the family of evolving curves is parametrized as follows:

$$\{\Gamma_t\}_{t \in [0, t_{\max})} = \{\mathbf{X}(u, t), u \in \mathbb{S}^1, t \in [0, t_{\max})\},$$

where $\mathbf{X} = \mathbf{X}(u, t) : \mathbb{S}^1 \times [0, t_{\max}) \rightarrow \mathbb{R}^3$ is a smooth mapping, $t_{\max} > 0$ is the terminal time and $\mathbb{S}^1 = \mathbb{R}/2\pi\mathbb{Z}$ is a unit circle. Then the unit arc-length parametrization s is given by $ds = g du$, where $g = |\partial_u \mathbf{X}|$. The unit tangent vector is given by $\mathbf{T} = \partial_s \mathbf{X}$. In the case when the curvature $\kappa = |\mathbf{T} \times \partial_s \mathbf{T}| > 0$ is strictly positive, we can define the so-called Frenet frame. It means that the unit normal and binormal vectors $\mathbf{N}$ and $\mathbf{B}$ can be uniquely defined as follows:

$$\mathbf{N} = \kappa^{-1} \partial_s \mathbf{T}, \quad \mathbf{B} = \mathbf{T} \times \mathbf{N}.$$

These unit vectors satisfy the following identities:

$$\mathbf{B} = \mathbf{T} \times \mathbf{N}, \quad \mathbf{T} = \mathbf{N} \times \mathbf{B}, \quad \mathbf{N} = \mathbf{B} \times \mathbf{T},$$

and the Frenet-Serret formulae:

$$\frac{d}{ds} \begin{pmatrix} \mathbf{T} \\ \mathbf{N} \\ \mathbf{B} \end{pmatrix} = \begin{pmatrix} 0 & \kappa & 0 \\ -\kappa & 0 & \tau \\ 0 & -\tau & 0 \end{pmatrix} \begin{pmatrix} \mathbf{T} \\ \mathbf{N} \\ \mathbf{B} \end{pmatrix},$$

where τ is the torsion of a curve. For $\kappa > 0$ the torsion τ is given by

$$\tau = \kappa^{-2} (\mathbf{T} \times \partial_s \mathbf{T}) \cdot \partial_s^2 \mathbf{T} = \kappa^{-2} (\partial_s \mathbf{X} \times \partial_s^2 \mathbf{X}) \cdot \partial_s^3 \mathbf{X}.$$

The time evolution of $\{\Gamma_t\}_{t \in [0, t_{\max})}$ is given by the geometric flow in the form of the following initial-value problem for the parametrization $\mathbf{X} = \mathbf{X}(u, t)$ in $\mathbb{R}^3$

$$\begin{aligned} \partial_t \mathbf{X} &= v_N \mathbf{N} + v_B \mathbf{B} + v_T \mathbf{T} & \text{in } \mathbb{S}^1 \times [0, t_{\max}), \\ \mathbf{X}|_{t=0} &= \mathbf{X}_0 & \text{in } \mathbb{S}^1, \end{aligned} \quad (1)$$

where $\mathbf{X}_0$ is the parametrization for the initial curve Γ_0 .

Firstly, we recall that the dynamic equations for unit vectors $\mathbf{N}, \mathbf{B}, \mathbf{T}$ and the local geometric quantities (i.e., the curvature κ , torsion τ) during the motion given by (1) in the following (see²⁷ for details).

- The unit vectors $\mathbf{N}, \mathbf{B}, \mathbf{T}$ forming the Frenet frame satisfy the evolution equations:

$$\begin{aligned} \partial_t \mathbf{T} &= (\partial_s v_N + \kappa v_T - \tau v_B) \mathbf{N} + (\partial_s v_B + \tau v_N) \mathbf{B}, \\ \kappa \partial_t \mathbf{N} &= -\kappa (\partial_s v_N + \kappa v_T - \tau v_B) \mathbf{T} + (\partial_s^2 v_B + v_N \partial_s \tau + 2\tau \partial_s v_N + \tau (\kappa v_T - \tau v_B)) \mathbf{B}, \\ \kappa \partial_t \mathbf{B} &= -\kappa (\partial_s v_B + \tau v_N) \mathbf{T} - (\partial_s^2 v_B + v_N \partial_s \tau + 2\tau \partial_s v_N + \tau (\kappa v_T - \tau v_B)) \mathbf{N}. \end{aligned} \quad (2)$$

- The local length element $g = |\partial_u \mathbf{X}|$ satisfy $\partial_t g = (-\kappa v_N + \partial_s v_T)g$.
- The curvature κ and torsion τ (for $\kappa(s, t) > 0$) satisfy the following system of evolution partial differential equations

$$\partial_t \kappa = \partial_s^2 v_N + \kappa^2 v_N + v_T \partial_s \kappa - \partial_s (\tau v_B) - \tau \partial_s v_B - \tau^2 v_N, \quad (3)$$

$$\begin{aligned} \partial_t \tau &= \kappa (\partial_s v_B + \tau v_N) + \partial_s (\kappa^{-1} (\partial_s^2 v_B + v_N \partial_s \tau + 2\tau \partial_s v_N + \tau (\kappa v_T - \tau v_B))) \\ &\quad + \tau (\kappa v_N - \partial_s v_T). \end{aligned} \quad (4)$$

Definition 1.^{21,26} (Trajectory surface) For given velocities v_T, v_N and v_B , an initial curve Γ_0 and terminal time $t_{\max}$, we define the trajectory surface $\sum_{t_{\max}}$ as $\sum_{t_{\max}} := \bigcup_{t \in [0, t_{\max})} \Gamma_t$.

Trajectory surfaces have been studied in^{1,21,23,26} for the following cases, such as the binormal flow (i.e., $v_T = v_N = 0, v_B = \kappa$), the curve shortening flow (i.e., $v_T = v_B = 0, v_N = \kappa$), minimal surface generating flow (i.e., $v_T = v_B = 0, v_N = \tau^{-\frac{1}{2}}$), inextensible flows (i.e., geometric flows with $\frac{\partial g}{\partial t} = 0$).

- (i) A Hasimoto surface Σ_1 is the surface traced out by a curve $\mathbf{X}(u, t)$ in $\mathbb{R}^3$ as it evolves over time according to this evolution equation:

$$\partial_t \mathbf{X} = \kappa \mathbf{B}, \quad (5)$$

which motion law is the binormal flow, also known as the localized induction approximation of the vortex filament flow, which has applications in the incompressible flows (see e.g.¹).

- (ii) A shortening trajectory surface Σ_2 is the surface traced out by a family of curves $\{\Gamma_t\}_{t \in [0, t_{max}]}$ in $\mathbb{R}^3$ as it evolves over time according to the the curve shortening flow:

$$\begin{aligned} \partial_t \mathbf{X} &= \kappa \mathbf{N}, & \text{in } \mathbb{S}^1 \times [0, t_{max}), \\ \mathbf{X}|_{t=0} &= \mathbf{X}_0, & \text{in } \mathbb{S}^1, \end{aligned} \quad (6)$$

where $\mathbf{X}_0$ is the parametrization for the initial curve Γ_0 in $\mathbb{R}^3$. Moreover, if the curve approaches singularity as t goes to t_{max} , the shortening trajectory surface will be an embedded disc without a point in the center of the original sphere (see e.g.²³).

- (iii) A minimal trajectory surface Σ_3 is the surface traced out by a family of curves $\{\Gamma_t\}_{t \in [0, t_{max}]}$ in $\mathbb{R}^3$ as it evolves over time according to the minimal surface generating flow (see²⁶):

$$\begin{aligned} \partial_t \mathbf{X} &= \tau^{-\frac{1}{2}} \mathbf{N}, & \text{in } \mathbb{S}^1 \times [0, t_{max}), \\ \mathbf{X}|_{t=0} &= \mathbf{X}_0, & \text{in } \mathbb{S}^1, \end{aligned} \quad (7)$$

where $\mathbf{X}_0$ is the parametrization for the initial curve Γ_0 in $\mathbb{R}^3$ with positive curvature and torsion.

- (iv) A $\sqrt{\tau}$ -normal trajectory surface Σ_4 is the surface traced out by a family of curves $\{\Gamma_t\}_{t \in [0, t_{max}]}$ in $\mathbb{R}^3$ as it evolves over time according to the $\sqrt{\tau}$ -normal geometric flow:

$$\begin{aligned} \partial_t \mathbf{X} &= \sqrt{\tau} \mathbf{N}, & \text{in } \mathbb{S}^1 \times [0, t_{max}), \\ \mathbf{X}|_{t=0} &= \mathbf{X}_0, & \text{in } \mathbb{S}^1, \end{aligned} \quad (8)$$

where $\mathbf{X}_0$ is the parametrization for the initial curve Γ_0 in $\mathbb{R}^3$ with positive curvature and nonnegative torsion.

The first and the second fundamental forms of the surface $\Sigma_{t_{max}}$ generated by (1) are given by

$$\begin{aligned} I &\equiv \begin{pmatrix} \mathcal{E} & \mathcal{F} \\ \mathcal{F} & \mathcal{G} \end{pmatrix} = \begin{pmatrix} \|\partial_u \mathbf{X}\|^2 & \langle \partial_u \mathbf{X}, \partial_t \mathbf{X} \rangle \\ \langle \partial_t \mathbf{X}, \partial_u \mathbf{X} \rangle & \|\partial_t \mathbf{X}\|^2 \end{pmatrix}, \\ II &\equiv \begin{pmatrix} \mathcal{L} & \mathcal{M} \\ \mathcal{M} & \mathcal{N} \end{pmatrix} = \begin{pmatrix} \langle \partial_u^2 \mathbf{X}, n \rangle & \langle \partial_u \partial_t \mathbf{X}, n \rangle \\ \langle \partial_t \partial_u \mathbf{X}, n \rangle & \langle \partial_t^2 \mathbf{X}, n \rangle \end{pmatrix}, \end{aligned}$$

where the unit normal vector n of the trajectory surface $\Sigma_{t_{max}}$ can be expressed as

$$n = \frac{\partial_u \mathbf{X} \times \partial_t \mathbf{X}}{\|\partial_u \mathbf{X} \times \partial_t \mathbf{X}\|} = \frac{-v_B \mathbf{N} + v_N \mathbf{B}}{\sqrt{v_B^2 + v_N^2}}.$$

Using evolution equations (2)-(4), the elements of the first fundamental form I read

$$\mathcal{E} = g^2, \quad \mathcal{F} = g v_T, \quad \mathcal{G} = v_N^2 + v_B^2 + v_T^2, \quad (9)$$

and we give the elements of I :

$$\begin{aligned} \mathcal{L} &= -\frac{g^2 \kappa v_B}{\sqrt{v_B^2 + v_N^2}}, \\ \mathcal{M} &= g \left(\frac{-v_B \partial_s v_N + v_N \partial_s v_B - \kappa v_T v_B}{\sqrt{v_B^2 + v_N^2}} + \tau \sqrt{v_B^2 + v_N^2} \right), \\ \mathcal{N} &= \frac{v_N \partial_t v_B - v_B \partial_t v_N + v_T [v_N (\partial_s v_B + \tau v_N) - v_B (\partial_s v_N + \kappa v_T - \tau v_B)]}{\sqrt{v_B^2 + v_N^2}} \\ &\quad + \kappa^{-1} \sqrt{v_B^2 + v_N^2} (\partial_s^2 v_B + v_N \partial_s \tau + 2\tau \partial_s v_N + \kappa \tau v_T - \tau^2 v_B). \end{aligned} \quad (10)$$

The Gaussian curvature K , mean curvature H , and principal curvatures $P_i, i = 1, 2$ of the trajectory surface $\sum_{t_{max}}$ as

$$K = \frac{\det II}{\det I} = \frac{\mathcal{L}\mathcal{N} - \mathcal{M}^2}{\mathcal{E}\mathcal{G} - \mathcal{F}^2}, \quad H = \frac{1}{2}\text{Tr}(II(I)^{-1}) = \frac{\mathcal{L}\mathcal{G} - 2\mathcal{M}\mathcal{F} + \mathcal{N}\mathcal{E}}{2(\mathcal{E}\mathcal{G} - \mathcal{F}^2)},$$

$$P_1 = H + \sqrt{H^2 - K}, \quad P_2 = H - \sqrt{H^2 - K},$$

respectively. By Eqs. (9) and Eqs. (10), we can write K and H in terms of the functions v_T, v_N, v_B, κ and τ as

$$K = \frac{\kappa v_B (v_B \partial_t v_N - v_N \partial_t v_B + v_T (-v_N (\partial_s v_B + \tau v_N) + v_B (\partial_s v_N + \kappa v_T - \tau v_B)))}{(v_B^2 + v_N^2)^2}$$

$$- \frac{v_B (\partial_s^2 v_B + 2\tau \partial_s v_N + \partial_s \tau \partial_s v_N + \kappa \tau v_T - \tau^2 v_B)}{v_B^2 + v_N^2} - \frac{(v_N \partial_s v_B - v_B \partial_s v_N - \kappa v_T v_B)^2}{(v_B^2 + v_N^2)^2}$$

$$- \frac{2\tau (v_N \partial_s v_B - v_B \partial_s v_N - \kappa v_T v_B)}{v_B^2 + v_N^2} - \tau^2,$$

$$H = - \frac{\kappa v_B (v_N^2 + v_B^2 + v_T^2)}{2(v_N^2 + v_B^2)^{\frac{3}{2}}} - v_T \left(\frac{v_N \partial_s v_B - v_B \partial_s v_N - \kappa v_B v_T}{(v_N^2 + v_B^2)^{\frac{3}{2}}} + \tau \right)$$

$$+ \frac{v_N \partial_t v_B - v_B \partial_t v_N + v_T (v_N (\partial_s v_B + \tau v_N) - v_B (\partial_s v_N + \kappa v_T - \tau v_B))}{2(v_N^2 + v_B^2)^{\frac{3}{2}}}$$

$$+ \frac{\partial_s^2 v_B + 2\tau \partial_s v_N + v_N \partial_s \tau + \kappa \tau v_T - \tau^2 v_B}{2\kappa \sqrt{v_B^2 + v_N^2}}.$$

For a regular curve $\gamma(x)$ on trajectory surface $\sum_{t_{max}}$ in $\mathbb{R}^3$, the geodesic torsion $\mathcal{T}_g$, geodesic curvature $\mathcal{K}_g$, and normal curvature $\mathcal{K}_n$ are given by

$$\mathcal{T}_g = \frac{\langle \dot{\gamma}, n \times \dot{\gamma} \rangle}{||\dot{\gamma}||^2}, \quad \mathcal{K}_g = \frac{\langle \ddot{\gamma}, n \times \dot{\gamma} \rangle}{||\dot{\gamma}||^2}, \quad \mathcal{K}_n = \frac{\langle \ddot{\gamma}, n \rangle}{||\dot{\gamma}||^2},$$

respectively, where $\dot{\gamma} = \frac{\partial \gamma}{\partial x}$, $\ddot{\gamma} = \frac{\partial^2 \gamma}{\partial x^2}$, $\dot{n} = \frac{\partial n}{\partial x}$.

Now, we can write the following important definitions:

Definition 2. ²⁸

- A regular surface $\sum$ is a developable (flat) surface if its Gaussian curvature $K \equiv 0$, whereas it is a minimal surface if its mean curvature $H \equiv 0$;
- For a regular curve $\gamma(t)$ on a trajectory surface $\sum_{t_{max}}$, the following facts are well-known:
 - 1) $\gamma(t)$ is a principal line if and only if the geodesic torsion $\mathcal{T}_g \equiv 0$;
 - 2) $\gamma(t)$ is an asymptotic line if and only if the normal curvature $\mathcal{K}_n \equiv 0$;
 - 3) $\gamma(t)$ is a geodesic curve if and only if the geodesic curvature $\mathcal{K}_g \equiv 0$.

Definition 3. The family of all u -parameter curves and the family of all t -parameter curves of a the trajectory surface $\sum_{t_{max}}$ in $\mathbb{R}^3$ are denoted by Λ^u and Ω^t , respectively.

Using evolution equations (2)-(4), we obtain the geodesic torsion ${}^t\mathcal{T}_g$, geodesic curvature ${}^t\mathcal{K}_g$, and normal curvature ${}^t\mathcal{K}_n$ of the t parameter curves $\mathbf{X}(u, t)$ on trajectory surface $\sum_{t_{max}}$ as

$${}^t\mathcal{T}_g = \frac{\langle \dot{\mathbf{X}}, n \times \dot{\mathbf{X}} \rangle}{||\dot{\mathbf{X}}||^2} = \frac{\rho[(v_N^2 + v_B^2)n_1 - (v_B n_3 + v_N n_2)v_T]}{v_N^2 + v_B^2 + v_T^2}, \quad (11)$$

$${}^t\mathcal{K}_g = \frac{\langle \ddot{\mathbf{X}}, n \times \dot{\mathbf{X}} \rangle}{||\dot{\mathbf{X}}||^2} = \frac{\rho[-(v_N^2 + v_B^2)\xi_1 + (v_B \xi_2 + v_N \xi_3)v_T]}{v_N^2 + v_B^2 + v_T^2}, \quad (12)$$

$${}^t\mathcal{K}_n = \frac{\langle \ddot{\mathbf{X}}, n \rangle}{||\dot{\mathbf{X}}||^2} = \frac{\rho(v_N \xi_3 - v_B \xi_2)}{v_N^2 + v_B^2 + v_T^2}, \quad (13)$$

respectively, where

$$\begin{aligned}\xi_1 &= \partial_t v_T - v_N(\partial_s v_N + \kappa v_T - \tau v_B) - v_B(\partial_s v_B + \tau v_N), \\ \xi_2 &= \partial_t v_N - v_B \kappa^{-1}(\partial_s^2 v_B + 2\tau \partial_s v_N + v_N \partial_s \tau + \tau \kappa v_T - \tau^2 v_B) + v_T(\partial_s v_N + \kappa v_T - \tau v_B), \\ \xi_3 &= \partial_t v_B + v_N \kappa^{-1}(\partial_s^2 v_B + 2\tau \partial_s v_N + v_N \partial_s \tau + \tau \kappa v_T - \tau^2 v_B) + v_T(\partial_s v_B + \tau v_N), \\ n_1 &= \rho[v_B(\partial_s v_N + \kappa v_T - \tau v_B) - v_N(\partial_s v_B + \tau v_N)], \\ n_2 &= -\rho \partial_t v_B - \kappa^{-1} \rho v_N(\partial_s^2 v_B + 2\tau \partial_s v_N + v_N \partial_s \tau + \tau \kappa v_T - \tau^2 v_B) - v_B \partial_t \rho, \\ n_3 &= \rho \partial_t v_N - \kappa^{-1} \rho v_B(\partial_s^2 v_B + 2\tau \partial_s v_N + v_N \partial_s \tau + \tau \kappa v_T - \tau^2 v_B) + v_N \partial_t \rho, \\ \dot{\mathbf{X}} &= \frac{\partial \mathbf{X}}{\partial t}, \ddot{\mathbf{X}} = \frac{\partial^2 \mathbf{X}}{\partial t^2}, \dot{n} = \frac{\partial n}{\partial t}, \rho = \frac{1}{\sqrt{v_B^2 + v_N^2}}.\end{aligned}$$

Similarly, the geodesic torsion ${}^u\mathcal{T}_g$, geodesic curvature ${}^u\mathcal{K}_g$, normal curvature ${}^u\mathcal{K}_n$ of the u parameter curves $\mathbf{X}(u, t)$ on trajectory surface $\sum_{t_{max}}$ as

$${}^u\mathcal{T}_g = \frac{\langle \mathbf{X}_u, n \times n_u \rangle}{\|\mathbf{X}_u\|^2} = \frac{\rho(v_B(\rho \tau v_B - (\rho v_N)_s) + v_N((\rho v_B)_s + \rho v_N \tau))}{g}, \quad (14)$$

$${}^u\mathcal{K}_g = \frac{\langle \mathbf{X}_{uu}, n \times \mathbf{X}_u \rangle}{\|\mathbf{X}_u\|^2} = \kappa g \rho v_N, \quad {}^u\mathcal{K}_n = \frac{\langle \mathbf{X}_{uu}, n \rangle}{\|\mathbf{X}_u\|^2} = -\kappa \rho v_B, \quad (15)$$

respectively, where

$$\mathbf{X}_u = \frac{\partial \mathbf{X}}{\partial u}(u, t), \mathbf{X}_{uu} = \frac{\partial^2 \mathbf{X}}{\partial u^2}(u, t), n_u = \frac{\partial n}{\partial u}(u, t), \rho = \frac{1}{\sqrt{v_B^2 + v_N^2}}.$$

For the sake of simplicity, we denote

$$\begin{aligned}\Omega_{\mathcal{T}_g}^t &= \{\mathbf{X} = \mathbf{X}(u, t) \in \Omega^t : {}^t\mathcal{T}_g = 0\}, \quad \Lambda_{\mathcal{T}_g}^u = \{\mathbf{X} = \mathbf{X}(u, t) \in \Lambda^u : {}^u\mathcal{T}_g = 0\}, \\ \Omega_{\mathcal{K}_g}^t &= \{\mathbf{X} = \mathbf{X}(u, t) \in \Omega^t : {}^t\mathcal{K}_g = 0\}, \quad \Lambda_{\mathcal{K}_g}^u = \{\mathbf{X} = \mathbf{X}(u, t) \in \Lambda^u : {}^u\mathcal{K}_g = 0\}, \\ \Omega_{\mathcal{K}_n}^t &= \{\mathbf{X} = \mathbf{X}(u, t) \in \Omega^t : {}^t\mathcal{K}_n = 0\}, \quad \Lambda_{\mathcal{K}_n}^u = \{\mathbf{X} = \mathbf{X}(u, t) \in \Lambda^u : {}^u\mathcal{K}_n = 0\}.\end{aligned}$$

3 | SOME GEOMETRIC PROPERTIES OF SOLUTIONS OF THE GEOMETRIC CURVE FLOWS IN SPACE

In this section, we obtain the Gaussian curvature, mean curvature, and principal curvatures of some trajectory surfaces including the Hasimoto surface, shortening trajectory surface, minimal trajectory surface, $\sqrt{\tau}$ -normal trajectory surface in $\mathbb{R}^3$ and give a necessary and sufficient conditions for t -parameter and u -parameter curves of some trajectory surfaces in $\mathbb{R}^3$ to be geodesics, asymptotic lines, and principal lines.

Some geometric properties and some characterizations of parameter curves of Hasimoto surfaces in $\mathbb{R}^3$ is given¹ and is briefly presented here for completeness.

Theorem 1. (see¹) Let $\mathbf{X} = \mathbf{X}(u, t)$ be a Hasimoto surface $\sum_1$ in $\mathbb{R}^3$.

(1) Then the Gaussian curvature K , mean curvature H , principal curvatures P_i ($i=1,2$) of the Hasimoto surface $\sum_1$ are

$$\begin{aligned}K &= -\frac{\kappa_{ss}}{\kappa}, \quad H = \frac{1}{2\kappa}(\frac{\kappa_{ss}}{\kappa} - \kappa^2 - \tau^2), \\ P_1 &= \frac{1}{2\kappa}(\frac{\kappa_{ss}}{\kappa} - \kappa^2 - \tau^2) + \frac{\sqrt{(\kappa^{-1}\kappa_{ss} - \kappa^2 - \tau^2)^2 + 4\kappa\kappa_{ss}}}{2\kappa}, \\ P_2 &= \frac{1}{2\kappa}(\frac{\kappa_{ss}}{\kappa} - \kappa^2 - \tau^2) - \frac{\sqrt{(\kappa^{-1}\kappa_{ss} - \kappa^2 - \tau^2)^2 + 4\kappa\kappa_{ss}}}{2\kappa}.\end{aligned}$$

respectively;

- (2) We give necessary and sufficient conditions for parameter curves $\mathbf{X}(u, t)$ of a Hasimoto surface Σ_1 in $\mathbb{R}^3$ to be principal lines, geodesics, and asymptotic lines, i.e.,

$$\begin{aligned}\Omega_{\mathcal{T}_g}^t &= \{\mathbf{X} \in \Omega^t : \tau = 0\}, & \Lambda_{\mathcal{T}_g}^u &= \{\mathbf{X} \in \Lambda^u : \tau = 0\}, \\ \Omega_{\mathcal{K}_g}^t &= \{\mathbf{X} \in \Omega^t : \kappa_s = 0\}, & \Lambda_{\mathcal{K}_g}^u &= \Lambda^u, \\ \Omega_{\mathcal{K}_n}^t &= \{\mathbf{X} \in \Omega^t : \kappa_{ss} - \kappa\tau^2 = 0\}, & \Lambda_{\mathcal{K}_n}^u &= \{\mathbf{X} \in \Lambda^u : \kappa = 0\}.\end{aligned}$$

- (3) The binormal flow (5) is inextensible flows (i.e., geometric flows with $\frac{\partial g}{\partial t} = 0$).

Proof. By using Eqs. (9), Eqs. (10) and Eqs. (11)-(15), we have

$$\begin{aligned}I &= \begin{pmatrix} g^2 & 0 \\ 0 & \kappa^2 \end{pmatrix}, II = \begin{pmatrix} -g^2\kappa & g\tau\kappa \\ g\tau\kappa & \kappa_{ss} - \kappa\tau^2 \end{pmatrix}, \\ {}^t\mathcal{T}_g &= -\tau, {}^t\mathcal{K}_g = \kappa_s, {}^t\mathcal{K}_n = \frac{\kappa_{ss} - \kappa\tau^2}{\kappa^2}, \\ {}^u\mathcal{T}_g &= \frac{\tau}{g}, {}^u\mathcal{K}_g = 0, {}^u\mathcal{K}_n = -\kappa, \partial_t g = (-\kappa v_N + \partial_s v_T)g = 0.\end{aligned}$$

This completes the proof. □

Theorem 2. Let $\mathbf{X} = \mathbf{X}(u, t)$ be a shortening trajectory surface Σ_2 in $\mathbb{R}^3$.

- (1) Then the Gaussian curvature K , mean curvature H , principal curvatures P_i ($i=1,2$) of the shortening trajectory surface Σ_2 are

$$\begin{aligned}K &= -\tau^2, H = \frac{2\tau\kappa_s + \tau_s\kappa}{2\kappa^2}, \\ P_1 &= \frac{2\tau\kappa_s + \tau_s\kappa}{2\kappa^2} + \frac{1}{2\kappa^2} \sqrt{(2\tau\kappa_s + \tau_s\kappa)^2 + 4\kappa^4\tau^2}, \\ P_2 &= \frac{2\tau\kappa_s + \tau_s\kappa}{2\kappa^2} - \frac{1}{2\kappa^2} \sqrt{(2\tau\kappa_s + \tau_s\kappa)^2 + 4\kappa^4\tau^2},\end{aligned}$$

respectively;

- (2) we give necessary and sufficient conditions for parameter curves $\mathbf{X}(u, t)$ of a shortening trajectory surface Σ_2 in $\mathbb{R}^3$ to be principal lines, geodesics, and asymptotic lines, i.e.,

$$\begin{aligned}\Omega_{\mathcal{T}_g}^t &= \{\mathbf{X} \in \Omega^t : \tau = 0\}, & \Lambda_{\mathcal{T}_g}^u &= \{\mathbf{X} \in \Lambda^u : \tau = 0\}, \\ \Omega_{\mathcal{K}_g}^t &= \{\mathbf{X} \in \Omega^t : \kappa_s = 0\}, & \Lambda_{\mathcal{K}_g}^u &= \{\mathbf{X} \in \Lambda^u : \kappa = 0\}, \\ \Omega_{\mathcal{K}_n}^t &= \{\mathbf{X} \in \Omega^t : 2\tau\kappa_s + \kappa\tau_s = 0\}, & \Lambda_{\mathcal{K}_n}^u &= \Lambda^u.\end{aligned}$$

Proof. By using Eqs. (9), Eqs. (10) and Eqs. (11)-(15), we have

$$\begin{aligned}I &= \begin{pmatrix} g^2 & 0 \\ 0 & \kappa^2 \end{pmatrix}, II = \begin{pmatrix} 0 & g\tau\kappa \\ g\tau\kappa & \kappa\tau_s + 2\tau\kappa_s \end{pmatrix}, \\ {}^t\mathcal{T}_g &= -\tau, {}^t\mathcal{K}_g = \kappa_s, {}^t\mathcal{K}_n = \frac{2\tau\kappa_s + \kappa\tau_s}{\kappa^2}, \\ {}^u\mathcal{T}_g &= \frac{\tau}{g}, {}^u\mathcal{K}_g = \kappa g, {}^u\mathcal{K}_n = 0, \partial_t g = (-\kappa v_N + \partial_s v_T)g = -\kappa^2 g.\end{aligned}$$

This finishes the proof of Theorem 2. □

Theorem 3. Let $\mathbf{X} = \mathbf{X}(u, t)$ be a minimal trajectory surface Σ_3 in $\mathbb{R}^3$.

- (1) Then the Gaussian curvature K , mean curvature H , principal curvatures P_i ($i=1,2$) of the minimal trajectory surface Σ_3 are

$$K = -\tau^2, H = 0, P_1 = \tau, P_2 = -\tau,$$

respectively;

(2) we give necessary and sufficient conditions for parameter curves $\mathbf{X}(u, t)$ of a minimal trajectory surface Σ_3 in $\mathbb{R}^3$ to be principal lines, geodesics, and asymptotic lines, i.e.,

$$\begin{aligned}\Omega_{\mathcal{T}_g}^t &= \emptyset, & \Lambda_{\mathcal{T}_g}^u &= \emptyset, \\ \Omega_{\mathcal{K}_g}^t &= \{\mathbf{X} \in \Omega^t : \tau_s = 0\}, & \Lambda_{\mathcal{K}_g}^u &= \emptyset, \\ \Omega_{\mathcal{K}_n}^t &= \Omega^t, & \Lambda_{\mathcal{K}_n}^u &= \Lambda^u.\end{aligned}$$

Proof. By using Eqs. (9), Eqs. (10) and Eqs. (11)-(15), we have

$$\begin{aligned}I &= \begin{pmatrix} g^2 & 0 \\ 0 & \frac{1}{\tau} \end{pmatrix}, II = \begin{pmatrix} 0 & g\sqrt{\tau} \\ g\sqrt{\tau} & 0 \end{pmatrix}, \\ {}^t\mathcal{T}_g &= -\tau, {}^t\mathcal{K}_g = -\frac{\tau_s}{2\tau^{\frac{3}{2}}}, {}^t\mathcal{K}_n = 0, \\ {}^u\mathcal{T}_g &= \frac{\tau}{g}, {}^u\mathcal{K}_g = \kappa g, {}^u\mathcal{K}_n = 0, \partial_t g = \frac{\kappa}{\sqrt{\tau}}.\end{aligned}$$

Note that $\kappa \neq 0$, $\tau > 0$, hence this completes the proof. $\square$

Theorem 4. Let $\mathbf{X} = \mathbf{X}(u, t)$ be a $\sqrt{\tau}$ -normal trajectory surface Σ_4 in $\mathbb{R}^3$.

(1) Then the Gaussian curvature K , mean curvature H , principal curvatures P_i ($i=1,2$) of $\sqrt{\tau}$ -normal trajectory surface Σ_4 are

$$K = -\tau^2, \quad H = \frac{\tau_s}{\kappa}, \quad P_1 = \frac{\tau_s + \sqrt{\tau_s^2 + \kappa^2 \tau^2}}{\kappa}, \quad P_2 = \frac{\tau_s - \sqrt{\tau_s^2 + \kappa^2 \tau^2}}{\kappa},$$

respectively;

(2) we give necessary and sufficient conditions for parameter curves $\mathbf{X}(u, t)$ of a $\sqrt{\tau}$ -normal trajectory surface Σ_4 in $\mathbb{R}^3$ to be principal lines, geodesics, and asymptotic lines, i.e.,

$$\begin{aligned}\Omega_{\mathcal{T}_g}^t &= \{\mathbf{X} \in \Omega^t : \tau = 0\}, & \Lambda_{\mathcal{T}_g}^u &= \{\mathbf{X} \in \Lambda^u : \tau = 0\}, \\ \Omega_{\mathcal{K}_g}^t &= \{\mathbf{X} \in \Omega^t : \tau_s = 0\}, & \Lambda_{\mathcal{K}_g}^u &= \emptyset, \\ \Omega_{\mathcal{K}_n}^t &= \{\mathbf{X} \in \Omega^t : \tau_s = 0\}, & \Lambda_{\mathcal{K}_n}^u &= \Lambda^u.\end{aligned}$$

Proof. By using Eqs. (9), Eqs. (10) and Eqs. (11)-(15), we have

$$\begin{aligned}I &= \begin{pmatrix} g^2 & 0 \\ 0 & \tau \end{pmatrix}, II = \begin{pmatrix} 0 & g\tau^{\frac{3}{2}} \\ g\tau^{\frac{3}{2}} & \frac{2\tau\tau_s}{\kappa} \end{pmatrix}, \\ {}^t\mathcal{T}_g &= -\tau, {}^t\mathcal{K}_g = \frac{\tau_s}{2\sqrt{\tau}}, {}^t\mathcal{K}_n = \frac{2\tau_s}{\kappa}, \\ {}^u\mathcal{T}_g &= \frac{\tau}{g}, {}^u\mathcal{K}_g = \kappa g, {}^u\mathcal{K}_n = 0, \partial_t g = -\kappa\sqrt{\tau}g.\end{aligned}$$

Note that $\kappa \neq 0$, $\tau \geq 0$, this finishes the proof of Theorem 4. $\square$

Remark 1. There the other types geometric curve flows (such as the geometric KdV curve flow or also called Fukumoto-Miyazaki model, the generalized bi-Schrödinger flow or also called Fukumoto-Moffatt model) in^{29,30}, as following

(i) A Fukumoto-Miyazaki surface is the surface traced out by a family of curves $\{\Gamma_t\}_{t \in [0, t_{max})}$ in $\mathbb{R}^3$ as it evolves over time according to the geometric KdV curve flow:

$$\begin{aligned}\partial_t \mathbf{X} &= \frac{1}{2}\kappa \mathbf{T} + \kappa_s \mathbf{N} + \kappa \tau \mathbf{B}, & \text{in } \mathbb{S}^1 \times [0, t_{max}), \\ \mathbf{X}|_{t=0} &= \mathbf{X}_0 & \text{in } \mathbb{S}^1,\end{aligned} \tag{16}$$

where $\mathbf{X}_0$ is the parametrization for the initial curve Γ_0 in $\mathbb{R}^3$ (see e.g.^{29,3}). Note that the geometric KdV curve flow (16) is inextensible flows, in fact,

$$\partial_t g = (-\kappa v_N + \partial_s v_T)g = (-\kappa \kappa_s + (\frac{1}{2}\kappa^2))g = 0.$$

- (ii) A Fukumoto-Moffatt surface is the surface traced out by a family of curves $\{\Gamma_t\}_{t \in [0, t_{max})}$ in $\mathbb{R}^3$ as it evolves over time according to the generalized bi-Schrödinger flow:

$$\begin{aligned} \partial_t \mathbf{X} &= \lambda \left\{ \kappa \mathbf{B} + \nu [\kappa^2 \tau \mathbf{T} + (2\kappa_s \tau + \kappa \tau_s) \mathbf{N} + (\kappa \tau^2 - \kappa_{ss}) \mathbf{B}] + \mu \kappa^3 \mathbf{B} \right\}, \\ \mathbf{X}|_{t=0} &= \mathbf{X}_0, \end{aligned} \quad (17)$$

where $\mathbf{X}_0$ is the parametrization for the initial curve Γ_0 in $\mathbb{R}^3$, λ, μ and ν are three real parameters (see e.g.^{30,4}). Notice again that the generalized bi-Schrödinger flow (17) is inextensible flows, this is because

$$\partial_t g = (-\kappa \nu_N + \partial_s \nu_T)g = (-\kappa \lambda \nu (2\kappa_s \tau + \kappa \tau_s) + \lambda \nu (\kappa^2 \tau)_s)g = 0.$$

Similarly, we also give some geometric properties and some characterizations of u -parameter curves and t -parameter curves of Fukumoto-Miyazaki surface and Fukumoto-Moffatt surface.

4 | CONCLUSION

In this paper we investigated the geometric properties of solutions of the geometric curve flows in Euclidean 3-space $\mathbb{R}^3$ with respect to the Frenet frame. This trajectory surfaces including the Hasimoto surface, shortening trajectory surface, minimal trajectory surface, $\sqrt{\tau}$ -normal trajectory surface may be useful for some specific applications in theoretical physics and fluid dynamics. It is of interest to analyze some properties (such as the long term behavior, upper bound for the area of the $\sqrt{\tau}$ -normal trajectory surface and terminating time) of space curves evolved by the $\sqrt{\tau}$ -normal geometric flow (8). These topics are beyond the scope of this paper, but may be considered in future work.

ACKNOWLEDGMENTS

The authors are very grateful to the anonymous referees for their valuable suggestions. The authors are supported by the Natural Science Foundation of Jiangxi Province (No. 20212BAB211005) and College Students' innovation and entrepreneurship training program (N0. S202210418008).

Financial disclosure

None reported.

Conflict of interest

The authors declare that there are no conflicts of interest regarding the publication of this paper.

DATA AVAILABILITY

No data was used for the research described in the article.

ORCID

Shi-ping Zhong <https://orcid.org/0000-0001-9183-7268>

References

1. N. H. ABDEL-ALL, R. A. HUSSEIN, T. YOUSSEF, *Hasimoto surfaces*, Life Sci. J., 9 (2012) 556-560.

2. M.E. AYDIN, A. MIHAI, A.O. OGRENMIS, M. ERGUT, *Geometry of the solutions of localized induction equation in the pseudo-Galilean space*, Adv. Math. Phys., 2015 (2015) 905978.
3. Q. DING, Y.D. WANG *Geometric KdV flows, motions of curves and the third order system of the AKNS hierarchy*, Int. J. Math., 22 (2011), 1013-1029.
4. Q. DING, W. WANG, Y.D. WANG, *The Fukumoto-Moffatt's model in the vortex filament and generalized bi-Schrödinger maps*, Phys. Lett. A, 375 (2011) 1457-1460.
5. M. ERDOGDU, M. OZDEMIR, *Geometry of Hasimoto surfaces in Minkowski 3-space*, Math. Phys. Anal. Geom., 17 (2014) 169-181.
6. M. ELZAWY, *Hasimoto surfaces in Galilean space*, J. Egypt. Math. Soc., 29 (2021) 5.
7. M. ERDOGDU, A. YAVUZ,, *Differential geometric aspects of nonlinear Schrödinger equation*, Communications Faculty of Sciences University of Ankara Series A1 Mathematics and Statistics, 70 (1) (2021) 510-521.
8. K. EREN , *New Representation of Hasimoto Surfaces with the Modified Orthogonal Frame*, Konuralp J. Math., 10(1) (2022) 69-72.
9. N. GÜRBÜZ, *Hasimoto surfaces according to three classes of curve evolution with Darboux frame in Euclidean space*, Gece Publishing, 2018.
10. H. HASIMOTO, *A soliton on a vortex filament*, J. Fluid Mech., 51 (1972) 477-485.
11. E. HAMOUDA, O. MOAAZ, C. CESARANO, S. ASKAR, A. ELSHARKAWY, *Geometry of Solutions of the Quasi-Vortex Filament Equation in Euclidean 3-Space $\mathbb{E}^3$* , Mathematics, 10(6) (2022) 891 1-11.
12. T.A. IVEY, *Helices, Hasimoto surfaces and Bäcklund transformations*, Can. Math. Bull., 43 (2000) 427-439
13. A. KELLECI, M. BEKTAS, M. ERGÜT, *The Hasimoto surface according to bishop frame*, Adiyaman Üniversitesi Fen Bilimleri Dergisi, 9(1) (2019) 13-22.
14. S.G. MOHAMED, *Binormal motions of inextensible curves in de-sitter space $\mathbb{S}^{2,1}$* , J. Egypt. Math. Soc., 25 (2017) 313318.
15. N. GÜRBÜZ, D.W. YOON, *Hasimoto surfaces for two classes of curve evolution in Minkowski 3-space*, Demonstratio Mathematica, 53(3) (2020) 277-284.
16. N.E. GÜRBÜZ, R. MYRZAKULOV, Z. MYRZAKULOVA, *Three anholonomy densities for three formulations with anholonomic coordinates with hybrid frame in $\mathbb{R}^{3,1}$* , Optik, 261 (2022). 169161
17. F. MARTÍN, A. SAVAS-HALILAJ, K. SMOCHYK, *On the topology of translating solitons of the mean curvature flow*, Calc. Var. Partial Differential Equations, 54 (2015), 2853-2882.
18. J. MINARČÍK, M. BENEŠ, *Nondegenerate homotopy and geometric flows*, Homology, Homotopy and Applications, 24(2) (2022) 255-264.
19. R.L. RICCA, *Rediscovery of Da Rios equations*, Nature, 352 (1991) 561562.
20. C. YU, H. SCHUMACHER, K. CRANE, *Repulsive Curves*, ACM Trans. Graph., 40 (2021) 121.
21. R. A. HUSSEIN, S. G. MOHAMED, *Generated surfaces via inextensible flows of curves in $\mathbb{R}^3$* , Journal of Applied Mathematics, 2016 (2016) 6178961.
22. M. BERGOU, M. WARDETZKY, S. ROBINSON, B. AUDOLY, E. GRINSUN, *Discrete elastic rods*, ACM Trans. Graphics, 27 (2008) 63:01-63:12
23. J. MINARČÍK, M. BENEŠ, *Long-term Behavior of Curve Shortening Flow in $\mathbb{R}^3$* , SIAM J. Math. Anal., 52(2) (2020) 1221–1231.

24. S. J. ALTSCHULER, *Singularities of the curve shrinking flow for space curves*, J. Differ. Geom., 34 (1991) 491-514.
25. S. J. ALTSCHULER, M. GRAYSON, *Shortening space curves and flow through singularities*, J. Differ. Geom., 35 (1992) 283-298.
26. J. MINARČÍK, M. BENEŠ, *Minimal surface generating flow for space curves of non vanishing torsion*, Discrete Contin. Dyn. Syst. Ser. B, 27(11) (2022) 6605-6617.
27. M. BENEŠ, MIROSLAV KOLAŘ, DANIEL ŠEVČOVIČ, *Qualitative and Numerical Aspects of a Motion of a Family of Interacting Curves in Space*, SIAM Journal on Applied Mathematics, 82(2) (2022) 549-575.
28. A.T. Ali, H.S.A. Aziz, A.H. Sorour, *Ruled surfaces generated by some special curves in Euclidean 3-Space.*, J. Egypt. Math. Soc., 21 (2013) 285-294.
29. Y. FUKUMOTO, T. MIYAZAKI, *Three-dimensional distortions of a vortex filament with axial velocity*, J. Fluid. Mech., 22 (1991) 369-416.
30. Y. FUKUMOTO, H.K. MOFFATT, *Motion and expansion of a visous vortex ring, I. A higer-order asymptotic formula for the velocity*, J. Fluid Mech., 417 (2000) 1-45.

How to cite this article: S.P. Zhong, Z.H. Zhao, and X.J. Wan, (2022), Geometry of solutions of the geometric curve flows in space, , .