

# A Soldier's Return to Duty after Minimally Invasive Correction of Complex Congenital Cardiovascular Disease

Tarin Phillips<sup>1</sup>, Adam Kisling<sup>1</sup>, and Robert Gallagher<sup>1</sup>

<sup>1</sup>Walter Reed National Military Medical Center

December 5, 2022

## Abstract

We report a case of incidental detection of severe aortic coarctation, severe secundum atrial septal defect, and bicuspid aortic valve in an active-duty military service member. A single complex minimally invasive procedure successfully corrected his coarctation and atrial septal defect allowing this patient to continue military service.

Title: A Soldier's Return to Duty after Minimally Invasive Correction of Complex Congenital Cardiovascular Disease

Authors: Tarin Phillips DO<sup>1</sup>, Adam Kisling MD<sup>1</sup>, & Robert Gallagher MD FACC FSCAI<sup>1</sup>

Affiliation:

<sup>1</sup> Walter Reed National Military Medical Center, Department of Cardiology, Bethesda, MD United States of America

Correspondence:

Tarin Phillips, DO [tarin.c.phillips.mil@health.mil](mailto:tarin.c.phillips.mil@health.mil)

Adam Kisling, MD [adam.j.kisling.mil@health.mil](mailto:adam.j.kisling.mil@health.mil)

Robert Gallagher, MD [robert.m.gallagher7.mil@health.mil](mailto:robert.m.gallagher7.mil@health.mil)

Acknowledgements: None

Funding: None

Conflict of Interest Statement: The authors have no competing interests to declare that are relevant to the content of this article.

Data Availability Statement: All data supporting this study's findings are available from the corresponding author upon reasonable request.

Informed Consent: Written informed consent in accordance with Walter Reed National Military Medical Center was obtained from the patient.

ORCID:

Tarin Phillips: 0000-0003-3390-4375

Adam Kisling: 0000-0002-4254-0641

Robert Gallagher: 0000-0001-8143-553X

## Abstract:

We report a case of incidental detection of severe aortic coarctation, severe secundum atrial septal defect, and bicuspid aortic valve in an active-duty military service member. A single complex minimally invasive procedure successfully corrected his coarctation and atrial septal defect allowing this patient to continue military service.

Keywords: atrial septal defect, aortic coarctation, bicuspid aortic valve

## Introduction:

The incidence of congenital heart disease (CHD) is roughly 40,000 cases in the United States every year [1]. Aortic coarctation (CoA) occurs in 4 of every 10,000 births and making up roughly 5-8% of all CHD [2,3]. Of those with a bicuspid aortic valve (BAV) only 7% have a concomitant CoA, but 75% of patients with a CoA also have BAV [3]. Atrial septal defects (ASD) occur in roughly 1 in every 1,860 births in the United States [2]. Genetic abnormalities are common in patients with CHD, and genetic testing for concomitant syndromes may be reasonable in patients with adult CHD. While there appears to be a genetic link between CoA and BAV, there is unifying etiology for the development of these conditions with a concomitant ASD [4].

CoA presentation is largely dependent on the severity ranging from asymptomatic in most cases to headaches, epistaxis, exertional intolerance, dizziness, and lower extremity claudication [5]. These symptoms are largely driven by increased afterload proximal to the narrowing and decrease blood flow distally leading to discrepant upper extremity hypertension, elevated upper to lower extremity pressure gradient, weak or absent femoral pulses [5]. Additionally, a prominent, nondisplaced apical impulse, loud A2, or continuous or systolic murmur radiating to the scapula due to significant collaterals may be appreciated on exam [5].

Most ASDs identified during adulthood are asymptomatic and found incidentally. A minority of these patient will develop a gradual decline in exercise capacity and occasionally overt dyspnea on exertion particularly if pulmonary arterial hypertension is present [6]. One of the ways ASDs are characterized is the degree of left-to-right shunting. Shunts that are hemodynamically significant enough to lead to physiologic sequelae should usually be closed. The significance of shunts is measured with a Qp:Qs which is the ratio of pulmonary blood flow to systemic blood flow. If the Qp:Qs is greater than 1.5, then the hemodynamic impact of the shunt will be significant. Intracardiac shunts and the subsequent increase in intracardiac circulatory volume can lead to chamber dilation particularly of the atria. In some patients this can contribute to atrial arrhythmias [6]. On exam patients with significant ASD will classically have a fixed split S2 due to increased pulmonary circulatory volume and the subsequent delayed closure of the pulmonic valve [6].

The presence of CoA, BAV with any complications, or ASD is disqualifying for any potential United States military recruit [7]. For retaining current service members, the United States Army allows exceptions for CHD that can be repaired with resolution of long-term risk and complications [7].

## Case Presentation:

A 20-year-old fit active-duty male with a history of growth delay as a child and bilateral upper extremity hypertension was rescued from the driver's seat of an M1A2 Abrams tank after suffering carbon monoxide poisoning from an exhaust system malfunction during an overseas deployment exercise. He was intubated in the field and transferred to higher level of care where there was concern for myocardial infarction. After stabilization and extubation, he was medically evacuated to a military treatment facility in the United States where his troponin level was normal, and his ECG was normal. However, a computed tomography scan (CT) of his chest revealed narrowing of the proximal descending thoracic aorta, and a transthoracic echocardiogram (TTE) was notable for a bicuspid aortic valve. Thoracic magnetic resonance imaging, seen in Figure 1, corroborated these findings and demonstrated right atrial enlargement concerning for atrial septal defect (ASD). Left and right heart catheterization showed severe CoA with a peak-to-peak gradient of 30 mmHg and extensive collateral artery formation. The catheterization also demonstrated a mildly elevated

mean pulmonary artery pressure and a pulmonary to systemic blood flow ratio of 1.5, reflecting a severe left to right shunt. CT coronary angiography measured the ASD at 7 mm.

After multidisciplinary heart team discussion, the patient underwent successful percutaneous transcatheter aortic coarctation stenting and ASD closure during a single procedure [8]. Immediate post-procedure coarctation stent gradient was 5 mmHg. Follow up TTE demonstrated no residual inter-atrial shunt, and CT aortic angiography demonstrated patent endovascular stent without complication, seen in Figure 2. The patient's upper extremity hypertension improved, and he was able to return to non-straining exercise without symptoms. Four months following the procedure, the patient completed a Bruce protocol treadmill stress test running for 12:36 minutes and reaching 14.5 METS with normal ECG and hemodynamic response. Given his CHD, it was recommended that he refrain from high intensity competitive athletics with forceful body contact and exercises involving straining. After appropriate recovery, the patient was able to return to duty and met cardiovascular military retention criteria.

### Discussion:

We present a case of complex CHD that was corrected with a single minimally invasive percutaneous procedure that allowed this service member to be retained for duty in the United States Army. In addition to multidisciplinary team discussion, it is important to consider the patient's wishes when making management decisions for CHD. Prior to making any treatment decisions, the patient was counseled thoroughly on the implications that his condition and its management would have on his health and career. In cases where multiple treatment options are reasonable and timing of intervention is flexible, a patient should be well informed, and their autonomy must be preserved regardless of the social implications of their decision. Our patient met current guidelines for the correction of his coarctation given his systemic hypertension, elevated peak to peak gradient, and severe collateralization [8]. The focal narrowing of the CoA was just distal to the subclavian artery making stenting technically challenging. Often, covered stents are used for CoA but this case was not amendable due to concern for occlusion of the subclavian artery. In our patient, a 36mm intrastent mega LD was use beginning proximal and extending distal to the subclavian artery correcting the focal narrowing. This temporarily reduced blood flow to the subclavian artery until a stent fenestration could be ballooned creating an intra-stent ostium for the subclavian artery. This is depicted in the post-procedural follow up CT aortic angiogram in Figure 2 with patent endovascular stent. Post-operatively this patient's upper extremity to lower extremity blood pressure discrepancy normalized, but he developed systemic hypertension, which is a common occurrence following CoA stenting that responded well to losartan. Regarding his secundum ASD, there was evidence of left to right shunt on multiple image modalities. This patient had normal pulmonary vascular resistance and a pulmonary artery systolic pressure less than half of the system pressure. This is important because ASD closure is contraindicated in adults with PA systolic pressure greater than two thirds systemic, pulmonary vascular resistance greater than two thirds systemic, or a net right-to-left shunt [8]. Qp:Qs measurement showed that there was a significantly higher pulmonary blood flow than systemic blood flow. This Qp:Qs [?]1.5 indicated that the shunt was sufficiently large to cause physiological sequelae. Additionally, there was already chamber dilation in the form of right atrial enlargement due to increased volume load. When hemodynamically significant ASDs are identified, it is important to assess the patient's functional capacity and whether the ASD is symptomatic as there is more clear benefit for patients who are functionally impaired by their ASD [8]. In this patient, a 10mm Amplatzer Septal Occluder was placed with usual technique without complication. Serial follow up echocardiograms demonstrated a well seated inter-atrial device without residual shunt. Bicuspid aortic valves are the most common congenital heart disease, and fusion of the left and right coronary cusps is the most common phenotype [3,9]. At the time of evaluation, this patient had not developed valvular dysfunction. The development of aortic stenosis often occurs earlier than tri-leaf patients and is suggested to be related to leaflet orientation [9]. Aortic regurgitation is also a common finding in BAV patients and is independent of valve stenosis [9]. At baseline, patient's with BAV generally have larger aortic size which predisposes development of early root and ascending aortic dilation [9]. Given these findings in conjunction with the patient's other congenital disease it is paramount that annual lifelong cardiology evaluation is completed. The presence of a bicuspid aortic valve is an important consideration when determining the timing and method for treatment of these

associated conditions. Additionally, it is recommended that all first-degree relatives to be screened for BAV and aortopathy with echocardiography [8].

This case highlights the patient-centered shared decision making afforded to congenital heart disease patients by the contemporary multidisciplinary heart team. The single transcatheter procedure performed for correction of this patient's aortic coarctation and atrial septal defect provided complete correction of all indicated congenital lesions while remarkably permitting his continued service in the United States Armed Forces. Author Contributions:

TP drafted the initial manuscript, collected data, performed literature search, and reviewed and revised the manuscript. AK conceptualized and designed the study, collected data, performed literature search, and reviewed and revised the manuscript. RG participated in case management, performed left and right heart catheterization, atrial septal defect repair, aortic coarctation angioplasty, and reviewed and revised the manuscript. All authors approved the final manuscript as submitted and agree to be accountable for all aspects of the work.

#### References:

1. Centers for Disease Control and Prevention. (2022, January 24). Data and statistics on congenital heart defects. Centers for Disease Control and Prevention. Retrieved August 28, 2022.
2. Mai CT, Isenburg JL, Canfield MA, et al. National population-based estimates for major birth defects, 2010-2014. *Birth Defects Res.* 2019;111(18):1420-1435. doi:10.1002/bdr2.1589
3. Sinning, Christoph et al. "Bicuspid aortic valve and aortic coarctation in congenital heart disease-important aspects for treatment with focus on aortic vasculopathy." *Cardiovascular diagnosis and therapy* vol. 8,6 (2018): 780-788. doi:10.21037/cdt.2018.09.20
4. Parker, Lauren E, and Andrew P Landstrom. "Genetic Etiology of Left-Sided Obstructive Heart Lesions: A Story in Development." *Journal of the American Heart Association* vol. 10,2 (2021): e019006. doi:10.1161/JAHA.120.019006
5. Kim, Yuli Y et al. "Aortic Coarctation." *Cardiology clinics* vol. 38,3 (2020): 337-351. doi:10.1016/j.ccl.2020.04.003
6. Bradley, Elisa A, and Ali N Zaidi. "Atrial Septal Defect." *Cardiology clinics* vol. 38,3 (2020): 317-324. doi:10.1016/j.ccl.2020.04.001
7. AR 40-501- U.S. Army Standards of Medical Fitness. Accessed 08 May 2022.
8. Stout, Karen K et al. "2018 AHA/ACC Guideline for the Management of Adults With Congenital Heart Disease: A Report of the American College of Cardiology/American Heart Association Task Force on Clinical Practice Guidelines." *Journal of the American College of Cardiology* vol. 73,12 (2019): e81-e192. doi:10.1016/j.jacc.2018.08.1029
9. Mordi, Ify, and Nikolaos Tzemos. "Bicuspid aortic valve disease: a comprehensive review." *Cardiology research and practice* vol. 2012 (2012): 196037. doi:10.1155/2012/196037

Figure 1. Aortic MRI demonstrating aortic coarctation with focal narrowing to 10mm with severely increased collateral blood supply to the lower descending aorta, bicuspid aortic valve, and dilated right atrium concerning for atrial septal defect.



Figure 2. Cardiac CT Angiogram 1 month following procedure demonstrating patent coarctation stent with intact flow to the subclavian artery and no evidence of aneurysm or other complication.

