

# Sink formation during shoot infection of tomato Micro-Tom by *Moniliophthora perniciosa*, the causal agent of cacao witches' broom disease, limits sugar supply for effective development of lateral roots

Antonio Figueira<sup>1</sup>, Daniele Paschoal<sup>1</sup>, Vitor Ometto<sup>2</sup>, João Vitor O. Mendes<sup>1</sup>, Laura Cazetta<sup>1</sup>, Esther Carrera<sup>3</sup>, Monica Lanzoni Rossi<sup>1</sup>, Juliana Aricetti<sup>4</sup>, Piotr Mieczkowski<sup>5</sup>, Gabriel Carvalho<sup>6</sup>, Igor Cesarino<sup>6</sup>, Simone Silva<sup>7</sup>, Rafael Ribeiro<sup>7</sup>, Paulo Teixeira<sup>2</sup>, and Eder da Silva<sup>1</sup>

<sup>1</sup>Universidade de Sao Paulo Centro de Energia Nuclear na Agricultura

<sup>2</sup>Universidade de Sao Paulo Escola Superior de Agricultura Luiz de Queiroz

<sup>3</sup>Universitat Politecnica de Valencia

<sup>4</sup>Centro Nacional de Pesquisa em Energia e Materiais

<sup>5</sup>The University of North Carolina at Chapel Hill

<sup>6</sup>Universidade de Sao Paulo Instituto de Biociencias

<sup>7</sup>Universidade Estadual de Campinas Instituto de Biologia

November 14, 2023

## Abstract

*Moniliophthora perniciosa* causes the witches' broom disease of cacao, and it can infect the tomato Micro-Tom (MT). Typical symptoms are stem swelling and shoot outgrowth, whereas reduction in root biomass is another side effect. We investigated whether the impairment of root growth derives from a hormonal imbalance or sink competition. Intense stem swelling coincided with a reduction in root biomass, predominantly of lateral roots. A few genes involved in hormone metabolism were activated; however, hormonal levels were not altered. Inoculation of the auxin highly-responsive *entire* genotype maintained the impaired root phenotype. Genes involved in root respiration, carbohydrate, amino acid and cell wall metabolism were repressed, whereas genes linked to water/nitrogen/phosphorous starvation were upregulated. Lower levels of sugars and amino acids suggested carbohydrate deprivation. Less <sup>13</sup>C accumulated in roots of infected MT, but not in the symptomless low-cytokinin MT-transgenic line that overexpresses *CYTOKININ OXIDASE-2* (*35S::AtCKX2*). We show evidence that the impairment of root development potentially derives from a reduction of photoassimilate supply by the establishment of a strong sink at the shoot symptomatic infection site, rather than hormonal imbalance. We speculate that this impact may contribute to the dramatic decrease in cocoa yields after *M. perniciosa* invasion.

## Hosted file

Paschoal et al. Strong sink.docx available at <https://authorea.com/users/29348/articles/686929-sink-formation-during-shoot-infection-of-tomato-micro-tom-by-moniliophthora-perniciosa-the-causal-agent-of-cacao-witches-broom-disease-limits-sugar-supply-for-effective-development-of-lateral-roots>







