

Yacoub valve-sparing aortic root replacement with a concomitant repair of Tetralogy of Fallot: a case report

Yunfei Ling¹, Zhongze Cao², Qi An³, and Yongjun Qian⁴

¹west china hospital

²West China School of Medicine, Sichuan University

³Affiliation not available

⁴West China Hospital of Medicine

March 07, 2024

Abstract

Here we report a case of a previously unoperated 35-year-old male TOF patient. He was admitted due to dyspnea, initial physical examination demonstrated cyanosis and heart murmurs. Echocardiography suggested secundum atrial septal defect, perimembranous ventricular septal defect and mild-severe aortic regurgitation. Computed tomography angiography indicated aortic root dilation and suspicious right coronary artery aneurysm. Concurrent repair of TOF and VSARR was performed successfully and the patient had a satisfactory recovery.

Title: Yacoub valve-sparing aortic root replacement with a concomitant repair of Tetralogy of Fallot: a case report

Running title: Yacoub technique in TOF with VSARR: a case report

Yunfei Ling¹, MD, PhD, Zhongze Cao², MD, Qi An¹, MD, Yongjun Qian, MD, PhD

¹Department of Cardiovascular Surgery, West China Hospital, Sichuan University, 37 Guoxue Lane, Chengdu, Sichuan 610041, China

²West China School of Medicine, Sichuan University, 37 Guoxue Lane, Chengdu, Sichuan 610041, China

Corresponding Author:

Dr. Qi An, MD

37 Guoxue Lane, Chengdu, Sichuan 610041, China

E-mail: anqi@scu.edu.cn

Abstract

Here we report a case of a previously unoperated 35-year-old male TOF patient. He was admitted due to dyspnea, initial physical examination demonstrated cyanosis and heart murmurs. Echocardiography suggested secundum atrial septal defect, perimembranous ventricular septal defect and mild-severe aortic regurgitation. Computed tomography angiography indicated aortic root dilation and suspicious right coronary artery aneurysm. Concurrent repair of TOF and VSARR was performed successfully and the patient had a satisfactory recovery.

Enlargement of the aorta is a severe sequela of tetralogy of Fallot (TOF), which can lead to fatal outcomes like aortic regurgitation (AR), dissection, aneurysm, or even rupture with fatal consequences, even in those

patients who have had a repair of TOF.[1] Valve sparing aortic root replacement (VSARR), also known as the Yacoub technique, is a surgical procedure that has the advantages of correcting AR, preserving native aortic valves, avoiding prosthetic valves and subsequent requirement for long-term anticoagulation treatment.[2] Several studies have demonstrated the effectiveness of VSARR in patients with AR complicated by aortic root dilation following a repair of congenital heart defects.[2, 3] To date, VSARR has not been performed on TOF patients undergone no intervention. Here we described a case of a previously unoperated TOF patient undergoing VSARR and concurrent repair of TOF successfully.

Patient Profile

A 35-year-old male previously diagnosed to have TOF at the age of 5 but did not undergo any intervention for an unknown reason, presented with dyspnea after physical exercises and occasional cyanosis of the lips. He reported no symptoms of chest pain, palpitation, amaurosis, syncope, swelling of limbs, or squatting position.

Initial echocardiography showed unoperated TOF. A dilated right atrium measuring 52 mm in diameter was evident, while the right ventricle appeared hypertrophic with a thickness of 14 mm. The infundibulum of the right ventricular outflow tract appeared severely narrowed with a diameter of 8 mm. The pulmonary artery measured 18 mm in diameter and the pulmonary artery valves were slightly restrictive. The left ventricle appeared normal in size and the ejection fraction was 73%. The aorta seemed to have moved forward and 50% of its area covered the interventricular septum. Furthermore, there was dilation of the aortic root (55.4 mm) (**Fig.1**) and a mild central AR (VC = 5.7 mm)(**Fig.2**) . The right coronary artery showed evidence of an anomaly with a diameter of 7.5mm. Moreover, there was a secundum atrial septal defect showing bidirectional low-speed shunt and a 15 mm perimembranous ventricular septal defect (VSD). Given the above findings, surgery was recommended to correct the above abnormalities.

Cardiopulmonary bypass was established between the distal ascending aorta and the vena cavae after a median sternotomy. Myocardial protection strategies including systemic hypothermia at 32, antegrade cold cardioplegia, and consistent retrograde cardioplegia through coronary sinus were applied. The right ventricular outflow tract was incised longitudinally and the excessive infundibular muscle was resected. The VSD extending towards the infundibular and perimembranous portion of the right ventricle was clearly visualized. An autologous pericardium was harvested for the closure of the VSD and secured with continuous sutures. The aortic root was then transected to explore the aortic annulus and aortic valves. There was no thickening or calcification found, and all the leaflets appeared equally. Following that, both the coronary arteries were freed, and all the aortic sinuses with the entire ascending aorta were resected. A 3 mm high aortic valvuloplasty ring was made out of a 26 mm artificial vessel. Then, the valvuloplasty ring, aortic root, and the proximal end of the artificial vessel were assembled by continuous suturing to reconstruct the Valsalva sinus. An end-to-end anastomosis was performed between the ascending aorta and the distal end of the artificial vessel with running sutures. Openings of 8 mm in size were made on the artificial vessel and both the coronary arteries were then anastomosed to the openings with continuous suturing using 5-0 Prolene. An autologous pericardium was utilized as a patch to enlarge the pulmonary artery trunk. Partial closure of the atrial septal defect was performed using running sutures, leaving a small remaining 3 mm opening. An intraoperative transesophageal echocardiogram was also performed, revealing trivial central AR and an unobstructed left ventricular outflow tract. Postoperatively, the right ventricular systolic pressure measured 45 mmHg, the pulmonary arterial pressure was 18 mmHg, and the peripheral blood pressure was 88/45 mmHg. The patient recovered from the surgery uneventfully and remained well upon last follow-up.

Discussion

Aortic dilation is a common sequela among patients with an unrepaired TOF. Studies have suggested that aortic dilation may be attributed to longstanding volume overload of the ascending aorta as a result of right ventricular outflow tract obstruction and right-to-left shunting across the ventricular septal defect.[4]·[5] VSARR is a well-established procedure that corrects AR as a result of aortic dilation, with recent studies suggesting that the modified David V technique that preserves sinus-like geometry may improve annular

stabilization and hemostasis in the long term compared with those of remodeling techniques (Yacoub/David II).[2] Also, VSARR has also been successfully performed on patients with congenital heart defects.[2, 3] Adachi and colleagues have reported a case of TOF repair with VSARR on a patient who has previously undergone a left original Blalock-Taussig shunt during infancy.[6] Our report represents the first to describe a successfully performed VSARR procedure on a previously unoperated adult TOF patient. In our patient, the long-standing VSD fortunately has not caused a significant impairment to the aortic cusps or annulus. With our previous experience, extreme care was taken to avoid disrupting the balanced aortic root or distorting the aortic annulus during VSD repair. To better preserve the sinuses and considering the hemodynamics of this patient, we chose the Yacoub technique to reconstruct the aorta.

Conclusion

In conclusion, valve-sparing aortic root replacement using the Yacoub technique for aortic root dilation and severe AR, with a concomitant repair of tetralogy of Fallot in previously unoperated adult patients is feasible with excellent surgical outcomes. Studies examining long-term outcomes are warranted to validate the effectiveness of the VSARR technique further while exploring other innovative surgical approaches in the management of TOF patients with aortic dilation.

Declarations

Ethics approval and consent to participate

No institutional ethics committee approval was obtained since this was a case report. Only written and informed consent was obtained from the patient. No patient health identification was recorded.

Consent for publication

The patient himself provided written and informed consent for the publication of data.

Competing interests

The authors declare that there is no conflict of interest regarding this manuscript.

Funding

This study was funded by Technology innovation research and development project of Chengdu Science and Technology Bureau (No. 2019-YF05-00183-SN) and 1-3-5 project for disciplines of excellence—Clinical Research Incubation Project, West China Hospital, Sichuan University(2018HXFH021).

Authors' contributions

YL, QA and YQ performed the surgery; YL and CZ participated in gathering and interpreting the data, images and medical history; all three authors participated in preparing and polishing the manuscript; all authors have approved this version of manuscript.

Acknowledgements

Authors would like to express our gratitude to our excellent surgical team, as well as to our patient and his family for their kind support and cooperation.

References:

1. Grotenhuis HB, Dallaire F, Verpalen IM, van den Akker MJE, Mertens L, Friedberg MK: **Aortic Root Dilatation and Aortic-Related Complications in Children After Tetralogy of Fallot Repair** . *Circ Cardiovasc Imaging* 2018, **11** (12):e007611.
2. Fraser CD, 3rd, Liu RH, Zhou X, Patel ND, Lui C, Pierre AS, Jacobs ML, Dietz HC, Habashi J, Hibino N *et al* : **Valve-sparing aortic root replacement in children: Outcomes from 100 consecutive cases** . *J Thorac Cardiovasc Surg* 2019, **157** (3):1100-1109.

3. Baliulis G, Ropponen JO, Salmon TP, Kaarne MO: **Valve-sparing aortic root replacement in adult patients previously operated for congenital heart defects: an initial experience** . *Eur J Cardio-thorac Surg* 2016, **50** (1):155-159.
4. Niwa K: **Aortic dilatation in complex congenital heart disease** . *Cardiovasc Diagn Ther* 2018, **8** (6):725-738.
5. Seki M, Kuwata S, Kurishima C, Nakagawa R, Inuzuka R, Sugimoto M, Saiki H, Iwamoto Y, Ishido H, Masutani S *et al* : **Mechanism of aortic root dilation and cardiovascular function in tetralogy of Fallot** . *Pediatr Int* 2016, **58** (5):323-330.
6. Adachi O, Suzuki T, Yoshioka I, Takahashi G, Akiyama M, Kumagai K, Tatebe S, Saiki Y: **Concomitant Valve-Sparing Aortic Root Replacement With Repair of Tetralogy of Fallot** . *Ann Thorac Surg* 2019, **108** (2):e99-e101.



