

# Fructose 1,6-bisphosphatase and gluconeogenesis control in an anoxia-tolerant turtle, *Trachemys scripta elegans*

Aakriti Gupta<sup>1</sup>, Anchal Varma<sup>1</sup>, and Kenneth Storey<sup>1</sup>

<sup>1</sup>Carleton University Institute of Biochemistry

April 05, 2024

## Abstract

The red eared slider (*Trachemys scripta elegans*) undergoes numerous changes to its physiological and metabolic processes to survive without oxygen. During anoxic conditions, its metabolic rate drops drastically to minimise energy requirements. The alterations in central metabolic pathways are often accomplished by the regulation of key enzymes. The regulation of one such enzyme, fructose-1,6-bisphosphatase (FBPase; EC 3.1.3.11), was characterized in the present study during anoxia in liver. FBPase is a crucial enzyme of gluconeogenesis. FBPase was purified from liver tissue in both control and anoxic conditions and subsequently assayed to determine the kinetic parameters of the enzyme. Studies revealed the relative degree of posttranslational modifications in FBPase from control and anoxic turtles. Further, this study demonstrated a significant decrease in the maximal activity in anoxic FBPase and decreased sensitivity to its substrate fructose-1,6-bisphosphate (FBP) when compared to the control. Immunoblotting demonstrated increased threonine phosphorylation (~1.4 fold) in the anoxic FBPase. Taken together these results suggest that phosphorylation of liver FBPase is an important step in suppressing FBPase activity ultimately leading to inhibition of gluconeogenesis in the liver of the red eared slider during anaerobic conditions.

## Fructose 1,6-bisphosphatase and gluconeogenesis control in an anoxia-tolerant turtle, *Trachemys scripta elegans*

Aakriti Gupta<sup>1</sup>, Anchal Varma<sup>1</sup>, Kenneth B. Storey\*

Institute of Biochemistry & Department of Biology, Carleton University, 1125 Colonel By Drive, Ottawa, Ontario K1S 5B6, Canada

<sup>1</sup> indicates equal authorship

**\*Corresponding author:**

Dr. Kenneth B. Storey

Institute of Biochemistry and Department of Biology,

1125 Colonel By Drive, Ottawa, ON, K1S 5B6

Tel: (613) 520-2600, ext. 3678

E-mail: KennethStorey@cunet.carleton.ca

## Abstract

The red eared slider (*Trachemys scripta elegans*) undergoes numerous changes to its physiological and metabolic processes to survive without oxygen. During anoxic conditions, its metabolic rate drops dras-

tically to minimise energy requirements. The alterations in central metabolic pathways are often accomplished by the regulation of key enzymes. The regulation of one such enzyme, fructose-1,6-bisphosphatase (FBPase; EC 3.1.3.11), was characterized in the present study during anoxia in liver. FBPase is a crucial enzyme of gluconeogenesis. FBPase was purified from liver tissue in both control and anoxic conditions and subsequently assayed to determine the kinetic parameters of the enzyme. Studies revealed the relative degree of posttranslational modifications in FBPase from control and anoxic turtles. Further, this study demonstrated a significant decrease in the maximal activity in anoxic FBPase and decreased sensitivity to its substrate fructose-1,6-bisphosphate (FBP) when compared to the control. Immunoblotting demonstrated increased threonine phosphorylation (~1.4 fold) in the anoxic FBPase. Taken together these results suggest that phosphorylation of liver FBPase is an important step in suppressing FBPase activity ultimately leading to inhibition of gluconeogenesis in the liver of the red eared slider during anaerobic conditions.

**Keywords :** *Trachemys scripta elegans* , fructose-1,6-bisphosphatase, enzyme regulation, gluconeogenesis, anoxia, metabolic rate depression, post-translational modification

**Abbreviations:** FBPase: Fructose-1,6-bisphosphatase, FBP: fructose-1,6-bisphosphate, PFK: phosphofructokinase, PEP: Phosphoenolpyruvate

## Introduction

The red-eared slider turtle, *Trachemys scripta elegans* , is one of the best-studied models of facultative anaerobiosis among vertebrate species. These turtles have the ability to survive for several months in oxygen deprived conditions when hibernating in cold water at temperatures of 0-5°C<sup>1</sup>. In contrast to mammals that show extreme sensitivity to brief anoxia exposures, turtles have physiological and biochemical adaptations to endure prolonged anoxia, including high glycogen reserves, antioxidant defences<sup>2</sup>, metabolic rate depression, and an ability to buffer high concentrations of lactic acid generated by anaerobic glycolysis by storing lactate into the shell<sup>3</sup>. Metabolism is reduced to about 10-20% of aerobic values<sup>4</sup> causing tissues to switch to a reliance on glycolysis alone for ATP production. Maintaining cellular energetics is one of the major concerns under anoxic conditions. This is because the conversion of glucose to lactate via glycolysis yields only 2 ATP per glucose molecule catabolized compared with the full aerobic oxidation of glucose to CO<sub>2</sub> and H<sub>2</sub>O that yields 36 ATP per glucose<sup>5</sup>. Therefore, studies of the molecular mechanisms that support long term anoxia tolerance are important for understanding this remarkable phenomenon.

One major contributor to anoxia tolerance is metabolic rate depression, a coordinated process that leads to the strong and coordinated suppression of most metabolic processes. Biochemical mechanisms contributing to metabolic arrest include changes in the enzyme specificity, posttranscriptional and posttranslational modifications, and reduced rates of protein biosynthesis<sup>5,6</sup>. Studies of stress responsive changes in the activities, kinetic properties, and regulatory mechanisms of enzymes involved in carbohydrate catabolism have provided insights into the metabolic adjustments used by turtles to suppress glycolysis under anoxic conditions<sup>7-9</sup>. These studies demonstrated organ specific changes in the kinetics and activities of enzymes including glycogen phosphorylase, phosphofructokinase, pyruvate kinase, lactate dehydrogenase, and fructose bisphosphate aldolase under anoxic conditions<sup>8-12</sup>. In addition, anoxia also triggered tissue specific changes in the activities of many other metabolic enzymes in *T. s. elegans*<sup>2</sup>. These studies have documented that global control of glycolytic and other enzymes is vital for turtles to adapt during oxygen depriving condition<sup>13-17</sup>.

During anaerobiosis, turtles utilize large reserves of liver glycogen for energy production. Hepatic glycogenolysis raises blood glucose levels and delivers glucose to other organs. Glucose can also be synthesized from noncarbohydrate precursors (such as amino acids) or during clearance of accumulated lactate by a process called gluconeogenesis that occurs primarily in liver. Limited work has been done with respect to the role that gluconeogenesis might play in fuel production in anoxia tolerant animals and further research in this area is warranted. Fructose-1,6-bisphosphatase (FBPase; EC 3.1.3.11) is a key regulatory enzyme in the gluconeogenic pathway and catalyzes the reverse reaction to that of phosphofructokinase (PFK) in glycolysis (Fig.1). Specifically, FBPase catalyzes the removal of a phosphate from fructose-1,6-bisphosphate to form fructose-6-phosphate, in the presence of metal ions, magnesium or manganese. FBPase and PFK act as

a rate-limiting enzymes to control the gluconeogenic versus glycolytic pathways, respectively, resulting in regulation of glucose homeostasis and energy metabolism<sup>18</sup>. FBPase is an allosteric enzyme (Fig. 1). Its activity is stimulated by high ATP concentrations whereas AMP and fructose 2,6-biphosphate are negative regulators<sup>19</sup>. Regulation of FBPase is also accomplished by reversible protein phosphorylation<sup>20,21</sup>. This strict regulatory network of FBPase and its importance in the gluconeogenic pathway makes it an intriguing enzyme to study in turtles, that accumulate huge amounts of lactate under anoxic conditions. The present investigation aims to explore the physical, kinetic, and regulatory properties of liver FBPase in an attempt to determine its role in anoxia survival.

## Materials and Methods

### Animals

Adult female red-eared sliders (*T. s. elegans*), 700–1,500 g, were acquired from a local supplier (Ottawa, Ontario) and held at  $5 \pm 1$  °C in large 50 L plastic tanks filled with dechlorinated tap water for 7–8 days before use. Control (normoxic) turtles were sampled from this condition. For anoxia exposure, turtles were transferred to large buckets containing water at  $5 \pm 1$  °C that had been previously bubbled for 1 h with N<sub>2</sub> gas via a port in the lid of the bucket. Two turtles were added per bucket and a wire mesh was fitted into the bucket about 5 cm below the water surface so that turtles remained submerged throughout the 20 h experimental anoxia exposure. Bubbling with nitrogen gas was continued for 1 h after turtles was added and was reinitiated again during sampling of the animals. All animals were killed by decapitation and liver tissue was rapidly dissected out, frozen in liquid nitrogen and stored at -80 °C until use. All animals were cared for in accordance with the guidelines of the Canadian Council on Animal Care and experimental procedures had the prior approval of the Carleton University Animal Care Committee (protocol #106937).

### Sample preparation

Samples of liver tissue from control and 20 h anoxic turtles were rapidly weighed and homogenised in ice-cold homogenization buffer A (25 mM MES buffer, pH 6.25 mM NaF, 2.5 mM EDTA, 2.5 mM EGTA, 5 mM  $\beta$ -mercaptoethanol and 10% v/v glycerol) in the ratio 1:5 w:v. A few crystals of phenylmethylsulphonyl fluoride were added to the buffer immediately before the homogenization. Samples were homogenised using a Polytron homogenizer (Brinkmann Instruments, Westbury, NY, USA) followed by centrifugation at 13,500 X g at 4 °C for 30 minutes. The supernatant was collected as crude homogenate and stored on ice.

### FBPase purification

FBPase was purified using a two-column procedure. A phosphocellulose column was prepared and equilibrated with 30 ml of buffer A. Supernatant was then loaded onto the column followed by washing with 40 mL of buffer A. FBPase then eluted using 30 ml of a 0 to 1 M KCl gradient containing 2.5 mM fructose-1,6-bisphosphate (FBP) in buffer A. Fractions were collected using an automated fraction collector (Gilson Medical Electronics, Inc., Middletown, WI, USA). The highest activity fractions were pooled and loaded on a carboxymethyl (CM) cellulose column pre-equilibrated with buffer A. The column was then washed with 20 ml of buffer A and the enzyme was eluted using a 30 ml linear salt gradient of 0 to 1 M KCl. The fractions with highest FBPase activity were pooled and then analyzed by sodium dodecyl sulfate polyacrylamide gel electrophoresis (SDS-PAGE) to demonstrate purity of the FBPase preparation.

### SDS PAGE and Coomassie staining

The purity of FBPase was confirmed by running SDS-PAGE using a protocol described previously by Smolinski *et. al.*<sup>22</sup>. Samples of FBPase for SDS-PAGE were prepared using equal aliquots of purified enzyme and SDS loading buffer mixed (100 mM Tris buffer, pH 6.8, 4% w:v SDS, 20% v:v glycerol, 0.2% w:v bromophenol blue) with 10% v:v  $\beta$ -mercaptoethanol added fresh. Samples were boiled for 5 minutes, then cooled on ice and stored in a freezer at -20 °C. Subsequently, enzyme samples were loaded into wells of 10% gel and run adjacent to a low molecular weight ladder (Pink Plus Prestained Protein Ladder, Frogga Bio, Toronto,

ON, catalog no. PM005–0500) for 55 minutes at 180 V. Silver staining was used following the procedure of Celiset. *al.*<sup>23</sup>. After the bands were clearly visible, the gel was imaged using Chemi-Genius Bio-Imaging system with GeneSnap software (Syngene, Frederick, MD). FBPase was further identified by excising the bands obtained from electrophoresis and sending them for mass spectrometry at the Proteomics Platform of the Quebec Genomics Center (Quebec, Canada).

### FBPase Assay

FBPase was assayed by monitoring NADP<sup>+</sup> reduction at 340 nm using a microplate reader (Multiskan Spectrum, Thermo Labsystems, Finland). Optimal assay conditions were 15 mM imidazole-HCl buffer, pH 7.2, 0.1 mM fructose-1,6-bisphosphate, 5 mM MgSO<sub>4</sub>, 0.2 mM NADP<sup>+</sup>, and 1 U each of phosphoglucosomerase and glucose-6-phosphate dehydrogenase in a total volume of 200  $\mu$ l. All assays were performed at 22 °C.  $K_m$  values were calculated using the above-mentioned assay conditions but with suboptimal concentrations of FBP (0.24  $\mu$ M).  $I_{50}$  values for AMP and ADP and  $K_a$  values for alanine, phosphoenolpyruvate (PEP) and ATP.Mg (1:1) were also determined using similar assay methods.

Soluble protein concentrations were determined using the Coomassie blue dye-binding method with the BioRad Laboratories prepared reagent and bovine serum as the standard. Spectrophotometric quantification was at 595 nm.

### Posttranslational Modification analysis via Western blotting

Posttranslational modifications (PTMs) were assessed via immunoblotting. The protocol followed was as in Gupta and Storey, 2020<sup>24</sup>. Following electrophoresis (as described above), proteins were transferred to a polyvinylidene difluoride (PVDF) membrane by electroblotting at a constant current of 160 mA for 90 min under transfer buffer (25 mM Tris, pH 8.5, 192 mM glycine, 10% v/v methanol). After transfer membranes were washed for 3  $\times$  5 min using TBST (Tris-buffered saline with Tween-20): 20 mM Tris base pH 7.6, 150 mM NaCl, 0.05% v/v Tween-20). Following washing, membranes were probed with primary antibody, diluted 1:1000 v/v in TBST with overnight incubation at 4 °C on a rocker.

Polyclonal antibodies were used to detect: (a) serine phosphorylation (catalogue# 618100); (b) threonine phosphorylation (catalog# 718200); (c) tyrosine phosphorylation (catalog# 136600), were purchased from Invitrogen, Carlsbad, CA, USA, (d) ubiquitination (catalog# ab19247 Abcam, Cambridge, UK); and (e) SUMOylation 1 (a gift from the Clinical Investigations Section Stroke Branch, NINDS, Bethesda, MD, USA).

After incubation, membranes were washed for 3  $\times$  5 min in TBST at room temperature and then incubated with secondary antibody (anti-rabbit IgG secondary antibody conjugated with horseradish peroxidase (catalog no. APA007P.2, Bioshop, Burlington, ON, Canada)) at 1:5000 v:v in TBST for 30 minutes. Membranes were again washed 3 times with TBST for 5 minutes per wash. Following the washes, immunoreactive bands on the membrane was detected using 600  $\mu$ l each of luminol and hydrogen peroxide and imaged using the ChemiGenius Bioimaging System (Syngene, MD, USA). Membranes were subsequently stained with Coomassie blue (0.25% w/v Coomassie brilliant blue, 7.5% v/v acetic acid, and 50% methanol) and the summed intensities of a group of Coomassie stained bands that were well separated from the band of interest and constant across all lanes were used to standardize the chemiluminescence of the immunobands of interest. Band intensities were quantified using GeneTools software (Syngene, Frederick, MD).

### *In silico* FBPase 3D homology modelling

The amino acid sequence of red-eared slider turtle FBPase (Genbank XP\_034629682.1) was used to predict the 3D structure of the enzyme using a homology modelling 3D protein structure prediction tool SWISS-MODEL (<https://swissmodel.expasy.org/>). Turtle FBPase was modelled against the X-ray crystal structure of human liver FBPase as it had the highest sequence coverage as compared with the turtle enzyme. Structural validations were performed using SWISS-MODEL and the structural assessment tool was used to analyse QMEAN scores, Ramachandran plots and MolProbity quality scores. Visualisation of the FBPase structure

in the presence of FBP and Mg was done using PyMOL Molecular Graphics System, Version 2.0 Schrödinger, LLC.

## Quantification and Statistical Analysis

All enzyme assays were analysed using a microplate analysis program (MPA) and enzyme kinetic parameters ( $K_m$ ,  $V_{max}$ ,  $I_{50}$ ,  $K_a$ ) were determined using a least squares regression computer program, Kinetics v. 3.5.1<sup>22</sup>. Data for western blots and kinetic parameters were analyzed using one-way analysis of variance (ANOVA), followed by Student's t-test or Tukey's post hoc test with  $p < 0.05$  accepted as a significant difference.

## Results

### Purification of Fructose 1,6 bisphosphatase

FBPase from *T.s. elegans* liver was purified to homogeneity in a two-step chromatographic process. The first step, elution of FBPase from a phosphocellulose column used a 0-1M KCl linear gradient with 2.5 mM FBP added. This was effective at removing many impurities and resulted in a 3.80-fold purification with 51.3% activity recovered (Table 1). The second step used a CM column and eluted FBPase using a 0-1 M KCl linear gradient resulting in a final fold purification of 8.7 and an 24.8% yield of enzyme activity (Table 1). Using SDS-PAGE followed by staining with silver demonstrated that the FBPase was purified to homogeneity. The purified FBPase was calculated to be ~37 kDa based on electrophoretic mobility compared to the protein standards loaded (Fig.2).

### Kinetic properties of FBPase

Enzymatic parameters were measured using purified FBPase from control and anoxia exposed turtles. The  $V_{max}$  of the enzyme from anoxic liver was significantly lower by 26% as compared with control liver. The enzymes also differed in multiple kinetic parameters (Table 2). The  $K_m$  FBP of FBPase increased significantly by 2.2 fold for the enzyme from anoxic liver, rising from 26.9  $\mu$ M for the control enzyme to 59.8  $\mu$ M for FBPase from anoxic liver. The effects of inhibitors on the relative activities of control and anoxic liver FBPase were also determined (Table 2). There were significant decreases in  $I_{50}$  values between control and anoxic FBPase for both ADP and AMP effects. The  $I_{50}$  for ADP inhibition of anoxic liver FBPase ( $8.3 \pm 0.6 \mu$ M) was significantly lower by 30% as compared to the control value ( $I_{50} = 11.8 \pm 0.3 \text{ mM}$ ). Similarly, a 41% decrease compared to control was observed for the  $I_{50}$  value for anoxic FBPase in presence of AMP, with the control AMP  $I_{50}$  being  $6.1 \pm 0.1 \mu$ M whereas anoxic FBPase showed an AMP  $I_{50}$  of  $3.6 \pm 0.3 \mu$ M (Table 2).

ATP.Mg, PEP and alanine were also investigated as activators of FBPase (Table 2). For ATP.Mg, the  $K_a$  value was significantly lower in anoxia ( $25 \pm 0.9 \mu$ M) as compared to that of controls ( $144.5 \pm 7.0 \mu$ M). By contrast,  $K_a$  of PEP was significantly higher in anoxia ( $3.3 \pm 0.2 \mu$ M) than in controls ( $0.33 \pm 0.09 \mu$ M). The activity of the anoxic enzyme in the presence of PEP decreased to 0.3% compared to control FBPase.  $K_a$  for L-alanine did not show any significant difference between the two conditions (Table 2).

### PTM western blots

Analysis of relative levels of PTMs was done using western blotting (Fig.3). Phosphorylation of threonine residues showed a 1.4-fold increase in FBPase from anoxic liver as compared to controls. However, SUMOylation and ubiquitination showed significant decrease to 73% and 22% of the control values respectively (Fig.3).

### Bioinformatics

A predicted 3D structure of red-eared slider turtle FBPase was generated using SWISS-MODEL. The primary sequence of the *T.s. elegans* enzyme was obtained from the NCBI (XP\_034629682.1) and the crystal structure of human liver FBPase was used as the template to obtain a structural model of the turtle enzyme (Fig.4A).

Visualisation of the FBPase structure in the presence of FBP substrate (red) and Mg (yellow) was done using Pymol and shows the position of these substrates at the active site (Fig.4B). The MolProbity structure-validation web service, a component of the SWISS-MODEL structure assessment tool, was used to provide an evaluation of model quality at both the global and local protein level. A low global MolProbity score of 1.33 reflects that crystallographic resolution is at a good, expected quality. The generated Ramachandran plot was analyzed with MolProbity that showed a Ramachandran favoured score of 95.62%, close to the ideal score of >98% (Fig.4C). The QMEAN score, a composite estimator of geometrical properties at the global and local protein level, was calculated to be 0.35 (Fig.4D). This score considers the C $\beta$  atoms interaction energy of -0.85, all-atom pairwise energy of -0.21, solvation energy of 1.22, and torsion angle energy of 0.09 (Fig.4D). Collectively these physics and statistical knowledge-based modelling scores are all near zero which indicates a model that is in the range of experimental models of the same size.

## Discussion

A lack of oxygen is a major challenge to the survival of many animals. The red-eared slider, *T.s. elegans*, is an anoxia tolerant species and can survive without oxygen for months during winter when hibernating in cold water<sup>1,25</sup>. Turtles also experience oxygen deprivation during breath-hold diving at all seasons and this escalates to prolonged periods of anoxia during overwintering in ice-locked ponds<sup>26</sup>. Under anoxic conditions, turtles rely on their glycogen reserves (stored mainly in liver) for energy and switch to lactic acid fermentation<sup>27</sup>. In addition, they lower their metabolic rate to only about 10% of normal<sup>9</sup>, making them model organisms to study remodelling of energy metabolism for extended periods of time<sup>28</sup>. Profound metabolic rate depression enable survival without oxygen for several months at cold temperatures. By rearranging key cellular processes, turtles can redirect available fuel/energy resources to prioritise pro-survival processes that are important during anoxia. Considering that glycogen is metabolised under anoxic conditions as an fuel/energy source, turtles must pre-load their organs with high glycogen stores during summer/autumn feeding<sup>7</sup>. Therefore, gluconeogenesis could be one of the active processes occurring under aerobic conditions. Gluconeogenesis is also important for the recovery of internal glycogen reserves when turtles transition back from prolonged anoxia to air-breathing. FBPase is the rate-limiting enzyme of gluconeogenesis. A study of its enzymatic mechanism would provide better insight into the metabolism restructuring by turtles under anoxic conditions. Indeed, the present study showed that liver FBPase from aerobic control and anoxic *T. s. elegans* displayed markedly different properties and further investigation linked these differences in enzyme properties with anoxia-responsive covalent modification of the enzyme.

The maximum activity of FBPase (expressed as U/mg protein) purified from liver of anoxic turtles was reduced significantly as compared to that of the control enzyme from aerobic turtles. This lower FBPase  $V_{max}$  suggests that a stable modification of the enzyme occurred in response to anoxia, potentially reducing FBPase function under anoxic cellular conditions where increased glycolysis is needed when ATP generation by oxidative phosphorylation is blocked. Liver FBPase also showed significant differences in kinetic parameters between control and anoxic turtles.  $K_m$  FBP was 2.2 fold higher in anoxic turtles as compared to controls (Table 2). This represents a significant decrease in substrate affinity and is indicative of a less active FBPase during anoxia. The results obtained here correspond well with the findings of previous studies. For example, Warren and Jackson<sup>3</sup> reported that under anoxia, gluconeogenesis is suppressed due to high energy costs. Gluconeogenesis might not be important during the early hours of anoxia; however, it would be required during the recovery from anoxia, when large amounts of lactate must be cleared<sup>6,11,12</sup>.

The activity of FBPase was negatively modulated by the allosteric inhibitors AMP and ADP<sup>29</sup>. Table 2 shows the  $I_{50}$  values for ADP and AMP.  $I_{50}$  means the concentration of an inhibitor that is needed to reduce enzyme activity by 50%. So, the current data represents that it takes an AMP concentration of 11.8  $\mu$ M to reduce the FBPase activity of the control enzyme by 50% but significantly less (8.3  $\mu$ M) to reduce the activity of anoxic FBPase. This shows that ADP is a stronger inhibitor of the anoxic form of FBPase as compared with the aerobic form. So, if AMP or ADP levels rise in anoxia (which they do) their higher levels

will have a stronger effect on the activity of anoxic FBPase than on aerobic FBPase. Such differences in the power of inhibitors to control an enzyme are often due to PTMs that change the conformation of the enzyme protein and thereby allow greater or lesser access to the enzyme by substrate or allosteric modulators.

Previous studies of liver FBPase from the little brown bat (*Myotis lucifugus*)<sup>30</sup> and white muscle of freshwater rainbow trout (*Oncorhynchus mykiss*)<sup>31</sup> showed similar responses to ADP and AMP under stress conditions<sup>30,32,33</sup>. The net effect of increased inhibition of anoxic FBPase in liver when compared to controls would result in the suppression of enzymatic activity under stress conditions. Furthermore, increased inhibition by ADP and AMP indicates that activity is closely regulated by adenylate energy status. FBPase being the rate-limiting enzyme of gluconeogenesis, adenylate levels would be important in controlling the rate of gluconeogenesis from substrates including glycerol-3-phosphate (resulting from triglyceride oxidation) or amino acids (from protein degradation) or lactate from glycogen catabolism<sup>34</sup>. This suggests that sensitivity to AMP and ADP was increased under the anoxic condition, so that activities of energy-expensive pathways such as gluconeogenesis would be suppressed to prevent fruitless carbon cycling and wasteful ATP hydrolysis under energy-limiting conditions. Further analysis of liver FBPase activity also showed its sensitivity to ATP-Mg, the enzyme from anoxic turtles showing a ~6-fold increase in its sensitivity to ATP as determined by a significantly decreased  $K_a$  value of the anoxic liver FBPase (Table 2). A previous study on anoxic turtle liver showed a decrease in glycolytic intermediates during anoxia and inhibition of glycolysis at the phosphofructokinase step to facilitate glycogenolysis<sup>33</sup>. Inhibition of glycolysis is also correlated with metabolic rate depression. Inhibiting both glycolytic rate (via PFK) and gluconeogenesis rate (via FBPase) would reduce overall carbohydrate metabolism both substantially and in a coordinated fashion. Therefore, the increase in affinity of ATP-Mg in the anoxic liver FBPase does not necessarily indicate an active anoxic FBPase. In fact, a study on turtle liver FBPase showed that anoxia had no effect on the total activity of FBPase, suggesting that FBPase activity might be regulated at the posttranslational level<sup>35</sup>. Current results suggest that initiation of gluconeogenesis would be more energy efficient under aerobic conditions to breakdown the lactate accumulated during anoxia. As such, perhaps FBPase activity is required during early stages of recovery; therefore, increased sensitivity towards ATP-Mg could hasten gluconeogenesis during the early hours of recovery.

Additional analysis of liver FBPase was done to assess effects of other putative modulators of enzyme activity. Phosphoenolpyruvate (PEP) was found to be the activator of liver FBPase.  $K_a$  values significantly increased in the presence of PEP in the anoxic liver compared to control values (Table 2). Flux through glycolysis is rerouted at the level of PEP. PEP concentration could coordinate regulation of FBPase and Phosphofructokinase (PFK) in one of the two ways: 1. gluconeogenesis is favored when the cell is rich in biosynthetic precursors and ATP, or 2. high concentrations of PEP can activate FBPase. It is apparent from previous work that short term control of hepatic gluconeogenesis by glucagon and insulin may be of physiological importance<sup>36</sup>. Studies on rat liver suggest that the carboxylation of pyruvate, the pyruvate kinase reaction, and the fructose 6 phosphate/fructose 1,6 bisphosphate cycle have been forwarded as probable regulatory sites of acute hormonal control of gluconeogenesis<sup>37,38</sup>. Pyruvate kinase is relatively the most important locus of endocrine control<sup>39,40</sup>. It is well established that glucagon exerts its effect at this level by stimulating cyclic AMP-dependent phosphorylation of pyruvate kinase<sup>41</sup> that decreases PK affinity for PEP, thereby decreasing flux through pyruvate kinase despite increased PEP to pyruvate concentration ratios during gluconeogenesis. As a consequence futile cycling between PEP and pyruvate is reduced and gluconeogenesis is effectively promoted<sup>42</sup>. Decrease in affinity of PEP towards the anoxic liver FBPase may suggest decrease in activity of enzyme towards PEP. The results obtained here further suggest another physiological function of FBPase, that is to prevent futile cycling during gluconeogenesis. Taken together, changes in substrate affinity and susceptibility to allosteric inhibitors and activators may be coordinately regulated to result in an overall decrease in FBPase function as part of anoxia-induced hypometabolism although still maintaining sufficient activity to support basal ATP production by anaerobic glycolysis.

From the data obtained on kinetic parameters it was apparent that structural differences might exist between the control and anoxic forms of FBPase. The potential reasons for structural differences could be PTMs, especially phosphorylation. Protein phosphorylation is known to be a recurrent way to reversibly

regulate enzyme activity in response to environmental stress in many animals<sup>43</sup>. Immunoblotting showed a significant increase of 1.4 fold in phosphorylation of threonine residues on FBPase from liver of anoxic turtles as compared with the enzyme from control turtles (Fig. 3). Threonine phosphorylation has been reported to suppress enzyme activities during anoxia and other forms of hypometabolism. For example, previous studies on the marine mollusc, *Busycon canaliculatum*<sup>44</sup> and the red eared-slider<sup>45</sup>, showed a correlation between increased pyruvate kinase phosphorylation and changes in enzyme properties that would suppress enzyme function. Furthermore, based on electrophoretic analysis of [<sup>32</sup>P] incorporation, it was suggested that O-phosphothreonine residue(s) were produced under anoxia<sup>46</sup>. By contrast, posttranslational modification by SUMOylation and ubiquitination decreased significantly to 22% and 73% of the control values during anoxia. In general, ubiquitination leads to protein degradation<sup>47</sup>. Furthermore, it has been shown that reduced ubiquitination stabilizes oxidized proteins<sup>48,49</sup>. Hence, under anoxic conditions, a decrease in ubiquitination levels can suggest an increase in enzyme stability in the energy-limited anoxic state. Similar to ubiquitination, the function of protein SUMOylation is less well understood. One study, showed that SUMOylation contributed to activation of pro-survival pathways during stress<sup>50</sup>. A study of muscle lactate dehydrogenase from *T.s. elegans* showed reduced ubiquitination and SUMOylation of the enzyme<sup>51</sup> in response to anoxia which further supports our findings for liver FBPase. While the consequences of ubiquitination and SUMOylation of turtle FBPase are still unknown, previous research has identified these PTMs as being crucial regulators of enzyme activity, protein-protein interactions, and protein stability<sup>52</sup>.

This evidence of altered FBPase function under anoxia is in accordance with comparable findings of anoxia-responsive changes in the properties of other enzymes of carbohydrate metabolism in turtle organs including glycogen phosphorylase, phosphofructokinase, and pyruvate kinase<sup>7,8</sup>. The anoxia-induced suppression of key energy-expensive processes (e.g. ion pumping, protein translation, urea synthesis, etc.) in isolated turtle hepatocytes<sup>53</sup> is also well documented. In these and other cases, including altered transcription factor responses under anoxia, the mechanism involved appear to be covalent modification by reversible protein phosphorylation<sup>54,55</sup>. These modifications at the binding sites of the enzyme can affect enzyme activity and sensitivity towards effectors under different conditions. Exposure to anoxia and long-term survival in an anoxic state requires regulatory mechanisms that enable a primary focus on only pro-survival processes under hypometabolic conditions. Being reversible processes, PTMs provide an economical solution to alter enzyme activity as opposed to more radical energy-expensive solutions such as proteolysis/resynthesis of enzymes over the anoxia/reoxygenation cycle. Changes in enzyme activities and catalytic properties due to PTMs in response to environmental stresses have been observed in multiple previous studies conducted on both vertebrate and invertebrate species<sup>22,45,56–58</sup>. Considering the above results, and previous studies, it can be said that PTMs are an important factor behind the differential activity of an enzyme under aerobic versus anoxic conditions.

## Conclusion

The data presented here show that turtle liver FBPase undergoes several significant changes in its functional and structural properties in response to anoxia. Differences in PTMs were linked with changes in the kinetic properties of FBPase. In addition to that the anoxic FBPase showed lower Vmax and decreased affinity for substrate as compared to control liver FBPase. The current study asserts posttranslational control over gluconeogenesis which further uphold the previous studies of metabolic responses to oxygen deprivation by anoxia tolerant species. It also provides new insights on metabolic reorganization that freshwater turtles undergo to cope with anoxia.

## Acknowledgment

The research was supported by a Discovery Grant (Grant # 6793) from the Natural Sciences and Engineering Research Council (NSERC) of Canada; KBS holds the Canada Research Chair in Molecular Physiology. We thank JM Storey for the editorial review of this manuscript.

## Conflict of interest

The authors declare no conflict of interest.

## References

1. Ultsch GR. The viability of nearctic freshwater turtles submerged in anoxia and normoxia at 3 and 10°C. *Comp Biochem Physiol Part A Physiol* . 1985;81(3):607-611. doi:10.1016/0300-9629(85)91035-7
2. Breedon SA, Hadj-Moussa H, Storey KB. Nrf2 activates antioxidant enzymes in the anoxia-tolerant red-eared slider turtle, *Trachemys scripta elegans* . *J Exp Zool Part A Ecol Integr Physiol* . 2021;335(4):426-435. doi:10.1002/jez.2458
3. Warren DE, Jackson DC. Lactate metabolism in anoxic turtles: an integrative review. *J Comp Physiol B* . 2008;178(2):133-148. doi:10.1007/s00360-007-0212-1
4. Storey KB. Anoxia tolerance in turtles: Metabolic regulation and gene expression. *Comp Biochem Physiol - A Mol Integr Physiol* . 2007. doi:10.1016/j.cbpa.2006.03.019
5. Storey KB. Molecular mechanisms of anoxia tolerance. *Int Congr Ser* . 2004;1275:47-54. doi:10.1016/j.ics.2004.08.072
6. Krivoruchko A, Storey KB. Turtle anoxia tolerance: Biochemistry and gene regulation. *Biochim Biophys Acta - Gen Subj* . 2015;1850(6):1188-1196. doi:10.1016/j.bbagen.2015.02.001
7. Brooks SPJ, Storey KB. Regulation of glycolytic enzymes during anoxia in the turtle *Pseudemys scripta* . *Am J Physiol Integr Comp Physiol* . 1989;257(2):R278-R283. doi:10.1152/ajpregu.1989.257.2.R278
8. Xiong ZJ, Storey KB. Regulation of liver lactate dehydrogenase by reversible phosphorylation in response to anoxia in a freshwater turtle. *Comp Biochem Physiol Part B Biochem Mol Biol* . 2012;163(2):221-228. doi:10.1016/j.cbpb.2012.06.001
9. Willmore WG, Cowan KJ, Storey KB. Effects of anoxia exposure and aerobic recovery on metabolic enzyme activities in the freshwater turtle *Trachemys scripta elegans* . *Can J Zool* . 2001;79(10):1822-1828. doi:10.1139/z01-149
10. Dawson NJ, Biggar KK, Storey KB. Characterization of Fructose-1,6-Bisphosphate Aldolase during Anoxia in the Tolerant Turtle, *Trachemys scripta elegans* : An Assessment of Enzyme Activity, Expression and Structure. Dzeja P, ed. *PLoS One* . 2013;8(7):e68830. doi:10.1371/journal.pone.0068830
11. Mehrani H, Storey KB. Enzymatic control of glycogenolysis during anoxic submergence in the freshwater turtle *Trachemys scripta* . *Int J Biochem Cell Biol* . 1995;27(8):821-830. doi:10.1016/1357-2725(95)00042-N
12. Storey KB. Metabolic adaptations supporting anoxia tolerance in reptiles: Recent advances. In: *Comparative Biochemistry and Physiology - B Biochemistry and Molecular Biology* . ; 1996. doi:10.1016/0305-0491(95)02043-8
13. Willmore WG, Storey KB. Antioxidant systems and anoxia tolerance in a freshwater turtle *Trachemys scripta elegans* . *Mol Cell Biochem* . 1997;170(1-2):177-185. doi:10.1023/a:1006817806010
14. Brooks SP, Storey KB. Glycolytic Enzyme Binding and Metabolic Control in Estivation and Anoxia in the Land Snail *Otala Lactea* . *J Exp Biol* . 1990;151(1):193-204. doi:10.1242/jeb.151.1.193
15. Brooks SPJ, Storey KB. Chapter 13 Is glycolytic rate controlled by the reversible binding of enzymes to subcellular structures? In: *Biochemistry and Molecular Biology of Fishes* . ; 1995:291-307. doi:10.1016/S1873-0140(06)80016-0
16. Brooks SPJ, Storey KB. Properties of Pyruvate Dehydrogenase from the Land Snail, *Otala lactea* : Control of Enzyme Activity during Estivation. *Physiol Zool* . 1992;65(3):620-633. doi:10.1086/physzool.65.3.30157973

17. Storey KB, Brooks SPJ. Chapter 6 The basis of enzymatic adaptation. In: *Principles of Medical Biology* . ; 1995:147-169. doi:10.1016/S1569-2582(06)80008-5
18. Benkovic SJ, DeMaine MM. Mechanism of Action of Fructose 1,6-Bisphosphatase. In: *Advances in Enzymology and Related Areas of Molecular Biology* . ; 2006:45-82. doi:10.1002/9780470122983.ch2
19. Hanson RW, Owen OE. Gluconeogenesis. In: Lennarz WJ, Lane MD, eds.*Encyclopedia of Biological Chemistry* . Elsevier; 2013:381-386. doi:10.1016/B978-0-12-378630-2.00040-2
20. Ekdahl KN. In vitro phosphorylation of fructose-1,6-bisphosphatase from rabbit and pig liver with cyclic AMP-dependent protein kinase.*Arch Biochem Biophys* . 1988;262(1):27-31. doi:10.1016/0003-9861(88)90164-6
21. Rakus D, Zarzycki M, Dzugaj A. Rabbit muscle fructose-1,6-bisphosphatase is phosphorylated in vivo. *Acta Biochim Pol* . 2003;50(1):115-121. doi:10.18388/abp.2003\_3718
22. Smolinski MB, Varma A, Green SR, Storey KB. Purification and Regulation of Pyruvate Kinase from the Foot Muscle of the Anoxia and Freeze Tolerant Marine Snail, *Littorina littorea* . *Protein J* . 2020;39(5):531-541. doi:10.1007/s10930-020-09934-9
23. Celis J, Trentemolle S, Gromov P. Gel-Based Proteomics High-Resolution Two-Dimensional Gel Electrophoresis of Proteins. Isoelectric Focusing and Nonequilibrium pH Gradient Electrophoresis. In: *Cell Biology* . Elsevier; 2006:165-174. doi:10.1016/B978-012164730-8/50207-0
24. Gupta A, Storey KB. Regulation of antioxidant systems in response to anoxia and reoxygenation in *Rana sylvatica* . *Comp Biochem Physiol Part B Biochem Mol Biol* . 2020;243-244:110436. doi:10.1016/j.cbpb.2020.110436
25. Jackson DC. Living without oxygen: lessons from the freshwater turtle. *Comp Biochem Physiol Part A Mol Integr Physiol* . 2000;125(3):299-315. doi:10.1016/S1095-6433(00)00160-4
26. Jackson DC. Hibernating without Oxygen: Physiological Adaptations of the Painted Turtle. *J Physiol* . 2002;543(3):731-737. doi:10.1113/jphysiol.2002.024729
27. Warren DE, Reese SA, Jackson DC. Tissue Glycogen and Extracellular Buffering Limit the Survival of Red-Eared Slider Turtles during Anoxic Submergence at 3degC. *Physiol Biochem Zool* . 2006;79(4):736-744. doi:10.1086/504617
28. Bundgaard A, James AM, Gruszczuk A V., Martin J, Murphy MP, Fago A. Metabolic adaptations during extreme anoxia in the turtle heart and their implications for ischemia-reperfusion injury. *Sci Rep* . 2019;9(1):2850. doi:10.1038/s41598-019-39836-5
29. Pilkis SJ, Granner DK. Molecular Physiology of the Regulation of Hepatic Gluconeogenesis and Glycolysis. *Annu Rev Physiol* . 1992;54(1):885-909. doi:10.1146/annurev.ph.54.030192.004321
30. Storey KB. Metabolic regulation in mammalian hibernation: Enzyme and protein adaptations. *Comp Biochem Physiol Part A Physiol* . 1997;118(4):1115-1124. doi:10.1016/S0300-9629(97)00238-7
31. Ferguson RA, Storey KB. Gluconeogenesis in trout (*Oncorhynchus mykiss* ) white muscle: purification and characterization of fructose-1,6-bisphosphatase activity in vitro. *Fish Physiol Biochem* . 1992;10(3):201-212. doi:10.1007/BF00004514
32. Storey KB. Regulation of liver metabolism by enzyme phosphorylation during mammalian hibernation. *J Biol Chem* . 1987;262(4):1670-1673. doi:10.1016/S0021-9258(19)75689-0
33. Storey KB. Investigations of the mechanisms of glycolytic control during hibernation. *Can J Zool* . 1987;65(12):3079-3083. doi:10.1139/z87-467
34. Rui L. Energy Metabolism in the Liver. In: *Comprehensive Physiology* . Wiley; 2014:177-197. doi:10.1002/cphy.c130024

35. Cowan KJ, Storey KB. Freeze-Thaw Effects on Metabolic Enzymes in Wood Frog Organs. *Cryobiology* . 2001;43(1):32-45. doi:10.1006/cryo.2001.2338
36. Williamson JR, Wright PH, Malaisse WJ, Ashmore J. Control of gluconeogenesis by acetyl CoA in rats treated with glucagon and anti-insulin serum. *Biochem Biophys Res Commun* . 1966;24(5):765-770. doi:10.1016/0006-291X(66)90391-3
37. Pilkis SJ, El-Maghrabi MR, McGrane MM, Pilkis J, Claus TH. The role of fructose 2,6-bisphosphate in regulation of fructose-1,6-bisphosphatase. *J Biol Chem* . 1981;256(22):11489-11495. doi:10.1016/S0021-9258(19)68427-9
38. Hers HG, Hue L. Gluconeogenesis and Related Aspects of Glycolysis. *Annu Rev Biochem* . 1983;52(1):617-653. doi:10.1146/annurev.bi.52.070183.003153
39. Sistare FD, Haynes RC. The interaction between the cytosolic pyridine nucleotide redox potential and gluconeogenesis from lactate/pyruvate in isolated rat hepatocytes. Implications for investigations of hormone action. *J Biol Chem* . 1985;260(23):12748-12753. doi:10.1016/S0021-9258(17)38940-8
40. Haynes RC, Picking RA. The role of inhibition of pyruvate kinase in the stimulation of gluconeogenesis by glucagon: A reevaluation. *Arch Biochem Biophys* . 1990;283(1):51-59. doi:10.1016/0003-9861(90)90611-2
41. Engstrom L. The Regulation of Liver Pyruvate Kinase by Phosphorylation—Dephosphorylation. In: *Current Topics in Cellular Regulation* . ; 1978:29-51. doi:10.1016/B978-0-12-152813-3.50006-9
42. Feliu JE, Hue L, Hers HG. Hormonal control of pyruvate kinase activity and of gluconeogenesis in isolated hepatocytes. *Proc Natl Acad Sci* . 1976;73(8):2762-2766. doi:10.1073/pnas.73.8.2762
43. Storey KB, Storey JM. Tribute to P. L. Lutz: putting life on 'pause' - molecular regulation of hypometabolism. *J Exp Biol* . 2007;210(10):1700-1714. doi:10.1242/jeb.02716
44. Whitwam RE, Storey KB. Pyruvate Kinase from the Land Snail *Otala Lactea* : Regulation by Reversible Phosphorylation During Estivation and Anoxia. *J Exp Biol* . 1990;154(1):321-337. doi:10.1242/jeb.154.1.321
45. Mattice AMS, MacLean IA, Childers CL, Storey KB. Characterization of pyruvate kinase from the anoxia tolerant turtle, *Trachemys scripta elegans* : a potential role for enzyme methylation during metabolic rate depression. *PeerJ* . 2018;6:e4918. doi:10.7717/peerj.4918
46. Plaxton WC, Storey KB. Purification and properties of aerobic and anoxic forms of pyruvate kinase from the hepatopancreas of the channelled whelk, *Busycotypus canaliculatum* . *Arch Biochem Biophys* . 1985;243(1):195-205. doi:10.1016/0003-9861(85)90788-X
47. Storey KB. Regulation of hypometabolism: insights into epigenetic controls. *J Exp Biol* . 2015;218(1):150-159. doi:10.1242/jeb.106369
48. Gomes A V., Zong C, Ping P. Protein Degradation by the 26S Proteasome System in the Normal and Stressed Myocardium. *Antioxid Redox Signal* . 2006;8(9-10):1677-1691. doi:10.1089/ars.2006.8.1677
49. Alva N, Panisello-Rosello A, Flores M, Rosello-Catafau J, Carbonell T. Ubiquitin-proteasome system and oxidative stress in liver transplantation. *World J Gastroenterol* . 2018;24(31):3521-3530. doi:10.3748/wjg.v24.i31.3521
50. Enserink JM. Sumo and the cellular stress response. *Cell Div* . 2015;10(1):4. doi:10.1186/s13008-015-0010-1
51. Dawson NJ, Bell RA V., Storey KB. Purification and Properties of White Muscle Lactate Dehydrogenase from the Anoxia-Tolerant Turtle, the Red-Eared Slider, *Trachemys scripta elegans* . *Enzyme Res* . 2013;2013:1-8. doi:10.1155/2013/784973
52. Geiss-Friedlander R, Melchior F. Concepts in sumoylation: a decade on. *Nat Rev Mol Cell Biol* . 2007;8(12):947-956. doi:10.1038/nrm2293

53. Hochachka PW, Buck LT, Dolft CJ, Landt SC. Unifying theory of hypoxia tolerance: Molecular/metabolic defense and rescue mechanisms for surviving oxygen lack (oxygen sensing/hypoxia defense/turtle hepatocytes/turtle brain). *Biochemistry* . 1996;93:9493-9498.
54. Krivoruchko A, Storey KB. Forever Young: Mechanisms of Natural Anoxia Tolerance and Potential Links to Longevity. *Oxid Med Cell Longev* . 2010;3(3):186-198. doi:10.4161/oxim.3.3.12356
55. Biggar KK, Storey KB. The emerging roles of microRNAs in the molecular responses of metabolic rate depression. *J Mol Cell Biol* . 2011;3(3):167-175. doi:10.1093/jmcb/mjq045
56. Childers CL, Storey KB. Post-translational Regulation of Hexokinase Function and Protein Stability in the Aestivating Frog *Xenopus laevis* . *Protein J* . 2016;35(1):61-71. doi:10.1007/s10930-016-9647-0
57. Green SR, Storey KB. Regulation of the  $\alpha$ -ketoglutarate dehydrogenase complex during hibernation in a small mammal, the Richardson's ground squirrel (*Urocitellus richardsonii* ). *Biochim Biophys Acta - Proteins Proteomics* . 2020;1868(9):140448. doi:10.1016/j.bbapap.2020.140448
58. Smolinski MB, Green SR, Storey KB. Glucose-6-phosphate dehydrogenase is posttranslationally regulated in the larvae of the freeze-tolerant gall fly, *Eurosta solidaginis* , in response to freezing. *Arch Insect Biochem Physiol* . 2019;102(4). doi:10.1002/arch.21618

### Table legends

**Table 1** . Purification of FBPase from liver from anoxic *T.s. elegans* . Two steps were used: (a) Phosphocellulose column chromatography with elution using a 0–1 M KCl gradient with 2.5 mM FBP added, and (b) CM column chromatography with elution using a 0–1 M KCl gradient.

**Table 2** . Kinetic parameters for the forward reaction of purified *T.s. elegans* liver FBPase, comparing control versus anoxia conditions.  $V_{max}$  and  $K_m$  values were determined using saturating FBP concentrations.  $K_a$  and  $I_{50}$  values were determined at subsaturating FBP concentration (0.24  $\mu$ M). Data are means  $\pm$  SEM,  $n = 4$ . Assays were performed at 22  $^{\circ}$ C. ‘\*’ indicates that values for anoxic FBPase are significantly different from the control ( $n=3-4$ ) as determined by Student’s t-test  $p < 0.05$ .

### Figure legends

**Fig. 1. An overview of the allosteric regulation of FBPase on gluconeogenesis.** The schematic diagram shows the regulatory mechanism of FBPase during conversion of FBP to Fructose 6-phosphate during gluconeogenesis. The diagram shows the activators and inhibitors for the enzyme. The Blue colour arrows represent gluconeogenesis and red colour arrows represent glycolysis. The activators are represented as ‘+’ sign in green circle and inhibitors are represented as ‘-’ sign in red circle. **PFK** : phosphofructokinase, **FBPase** : Fructose-1,6 bisphosphate, **FBP** : Fructose 1,6 bisphosphate, **PEP** : phosphoenolpyruvate. Credit “Created with BioRender.com.”

**Fig. 2: 10% SDS-PAGE gel showing liver FBPase purification** . Lanes are: (1) Molecular weight ladder (Froggabio); (2) crude homogenate; (3) pooled peak fractions after elution from phosphocellulose; (4) pooled peak fractions after elution from CM column.

**Fig. 3 . Western blot analysis to evaluate posttranslational modifications in *T.s. elegans* liver FBPase from control versus anoxia conditions.** Relative band intensity is defined as the quotient of the intensity of the chemiluminescent signal divided by the band intensity observed after staining with Coomassie brilliant blue. ‘\*’ indicates that values for anoxic FBPase are significantly different from the control ( $n=3-4$ ) as determined by Student’s t-test  $p < 0.05$ .

**Fig. 4: *T.s. elegans* FBPase 3D homology structure and structure assessments** . (A ) Predicted homology model of *T.s. elegans* FBPase generated using SWISS-MODEL with the *T.s. elegans* FBPase amino acid sequence (NCBI) and X-ray crystallized human liver FBPase as the structural template. Each chain of the tetramer is in a different color. (B ) Visualisation of the FBPase structure with FBP and Mg was done using PyMOL Molecular Graphics System, Version 2.0 Schrödinger, LLC. (C ) Ramachandran plot of the

amino acid torsional angles for *T.s. elegans* FBPase 3D homology model, shows favorable  $\beta$ -sheet and  $\alpha$ -helix angles. (D) QMEAN structure scores as determined by SWISS-MODEL. All tests show numbers that are close to zero in the white area of the bar. Values close to or higher than zero indicate model scores that are expected or better than experimental structures of similar size.

**Table 1:**


Purification Step	Total Protein (mg)	Total Activity (mU)	Specific Activity (mU/mg)	Fold Purification	% Yield
Crude	21.07	94.25	4.47	---	---
PC	2.87	48.75	16.97	3.80	51.72
CM	0.59	23.4	39.13	8.75	24.82

**Table 2:**


**Hosted file**

image2.emf available at <https://authorea.com/users/734884/articles/711590-fructose-1-6-bisphosphatase-and-gluconeogenesis-control-in-an-anoxia-tolerant-turtle-trachemys-scripta-elegans>

**Figure 1:**



**Figure 2:**



**Figure 3**




Figure 4

