

Ana o 3 sIgE and diagnostic algorithms reduce overall cost of diagnosis of cashew allergy in children compared to skin prick test alone: a cost comparison analysis

Tim Brettig¹, Kim Dalziel¹, Jennifer Koplin¹, Thanh Dang¹, Lars Lange², Vicki McWilliam¹, Sakura Sato³, Savvas Savvastianos⁴, and Kirsten Perrett P¹

¹Murdoch Children's Research Institute

²GFO Kliniken Bonn Betriebsstatte St Marien

³Kokuritsu Byoin Kiko Sagamihara Byoin

⁴Allergy Department 2nd Pediatric Clinic University of Athens Greece

March 07, 2024

Abstract

Background: In the absence of a clear clinical history of reaction, diagnosis of cashew allergy using skin prick tests (SPT) or cashew-specific IgE requires a high number of oral food challenges (OFC). We recently showed that Ana o 3 sIgE alone, or a two-step diagnostic algorithm using cashew sIgE followed by Ana o 3 sIgE can reduce need for OFC. We aimed to determine if either of these approaches can provide a cost reduction to the health system compared to cashew SPT alone. **Methods:** Pooled individual level data from 6 studies was used to determine diagnostic accuracy and OFC rate. Two studies used cashew SPT (n=567, 198 allergic), with 95% positive and negative predictive values of [?]12mm and <3mm. Four studies were included in the pathways for Ana o 3 sIgE alone or a 2-step algorithm incorporating cashew and Ana o 3 sIgE (n=271, 156 allergic). Cut-offs used were [?]8.5kUA/L and [?]0.1kUA/L for cashew sIgE and [?]0.35kUA/L and [?]0.1kUA/L for Ana o 3 sIgE. Costs were constructed based on unit prices from hospital inpatient admissions, expenses incurred by families, individual patient data on allergic reaction types and rates and adrenaline autoinjector carriage, applying a health system perspective. **Results:** Modelled data through the Ana o 3 pathway resulted in a 46.43% cost reduction (\euro307,406/1000 patients) compared to using cashew SPT alone (\euro573,854/1000 patients). The 2-step algorithm resulted in a 44.94% cost reduction compared to SPT alone (\euro315,952.82/1000 patients). Both the Ana o 3 pathway and 2-step algorithm resulted in a 79-80% reduction in OFCs compared to SPT. **Conclusions:** Using Ana o 3 as a standalone test for cashew allergy diagnosis or a 2-step algorithm incorporating cashew sIgE and Ana o 3 sIgE is accurate and results in a large reduction in both OFCs and health system costs compared to cashew SPT alone.

Hosted file

Economic impact of ana o 3 v1.docx available at <https://authorea.com/users/488711/articles/713568-ana-o-3-sige-and-diagnostic-algorithms-reduce-overall-cost-of-diagnosis-of-cashew-allergy-in-children-compared-to-skin-prick-test-alone-a-cost-comparison-analysis>

Hosted file

Figures v1.docx available at <https://authorea.com/users/488711/articles/713568-ana-o-3-sige-and-diagnostic-algorithms-reduce-overall-cost-of-diagnosis-of-cashew-allergy-in-children-compared-to-skin-prick-test-alone-a-cost-comparison-analysis>

Hosted file

Tables v3.docx available at <https://authorea.com/users/488711/articles/713568-ana-o-3-sige-and-diagnostic-algorithms-reduce-overall-cost-of-diagnosis-of-cashew-allergy-in-children-compared-to-skin-prick-test-alone-a-cost-comparison-analysis>