

Sunil Gurrapu¹, Ashwini Ranada¹, and J P Patra¹

¹National Institute of Hydrology

December 26, 2022

Abstract

Indian summer monsoon rainfall is strongly influenced by various large-scale atmosphere-ocean oscillations including Pacific Decadal Oscillation (PDO), and El Niño-Southern Oscillation (ENSO). Researchers have shown that the negative phase of PDO or La Niña episodes of ENSO produce higher magnitude rainfall and hence relatively wetter years. So, it is imperative to have better knowledge of flood characteristics in the Indian watersheds for optimal planning and design of various infrastructure, and for optimal planning and management of reservoir operations. Traditionally, such information is estimated using flood frequency analysis (FFA) for small and medium sized projects. However, the adequacy of traditionally accepted assumption that the annual peak flows are independent and identically distributed (*i.i.d.*) is questioned globally (Gurrapu et al., 2016; Milly et al., 2008). This study evaluates the influence of PDO and ENSO and flood characteristics in Godavari River, India. The results indicate that the flood magnitude and frequency at majority of the selected gauges in Godavari River Basin is significantly influenced by PDO and ENSO, higher magnitude floods are associated with negative phase of PDO and La Niña episodes of ENSO. A few gauges are inversely related to these teleconnections. The influence of these teleconnections on regional climate is spatially variable across India and so are the contrasting results. The results from this study benefits the design engineers for efficient design of water resources infrastructures and water managers for optimal planning and management of reservoir operations.

References

- The Influence of the Pacific Decadal Oscillation on Annual Floods in the Rivers of Western Canada. (2016). *JAWRA Journal of the American Water Resources Association*, 52(5), 1031–1045. <https://doi.org/10.1111/1752-1688.12433>
- Stationarity Is Dead: Whither Water Management?. (2008). *Science*, 319(5863), 573–574. <https://doi.org/10.1126/science.1151915>

The Effects of Pacific Decadal Oscillation and El Niño-Southern Oscillation on Annual Floods in Godavari River Basin, India

Sunil Gurrapu¹, *, Ashwini Ranada¹, J P Patra¹


¹ National Institute of Hydrology, Roorkee, Uttarakhand – 247 667, India

* gurrapus@uregina.ca


INTRODUCTION

Indian summer monsoon rainfall is strongly influenced by large-scale atmosphere-ocean oscillations including Pacific Decadal Oscillation (PDO), El Niño-Southern Oscillation (ENSO). Researchers have shown that the negative phase of PDO or La Niña episodes of ENSO produce higher magnitude rainfall and hence relatively wetter years. This study is motivated by the observation that such teleconnections are not yet a key ingredient in planning and design of regional water resources and/or transportation infrastructure. This study is first of its kind to evaluate the impact of these low frequency oscillations, which are known to substantially control the magnitude and frequency of ISMR, on the annual peak flows in Indian watersheds.


(a) Godavari River Basin




Locations of the selected streamflow Gauging station spread across Godavari River Basin




Streamflow hydrographs at a gauging station in Godavari River basin in response to the extreme PDO events, i.e. 1987 (Positive) and 2008 (Negative).




Variability of the Pacific Decadal Oscillation (PDO) averaged over August to October (1901 – 2010), together with a 5-year running mean (red line).




Variability of the El Niño-Southern Oscillation (ENSO) as defined by the June Monthly Southern Oscillation Index (SOI)




Log-Normal expected quantiles and their 90% confidence interval for the annual peak flows in selected gauging stations of Godavari River basin stratified according to cool and warm phases of PDO.



Quantile-Quantile (Q-Q) plots based on the annual peak flows stratified according to the PDO at 46 streamflow gauging stations located across Godavari River Basin. The length of data at gauging station is shown in the upper left corners.



Histogram of flood ratios, flood quantiles in La Niña episodes to those in El Niño episodes, for several return periods



Histogram of flood ratios, flood quantiles in negative PDO phase to those in positive PDO phase, for several return periods

CONCLUSIONS

The results indicate that the annual peak flows in the gauges spread across Godavari River basin indicate that the magnitude and frequency of annual peak flows is substantially influenced by the phase of PDO, ENSO, or IOD. In majority of the gauges, higher magnitude floods seem to be more common during the negative phase of PDO or during the La Niña episodes of ENSO pattern. Overall, the results from this study highlight the potential inadequacy of *i.i.d.* assumption and is not tenable where the hydroclimatology is strongly influenced by the low-frequency atmosphere-ocean oscillations. These results are in agreement with the observations made by other researchers across the globe. This is a manifest in Godavari and Narmada River basins in India and other parts across the globe including western Canada, California and eastern Australia. The extent of this problem in other Indian watersheds remains to be explored. Any region with a strong teleconnection with such large-scale atmosphere-ocean oscillations may be subject to under- or over-estimation of the design flood. Therefore, the knowledge of the regional hydroclimate with regards to phases of the large-scale atmosphere-ocean oscillations should be considered prior to estimating design flood. Furthermore, the effect of other atmosphere-ocean oscillations, e.g. North Atlantic Oscillation (NAO), Arctic Oscillation (AO) and Atlantic Multi-Decadal Oscillation (AMO), on extreme hydrology of Indian watersheds needs to be explored.