

# Supporting Information for “Seasonal Forecasts of the Exceptional Northern Hemisphere Winter of 2020”

Simon H. Lee<sup>1</sup>, Zachary D. Lawrence<sup>2,3</sup>, Amy H. Butler<sup>2,4</sup>, and Alexey Y.

Karpechko<sup>5</sup>

<sup>1</sup>Department of Meteorology, University of Reading, Reading, UK

<sup>2</sup>Cooperative Institute for Research in Environmental Sciences (CIRES), University of Colorado, Boulder, Colorado, USA

<sup>3</sup>NOAA Physical Sciences Laboratory (PSL), Boulder, Colorado, USA

<sup>4</sup>NOAA Chemical Sciences Laboratory (CSL), Boulder, Colorado, USA


<sup>5</sup>Finnish Meteorological Institute, Helsinki, Finland

## Contents of this file


1. Table S1
2. Figures S1 to S2

**Table S1.** Details of the 6 seasonal forecast models used in this study. In a “burst” ensemble the entire ensemble is produced at the same time, while “lagged” ensembles are produced as a collection of smaller ensembles over several days. The number in parentheses indicates the hindcast ensemble size. Accessed from <https://confluence.ecmwf.int/display/CKB/Description+of+the+C3S+seasonal+multi-system>.

Center	Model	Ens. Type	Ens. Size (Hindcasts)	Resolution
CMCC	CMCC-SPS3	Burst	50 (40)	1° lat-lon L46
DWD	GCFS 2.0	Burst	50 (30)	T127 L95
ECMWF	SEAS5	Burst	51 (25)	Tco319 L91
Météo-France	System 7	Lagged	51 (25)	TL359 L91
NCEP	CFSv2	Lagged	120 (24)	T128 L64
UKMO	GloSea5-GC2	Lagged	60 (28)	N216 L85



**Figure S1.** As Figure 1 (a–h) in the main text, but for forecasts nominally initialized on 1 October 2019.



**Figure S2.** As Figure S1 but for forecasts nominally initialized on 1 November 2019.