

Table 3: ECMO Characteristics

Characteristics	Overall	Survivor	Non-Survivor	p
Cannulated < 24 hours from admission (%)	23 (48.9)	8 (66.7)	15 (42.9)	0.193
Peripheral Cannulation (%)	44 (93.6)	12 (100.0)	32 (91.4)	0.560
Targeted Temperature Management (%)	22 (46.8)	6 (50.0)	16 (45.7)	1.000
Mechanical circulatory support prior to ECMO (%)	14 (29.8)	6 (50.0)	8 (22.9)	0.140
Revascularization pursued (%)	19 (40.4)	6 (50.0)	13 (37.1)	0.506
Venting ventricular assist device (%)	6 (12.8)	2 (16.7)	4 (11.4)	0.637
Time on ECMO in days (median, [IQR])	3.00 [1.64, 6.42]	3.85 [2.93, 7.75]	2.73 [1.31, 5.06]	0.110
ICU length of stay in days (median, [IQR])	10.00 [2.00, 21.00]	28.50 [13.50, 39.50]	7.00 [1.50, 15.50]	0.001*
Hospital length of stay in days (median, [IQR])	16.00 [4.50, 27.00]	35.50 [21.75, 49.50]	8.00 [3.00, 17.50]	<0.001*