

TABLE 1: Prognostic variables on univariate analysis

| Prognostic variable | DFS probability | <i>p</i> value | OS probability | <i>p</i> value | hazard ratio | <i>p</i> value |
|-------------------------------|-----------------|----------------|----------------|----------------|--------------|----------------|
| Age | | | | | | |
| < 2 years | 19.8 | 0.96 | 23.3 | 0.92 | 1.02 | 0.927 |
| > 2 years | 15.3 | | 15.2 | | | |
| Primary site | | | | | | |
| Adrenal | 14.7 | 0.178 | 17.6 | 0.07 | 1.00 | 0.99 |
| Cervical | 30.0 | | 30.0 | | | |
| Post. mediastinal | 50.0 | | 100.0 | | | |
| Paraspinal | 13.3 | | 13.3 | | | |
| Abdominal/ retroperitoneal | 33.3 | | 33.3 | | | |
| Multifocal | 25.0 | | 25.0 | | | |
| Metastatic site | | | | | | |
| Bone marrow | 18.0 | 0.865 | 20.7 | 0.87 | 1.3 | 0.861 |
| Bones | 9.1 | | 9.1 | | | |
| Lymph nodes | 25.0 | | 25.0 | | | |
| Multiple sites | 16.3 | | 15.9 | | | |
| LDH | | | | | | |
| < 750 U/L | 50.0 | 0.028 | 46.9 | 0.032 | 2.29 | 0.013 |

| | | | | | | |
|------------------------|------|-------|------|-------|------|-------|
| > 750 U/L | 18.6 | | 16.2 | | | |
| Chemotherapy | | | | | | |
| Chemo A+ Metronomic | 55.6 | | 55.6 | | | |
| Chemo A | 23.6 | 0.001 | 27.2 | 0.001 | 6.03 | 0.001 |
| Chemo B | 8.3 | | 8.1 | | | |
| Response | | | | | | |
| Good | 27.8 | 0.001 | 29.7 | 0.001 | 6.96 | 0.001 |
| Poor | 3.3 | | 3.3 | | | |
| Surgery | | | | | | |
| No surgery | 5.7 | 0.001 | 7.5 | 0.001 | 1.32 | 0.001 |
| Biopsy only | 0.0 | | 0.0 | | | |
| Debulking | 66.7 | | 66.7 | | | |
| Excision | 43.8 | | 43.8 | | | |