

Table 2. Characteristics after ablation

	Group 1 n=11	Group 2 n=21	p value
echocardiographic parameters – median (IQR)			
follow-up period of echocardiography	199(165-223)	276(115-402)	0.968
LAD - mm	41.3(37.5-43.7)	37.6(33.2-43.2)	0.275
LAV - ml	58.0(55.5-66.5)	45.9(37.8-51.8)	0.0099
LAVI - ml/m ²	32.9(29.0-42.2)	23.9(20.8-27.5)	0.0073
LVEF - %	48.8(38.0-54.4)	64.1(61.8-66.9)	0.0000323
LV end-diastolic diameter - mm	53.5(49.8-57.5)	47.4(42.9-50.2)	0.0012
LV end-systolic diameter - mm	37.9(37.1-47.5)	30.9(28.1-33.7)	0.000304
NT-proBNP - median (IQR) pg/ml	337(72-1242)	99(51-134)	0.275
Recurrence of Arrhythmia - no.(%)	2(18.2)	7(33.3)	0.441
Period from ablation to recurrence - median (IQR)	635	176(105-397)	0.667

Abbreviations: IQR, interquartile range; LAD, left atrial diameter; LAV, left atrial volume; LAVI, left atrial volume index; LVEF, left ventricular ejection fraction; LV, left ventricular