Table 7. Absorbance at 670 nm (Chlorophyll) noted in the 'Arbequina'. 'Picual' and 'Verdial' olive oils extracted from fruit, picked with a Manual Inverted Umbrella (R1) and in a traditional way (R2) and stored during 0, 4, 8, and 14 days at 5 °C (C1) and ambient temperature (C2)[a].

	CHLOROPHYLL					
ST(days); R (1.2); C (1.2)	ARBEQUINA		PICUAL		VERDIAL	
	year 1	year 2	year 1	year 2	year 1	year 2
0; 1; 1	9.67 ± 0.93 **B** b x	12.48 ± 1.70 **B** x	4.31 ± 0.70 **B** a x	13.08 ± 1.73 b x	2.17 ± 0.10 **C**	23.35 ± 3.10 **AB**
0; 1; 2	9.67 ± 0.93 **A** b x	12.48 ± 1.70 **B** x	4.31 ± 0.70 **B** a x	13.08 ± 1.73 **A** b x	2.17 ± 0.10 **B**	23.35 ± 3.10 **A**
0; 2; 1	12.64 ± 0.90 **AB** a y	15.16 ± 0.51 y	2.64 ± 0.30 **B** b y	18.53 ± 0.70 **A** a y	2.63 ± 0.50 **B**	24.23 ± 2.28
0; 2; 2	12.64 ± 0.90 **A** a y	15.16 ± 0.51 **A** y	2.64 ± 0.30 **B** b y	18.53 ± 0.70 **A** a y	2.63 ± 0.50 **B**	24.23 ± 2.28 **A**
4; 1; 1	12.80 ± 1.75 **AB** ab α	11.70 ± 0.14 **B** x α	3.54 ± 0.08 **B** bc x α	13.28 ± 2.10 b x α	4.49 ± 0.64 **A** ab x α	18.51 ± 1.97 **B**
4; 1; 2	8.60 ± 1.08 **AB** c β	10.11 ± 0.69 **B** x β	4.11 ± 0.26 **B** a x β	10.80 ± 1.02 **AB** b x β	3.77 ± 0.23 **A** b x β	18.90 ± 1.77 **AB**
4; 2; 1	14.24 ± 0.88 **A** a α	16.84 ± 2.51 y α	2.96 ± 0.29 **B** c y α	19.32 ± 2.77 **A** a y α	5.71 ± 0.29 **A** a y α	22.32 ± 2.26
4; 2; 2	10.47 ± 1.32 **AB** bc β	9.85 ± 0.05 **B** y β	3.28 ± 0.22 **B** c y β	14.26 ± 0.96 **B** b y β	4.36 ± 0.69 **A** b y β	17.65 ± 1.46 **B**
8; 1; 1	13.87 ± 1.12 **A** a x α	17.87 ± 1.58 **A** a α	3.90 ± 0.21 **B** a x	13.12 ± 2.97 b x α	4.34 ± 0.21 **A** b x α	24.85 ± 1.35 **A** a α
8; 1; 2	7.80 ± 0.40 **AB** b x β	9.69 ± 0.71 **B** b β	3.77 ± 0.28 **B** a x	9.46 ± 0.86 **B** b x β	4.19 ± 0.30 **A** b x β	16.37 ± 0.57 **B** b β
8; 2; 1	14.35 ± 1.56 **A** a y α	17.76 ± 2.23 a α	2.06 ± 0.19 **A** c y	18.92 ± 0.68 **A** a y α	5.26 ± 0.20 **A** a y α	27.54 ± 1.49 a α
8; 2; 2	11.92 ± 1.39 **AB** a y β	10.60 ± 0.27 **B** b β	2.82 ± 0.13 **A** b y	11.09 ± 1.03 **C** b y β	4.13 ± 0.39 **A** b y β	14.91 ± 0.18 **B** b β
14; 1; 1	11.31 ± 1.00 **AB** a α	20.22 ± 0.97 **A** a x α	3.00 ± 0.21 **A**	12.48 ± 1.71 a α	3.26 ± 0.28 **B** c x	27.92 ± 2.03 **A** a α
14; 1; 2	6.00 ± 1.64 **B** b β	15.85 ± 1.22 **A** b x β	3.40 ± 0.22 **A**	8.02 ± 0.37 **B** b β	4.36 ± 0.23 **A** b x	16.77 ± 1.24 **B** b β
14; 2; 1	10.57 ± 1.17 **B** a α	17.30 ± 1.38 ab y α	2.60 ± 0.74 **A**	13.57 ± 1.59 **B** a α	5.54 ± 0.30 **A** a y	24.75 ± 2.28 a α
14; 2; 2	8.96 ± 1.39 **B** ab β	14.25 ± 1.09 **A** b y β	3.40 ± 0.53 **A**	8.68 ± 0.82 **D** b β	4.59 ± 0.55 **A** ab y	18.76 ± 1.34 **B** b β
Storage Time (ST)	.000	.000	.035	.000	.000	.000
Treatment (T)	.000	.000	.000	.000	.000	.000
ST × T	.002	.000	.003	.001	.001	.000
Harvesting (R)	.000	.036	.000	.000	.000	.343
Conservation (C)	.000	.000	.005	.000	.001	.000
ST × R	.274	.000	.000	.005	.039	.677
ST × C	.000	.000	.302	.000	.003	.000
R × C	.008	.314	.268	.053	.000	.376
ST × R × C	.119	.010	.321	.332	.022	.012

[a] In each variable the values of different treatments followed by different letters are significantly different according to the Tukey test ($P < 0.05$). Absence of letters means no significant effect due to treatment according to one-way ANOVA ($P < 0.05$). In each column, values at different storage times (ST) and the same harvesting method (R) and conservation method (C), followed by different upper bold case letters are significantly different; four values at each ST, followed by different lower case letters (a, b, c, d) are different; two values at the

same ST and same conservation method (C), but different harvesting method (R), followed by lower case letters (x or y), are different; two values at the same ST and same R, but different C, followed by different Greek letters are significantly different. Each value is the mean ± SD of 3 replicates.