

Surgical step	Our technique	One-step conservative surgery.^{7,8,9}	Triple-P technique.²⁸	Stepwise surgical approach.¹⁰
<i>Blood loss :</i>				
<i>Median (range)</i>				
<i>Intraoperative</i>	1400 (500-2000)			1700 (600-2400)
<i>Postoperative</i>	100 (80-350)			570 (400-1300)
<i>Mean +/- SD</i>				
<i>-Intraoperative</i>	1.3 +/- .36		1.70 +/- 0.95	
<i>-total</i>	1.4 +/- .36		2.31 +/- 1.38	
<i>>1500 CC (%)</i>	3/20 (15%)	94/326 (28.8%)		
<i>Blood transfusion:</i>				
<i>Median (range)</i>	1.5 (0-3)			4 (2-6)
<i>-Number of patients requiring transfusion</i>	12/ 20 (60%)		9/19 (47.4%)	
<i>-Total RBCs units transfused</i>	24		40	
<i>Hb reduction</i>	1.1 +/- 0.59		2.16 +/- 1.43	
<i>PPH</i>	0/20 (0%)		3/19 (15.8%)	
<i>CsH required/ failed conservation</i>	2/20 (10%)	96/326 (29.45%)	0 (0%)	6/71 (8.5%)
<i>UB injury</i>	1/20 (5%)	47/326 (14.4%)	0 (0%)	9/71 (12.7%)
<i>Operative time</i>				
<i>Median (range)</i>	120 (75-210)			85 (70-140)
<i>Hospital stay</i>	1.3 +/- 0.73		4.75 +/- 2	
<i>Vesico-uterine fistula</i>	0 (0%)		0 (0%)	1/71 (1.4%)
<i>Paralytic ileus</i>	1/20(5%)	13/326 (3.99%)		

<i>Arterial thrombosis</i>	0 (0%)	1/326 (0.3%)	1/19 (5.2%)	0 (0%)
----------------------------	--------	--------------	-------------	--------