

<b>Demographic data (unit)</b>		<b>Median (IQR)</b>
Age (years)		31 (6.5)
Parity (number)		2 (1)
Previous caesarian sections (number)		2 (1)
Gestational age at termination (weeks)		36 (0)
<b>Surgical outcomes (unit)</b>		<b>Median (IQR)</b>
Intraoperative blood loss (cubic centimeters (CC))		1400 (400)
Postoperative blood loss (CC)		100 (50)
Total blood loss (CC)		1500 (387.5)
Total blood transfusion (packed red blood cell units)		1.5 (2)
Hemoglobin drop (gram/ deciliter)		1.1 (0.65)
Operative time (minutes)		120 (37.5)
Hospital stay (days)		1 (0)
APGAR score minute 1		9 (1)
APGAR score minute 5		10 (1)
<b>Surgical outcomes</b>		<b>Number (percentage)</b>
Cervical invasion		3/ 20 (15%)
Failed conservative surgery (CsH)		2/ 20 (10%)
Urinary bladder injury		1/ 20 (5%)
Postoperative paralytic ileus		1/ 20 (5%)
Peritoneal inclusion cyst (pseudocyst) due to intra-abdominal adhesions		1/ 20 (5%)
Symptomatic uterine niche (>1 CC)		1/ 20 (5%)
Wound infection		1/20 (5%)
Blood transfusion needed		12/20 (60%)
Intraoperative blood loss > 1500 CC		3/20 (15%)
Neonatal intensive care unit (NICU) admission		6/20 (30%)
Ureteric injury		0
Maternal intensive care unit admission		0
Deep venous thrombosis		0
Severe intrauterine adhesions		0
Postoperative urinary retention		0
Vesico-uterine/ vaginal fistula		0
Arterial thrombosis		0
Postpartum hemorrhage		0
Postpartum fever (sepsis)		0
Maternal mortality		0
Serious morbidity: Renal failure requiring dialysis Respiratory failure Hysterectomy		0
Placenta previa type by ultrasound	Complete centralis	6/20 (30%)
	Incomplete centralis	14/20 (70%)

Preoperative radiological diagnosis	Total percreta	5/20 (25%)
	Total increta, partial percreta	15/20 (75%)
Intraoperative clinical diagnosis	Total percreta	4/20 (20%)
	Total increta, partial percreta	14/20 (70%)
	Total accrete, partial increta	2/20 (10%)
Postoperative histopathological diagnosis	Placenta percreta	18/20 (90%)
	Placenta increta	2/20 (10%)
Past medical history	Free	14/20 (70%)
	Gestational Anemia	3/20 (15%)
	Gestational hypertension	1/20 (5%)
	Pregestational hypertension	1/20 (5%)
	Pregestational diabetes mellitus	1/20 (5%)
Termination time and setting	Elective (near 36 weeks)	18/20 (90%)
	Emergency (31 weeks)	1/20 (5%)
	Emergency (34 weeks)	1/20 (5%)
NICU admission	Re-admission for hyperbilirubinemia	3/20 (15%)
	Transient tachypnea of newborn	2/20 (10%)
	Prematurity	2/20 (10%)