

TABLE 2 Median dietary intake and glycemic indices at study entry, end of induction, and end of month-6					
	Study Entry (N=23) Median [Minimum, Maximum]	End of Induction (N=23) Median [Minimum, Maximum]	End of Month-6 (N=20) ^a Median [Minimum, Maximum]	Across all three timepoints	From study entry to end of month-6
				<i>P</i> -value	<i>P</i> -value
Calories	1360 [479, 2637]	2005 [868, 3939]	1688 [560, 2436]	0.013	0.250
Protein (% of kcal)	17 [9, 26]	19 [13, 36]	18 [12, 27]	0.073	0.027
Carbohydrates (% of kcal)	51 [39, 72]	46 [23, 58]	50 [36, 60]	0.035	0.064
Fat (% of kcal)	32 [15, 41]	36 [24, 44]	33 [25, 46]	0.063	0.090
Saturated fat (% of kcal)	11 [3, 16]	12 [7, 20]	10 [7, 18]	0.300	0.120
Sugar (g/1000 kcal)	55 [21, 94]	39 [13, 66]	48 [17, 78]	0.0051	0.003
Dietary Fiber (g/1000 kcal)	8 [4, 13]	9 [5, 19]	10 [4, 18]	0.210	0.083
Glycemic Index	57 [46, 66]	56 [48, 71]	58 [49, 64]	0.920	0.900
Glycemic Load (g/1000 kcal)	64 [46, 94]	61 [26, 85]	62 [50, 81]	0.030	0.053
^a Three patients had missing diet data at month-6.					