

Table 4: Association between demographic, clinical, laboratory and radiological parameters with mortality

	Alive (n=24)	Death (n=9)	Effect size (95% CI)
Mean (SD) age in months	9.6 (4.5)	9.1(3.9)	0.97 (0.83 to 1.15)
Male	20 (83.3)	6 (66.7)	0.54 (0.13 to 2.15)
Duration of illness in days, mean (SD)	10.8 (7.8)	20.2 (13.8)	0.02 (0.003 to 0.03)*
SpO2 <90%	12 (50)	7 (77.8)	2.58 (0.54 to 12.41)
Anemia (Hb<10 gm/dl)	13 (54.2)	5 (55.6)	1.04 (0.28 to 3.87)
TLC <15000	12 (50)	6 (66.7)	1.67 (0.42 to 6.66)
Platelet <1.5 lac	2 (8.33)	1 (11.1)	1.25 (0.15 to 9.99)
CRP >=25	6 (25)	5 (55.6)	2.5 (0.67 to 9.31)
SGPT>=100	6 (25)	5 (55.6)	2.5 (0.67 to 9.31)
MV required	16 (66.7)	9 (100)	---
Inotropes required	8 (33.3)	6 (66.7)	2.71 (0.68 to 10.85)
CXR pathcy opacities	7 (29.2)	8 (88.9)	9.6 (1.2 to 76.76)*

Figures are number (percentages) unless stated otherwise

*significant