

<b>Table 1. Baseline characteristics</b>			
<b>Variable</b>	<b>Thrombocytopenia (n=2,402, 10.4%)</b>	<b>Normal platelet count (n=20,803)</b>	<b>All patients (n=23,205)</b>
<b>Maternal age (±SD)</b>	32.8 (4.8)	32.1 (5.0)	32.2 (5.0)
<b>Pre-pregnancy BMI (IQR)</b>	23.0 (20.9-25.8)	23.5 (21.2-26.9)	23.4 (21.2-26.8)
<b>Gestational age at delivery (IQR)</b>	39.6 (37.7-40.8)	39.7 (38.1-40.7)	39.7 (38.0-40.7)
<b>Gravidity (±SD)</b>	2.3 (1.5)	2.3 (1.5)	2.3 (1.5)
<b>Nulliparous, n (%)</b>	1,137 (47.3)	9,726 (46.8)	10,863 (46.8)
<b>Multiple gestation, n (%)</b>	209 (8.7)	723 (3.5)	932 (4.0)
<b>Birth weight in grams (IQR)</b>	3,339 (2,783-3,756)	3,348 (2,866-3,720)	3,346 (2,857-3,725)
<b>Macrosomia, n (%)</b>	332 (13.8)	2,070 (10.0)	2,402 (10.4)
<b>Mode of delivery, n (%)</b>			
Normal vaginal	1,473 (61.3)	14,215 (68.3)	15,688 (67.6)
Instrumental	179 (7.5)	1,538 (7.4)	1,717 (7.4)
Primary caesarean	449 (18.7)	2,609 (12.5)	3,058 (13.2)
Emergency caesarean	301 (12.5)	2,441 (11.7)	2,742 (11.8)
<b>Induction of labour, n (%)</b>	608 (25.3)	4,832 (23.2)	5,440 (23.4)
<b>Pregnancy complications, n (%)</b>			
Manual placenta removal	141 (5.9)	1,121 (5.4)	1,262 (5.4)
Perineal or birth tract injuries	83 (3.5)	631 (3.0)	714 (3.1)
Preeclampsia	114 (4.7)	542 (2.6)	656 (2.8)
<b>Prior SPPH, n (%)</b>	102 (4.2)	895 (4.3)	997 (4.3)
<b>Prior caesarean delivery, n (%)</b>	275 (11.4)	2,319 (11.1)	2,594 (11.2)
<b>Platelet count (IQR)</b>	134 (116-143)	223 (186-274)	
<b>Platelet disorders, n (%)</b>			
Gestational thrombocytopenia or thrombocytopenia of unknown origin	2,206 (91.8)	-	2,206 (9.5)
HELLP syndrome	171 (7.1)*	84 (0.4)	255 (1.1)
ITP	20 (0.8)*	10 (<0.1)	30 (0.1)
Thrombopathy	4 (0.2)	16 (0.1)	20 (0.1)
Other	2 (0.1)	3 (<0.1)	5 (<0.1)
<b>Platelet transfusion between last platelet count before delivery, and 24 hours after delivery, n(%)</b>	35 (1.5)	28 (0.1)	63 (0.3)