

Table 1. Demographics and clinical data of the study patients.

Variable	Value
Age (year)	64,26±10.6
Gender (F/M), n (%)	28 (37.8%)/46 (62.2%)
SOFA score	11.2±3.07
Type of surgery n (%)	
CABG	32 (43.2%)
CABG+Valve replacement	12 (16.2%)
AVR+MVR	11 (14.9%)
Abdominal aortic aneurysm repair	2 (2.7%)
Aortic aneurism and dissection repair	11 (14.9%)
EVAR	1 (2.7%)
Aortabifemoral surgery	2(2.7%)
Heart Tx	1(1.4%)
Medical history n (%)	
Respiratory disease	29 (31.1%)
Diabetes mellitus	11 (29.7%)
Renal dysfunction	19 (25.7%)
Hyperlipidemia	18 (24.3%)
Hypertension	26 (35.1%)
Postoperative days until infection	18.5±11.6
Infection site, n (%)	
Respiratory	58 (78.4%)
Surgical site	11 (10.8%)
Bloodstream	3 (4.1%)
Urinary tract	2 (2.7%)
SICU stay after infection (days)	46.97±30.4
Days of mechanical ventilation after infection	35.04±24.3
Mortality, n (%)	57 (77%)
7 days	17(%23.1)
30 days	42 (%57.3)