

	<b>Normal</b>	<b>Obstructive</b>	<b>NSVL</b>	<b>Restrictive</b>	<b>p-value</b>
Total Number Subjects	188	27	19	19	
Age (years), mean (SD)	15.7 (3.3)	16.3 (1.6)	17.7 (4.3)	15.1 (1.1)	<i>0.041</i>
Body Mass Index, mean (SD)	18.9 (3)	19.9 (2.9)	19.1 (2.4)	18.9 (2.1)	0.40
Haller Index, median (Q1, Q3)	4.8 (4, 5.9)	4.5 (4, 5.3)	5.0 (4.3, 6.2)	4.8 (4, 5.4)	0.60
Correction Index, mean (SD)	34.4 (14.1)	31.2 (16.8)	35.0 (14.8)	31.2 (10.4)	0.59
Dyspnea with Exercise, % (n)	64.4% (121)	63.0% (17)	68.4% (13)	42.1% (8)	0.27
Chest Pain, % (n)	42.6% (80)	48.2% (13)	42.1% (8)	10.5% (2)	<i>0.045</i>
Peak VO <sub>2</sub> , mean (SD)	89.9 (16.5)	87.3 (16.1)	85.9 (15.9)	84.5 (18.7)	0.42
Peak O <sub>2</sub> Pulse, mean (SD)	92.0 (17.6)	89.4 (17.6)	86.3 (17.2)	85.7 (18.7)	0.28
Breathing Reserve, mean (SD)	.27 (12.8)	.33 (11.1)	.32 (13.4)	.26 (15.1)	0.08
Vt/IC, mean (SD)	.67 (0.11)	.66 (0.12)	.69 (0.16)	.69 (0.11)	0.73
Respiratory Rate, mean (SD)	52.5 (8.9)	50.1 (7.8)	49.8 (10.3)	52.7 (9.0)	0.35

**Table 2.** Symptoms, anatomic severity indices, and CPET outcome variables by normal and abnormal pulmonary function testing. Abbreviations: NSVL (nonspecific ventilatory limitation), VO<sub>2</sub> (oxygen consumption), O<sub>2</sub> pulse (oxygen pulse), Vt (tidal volume), IC (inspiratory capacity).