

**TABLE 1: summary of the details and data of the included subjects**

<b>Sociodemographic details</b>	
<b>Age ( years )</b>	35.1 (24-47 )
<b>Males</b>	44 (75.8 )
<b>Females</b>	14 (14.2 )
<b>Body weight ( kg )</b>	49.5 (42-60)
<b>BMI( kg/m2 )</b>	19.6 (17.1-24.1 )
<b>Co-infections</b>	
<b>HIV</b>	Negative
<b>HCV</b>	Negative
<b>HBV</b>	Negative
<b>Lab parameters</b>	
<b>ESR (mm/hr))</b>	52.5 ( 40-69 )
<b>Heamoglobin ( gm/ dl )</b>	11.3 ( 7.8-13.2 )
<b>ALT (sgpt) ( iu/l )</b>	60.0 (56-65)
<b>AST (sgot) ( iu/l )</b>	46.3 (41-56)
<b>Clinical factors</b>	
<b>Past history of Tuberculosis</b>	
<b>Yes</b>	52
<b>No</b>	06
<b>Co -medication</b>	
<b>Pyrazinamide</b>	58
<b>Cycloserine</b>	58
<b>Kanamycin</b>	58
<b>Linezolid</b>	58
<b>Ethionamide</b>	58
<b>Paracetamol</b>	05
Abbreviations: BMI- Body mass index, HIV-Human immunodeficiency virus infection, HCV- Hepatitis C virus, HBV- hepatitis B virus, ESR- Erythrocyte sedimentation rate, ALT-alanine transaminase, AST- aspartate aminotransferase.	