

Table1 : Social demographic factors were compared among non-snorers,occasional snore s and habitual snores

Social Demography	Non-snorers (n=402)	Occasional Snore s (n=683)	Habitual Snore s (n=463)	<b>P</b>
<b>Age</b>				<b>0.002</b>
< 7Y	159 (39.6)	310 (45.4)	240 (51.8)	
≥7Y	243 (60.4)	373 (54.6)	223 (26.6)	
<b>Sex</b>				<b>0.089</b>
Boy	207 (51.5)	427 (62.5)	292 (31.5)	
Girl	195 (48.5)	256 (37.5)	171 (27.5)	
<b>Race</b>				<b>0.326</b>
Han	382 (96.5)	649 (97.6)	450 (30.4)	
Other Race	14 (3.5)	16 (2.4)	9 (23.1)	
<b>Living Districts</b>				<b>0.141</b>
City	169 (42.0)	330 (48.3)	231 (31.6)	
Suburb	67 (16.7)	131 (19.2)	84 (29.8)	
County seat	67 (16.7)	150 (22.0)	97 (30.9)	
Town	51 (12.7)	38 (5.6)	30 (25.2)	
Countryside	48 (11.9)	34 (5.0)	21 (20.4)	

Table2 : Mothers with pregnancy complications were compared among non-snorers, occasional snore s and habitual snore s

Disease of Pregnancy	Non-snorers (n=402)	Occasional Snore s (n=683)	Habitual Snore s (n=463)	<b>P</b>
Yes	76 (18.9)	168 (24.6)	77 (16.6)	<b>0.006</b>
No	326 (81.1)	415 (75.4)	386 (83.4)	

Table3 : The upper respiratory disease within 6 months were compared among non-snorers, occasional snore s and habitual snore s

Upper respiratory disease	Non-snorers (n=402)	Occasional Snorers (n=683)	Habitual Snorers (n=463)	P
Yes	49 (12.4)	156 (26.7)	196 (43.8)	<0.001
No	347 (87.6)	429 (73.3)	267 (24.3)	
Tonsillitis	19 (4.7)	82 (12.0)	115 (53.2)	<0.001
Chronic Rhino-Sinusitis	30 (7.5)	73 (10.7)	62 (37.6)	0.023
Allergic Rhinitis	23 (5.7)	71 (10.4)	62 (39.7)	0.005
Pneumonia (bronchitis)	21 (5.2)	76 (14.3)	98 (43.2)	0.012
Asthma	4 (1.0)	11 (1.6)	5 (1.1)	0.629

Table4: Passive smoking exposure were compared among non-snorers, occasional snorers and habitual snorers

Passive Smoking Exposure	Habitual Snorers		P
	Yes	No	
Smoking During Pregnancy	25 (43.1)	438 (29.4)	<b>0.025</b>
Exposure to Secondhand Smoke During Pregnancy	76 (41.8)	387 (28.3)	<b>&lt;0.001</b>
Exposure to Secondhand Smoke	72 (42.9)	391 (28.3)	<b>&lt;0.001</b>