

eHealth tools for childhood cancer survivorship care: A qualitative analysis of survivors', parents', and general practitioners' views.

Maria Schaffer (BA (Hons)).^{1,2} Jordana K. McLoone (PhD).^{1,2} Claire E. Wakefield, (PhD, MPH)^{1,2}, Mary-Ellen Brierley (BPsych (Hons)).^{1,2} Afaf Girgis, A.,³ Maria C McCarthy(PhD).^{4,5} Elysia Thornton-Benko (PhD).⁶ Raymond J Chan.,⁷ Karen A Johnston (MN-NursPrac).^{1,2} Richard J Cohn (FRACP).^{1,2} and Christina Signorelli (PhD).^{1,2}

on behalf of the **ANZCHOG Survivorship Study Group***

1. Kids Cancer Centre, Sydney Children's Hospital, Randwick, NSW, Australia
2. School of Women's and Children's Health, UNSW Medicine, University of New South Wales, Sydney, NSW, Australia
3. Centre for Oncology Education and Research Translation, Ingham Institute for Applied Medical Research, South Western Sydney Clinical School, UNSW Sydney, NSW, Australia
4. Murdoch Children's Research Institute, Parkville, Victoria 3052, Australia
5. Children's Cancer Centre, Royal Children's Hospital, Parkville, Victoria 3052, Australia
6. Bondi Road Doctors, Bondi Junction, NSW, Australia
7. Cancer and Palliative Care Outcomes Centre and School of Nursing, Queensland University of Technology (QUT), Kelvin Grove, Queensland 4059, Australia
8. Division of Cancer Services, Princess Alexandra Hospital, Woolloongabba, Queensland 4102, Australia.

*Correspondence to:

Dr Christina Signorelli. Behavioural Sciences Unit, Kids Cancer Centre, Sydney Children's Hospital, High St, Randwick, NSW 2031 AUSTRALIA. Ph: +612 9382 3115; F: +612 9382 1789; E: c.signorelli@unsw.edu.au

Text Word Count: 3,000;

Abstract word count: 250;

Brief running title: eHealth tools for childhood cancer survivorship care

Key words: eHealth, Survivorship, Pediatrics

Tables: 4

Figures: 2

Abbreviations	
GP	General practitioners
SCP	Survivorship care plans
ALL	Acute Lymphoblastic Leukemia

*The members of the **ANZCHOG Survivorship Study Group** in alphabetical order: Dr Frank Alvaro, Prof Richard Cohn, Dr Rob Corbett, Dr Peter Downie, Ms Karen Egan, Ms Sarah Ellis, Prof Jon Emery, Dr Joanna Fardell, Ms Tali Foreman, Dr Melissa Gabriel, Prof Afaf Girgis, Ms Kerrie Graham, Ms Karen Johnston, Dr Janelle Jones, Dr Liane Lockwood, Dr Ann Maguire, Dr Maria McCarthy, Dr Jordana McLoone, Dr Francoise Mechinaud, Ms Sinead Molloy, Ms Lyndal Moore, Dr Michael Osborn, Dr Christina Signorelli, Dr Jane Skeen, Dr Heather Tapp, Ms Tracy Till, Ms Jo Truscott, Ms Kate Turpin, Prof Claire Wakefield, Ms Jane Williamson, Dr Thomas Walwyn, and Ms Kathy Yallop.

Financial disclosure: The authors have no financial relationships relevant to this article to disclose.

Funding: Christina Signorelli is supported by The Kids' Cancer Project. Claire Wakefield is supported by a Career Development Fellowship from the NHMRC of Australia (APP1143767). The Behavioural Sciences Unit (BSU) is proudly supported by the Kids with Cancer Foundation. The BSU's survivorship research program is funded by the Kids Cancer Alliance, The Kids' Cancer Project and a Cancer Council NSW Program Grant (PG16-02) with the support of the Estate of the Late Harry McPaul. ANZCHOG is supported by the Australian Government, through Cancer Australia's Support for Cancer Clinical Trials program. These funding bodies did not have any role in the study, nor did they have a role in the writing of the manuscript or the decision to submit it for publication.

Conflict of interest: No author has any conflicts of interest to disclose in relation to this manuscript.

ABSTRACT

Purpose: The long-term management of childhood cancer survivors is complex. Electronic health (eHealth) technology has the potential to significantly improve the management of late effects for childhood cancer survivors and assist their General Practitioners (GP) to coordinate their care. We assessed the acceptability of and perceived benefits and barriers to eHealth use amongst survivors, parents, and GPs.

Methods: We conducted semi-structured telephone interviews with survivors of childhood cancer, parents of childhood cancer survivors and their nominated GP. We described a hypothetical eHealth tool and asked participants how likely they would use the tool to manage their survivorship care and their perceived benefits and concerns for use of the tool. We used content analysis to synthesise the data using QSR NVivo12.

Results: We interviewed 31 survivors (mean age: 27.0), 29 parents (survivors' mean age: 12.6) and 51 GPs (mean years practising: 28.2). Eighty-five percent of survivors and parents (n=51), and 75% of GPs (n=38) indicated that they would be willing to use an eHealth tool. Survivors and parents reported that an eHealth tool would increase their confidence in their ability, and their GP's ability, to manage their survivorship care. GPs agreed that an eHealth tool would provide easier access to survivors' medical information and increase their capacity to provide support during survivorship. 7% of GPs (n= 4) and 43% of survivors (n=26) reported being hesitant to use eHealth tools due to privacy/security concerns.

Conclusion: Overall, eHealth tools were reported to be acceptable to childhood cancer survivors, their parents, and their GPs.

Survivors of childhood cancer live on average 60 years post-diagnosis,¹ during which their risk of developing cancer and/or treatment-related late effects increases.² Late effects may include second malignant neoplasms, life-threatening organ dysfunction, or poor psychosocial functioning.³ Many late effects can be prevented, managed or treated with appropriate surveillance through long-term follow-up care, and patient education of risk-factors for the development of late effects.⁴ However, around 45% of patients disengage from follow-up care during survivorship,⁵ due to survivors ageing out of paediatric healthcare,⁶ limited follow-up clinics/specialists,⁷ distance to clinics,⁸ survivors' limited knowledge of late effects⁹, and hospitals prioritising resources for high-risk survivors.¹⁰ Consequently, many survivors become reliant on their general practitioner (GP) to manage their care, or forego survivorship-specific care.

GPs can assist survivors to manage their late effects by scheduling screening tests, providing preventative lifestyle advice for late effects (e.g. diabetes and cardiovascular disease), and making referrals to specialists for complex co-morbidities.¹¹ Despite GPs willingness, they report insufficient knowledge and confidence providing paediatric cancer follow-up care.¹² Survivors also report a lack of confidence in their GP's provision of cancer-related care.¹³

Survivorship Care Plans (SCPs) may support GPs to deliver shared care with oncologist and other related specialists. SCPs provide personalised summaries of survivors' cancer history and treatment(s), and recommended screening or management.¹⁴ However, there is mixed evidence that SCPs improve childhood cancer survivors' long-term outcomes (e.g. survivors' knowledge, adherence to follow-up recommendations).^{15,16} As typically static, paper-based documents, SCPs easily become misplaced or outdated, across several decades of survivorship.

Cost-effective and high-quality electronic and mobile health tools (herein “eHealth tools”) have the potential to improve long-term survivorship care due to their inherently dynamic systems. eHealth interventions involve the promotion, prevention, treatment and maintenance of health (e.g. online programs), using technology (e.g. mobile applications) for the delivery of health-related services.¹⁷ There is a growing interest in using technologies to mitigate late effects in childhood cancer survivors. Cancer patients and survivors report positive attitudes towards eHealth tools for care management,^{18,19} preferring eHealth tools that enable active and frequent self-monitoring and more convenient survivorship care delivery.

Previous studies assessing patient perspectives have been broad, and varied in eHealth intervention type and content, as well as participant type (i.e. adult and child, in-patients’ and out-patients’ perspectives). Additionally, the acceptability of eHealth tools to healthcare professionals (including GPs) or parents of childhood cancer survivors remains unexplored, as do the preferences of survivors, parents and healthcare professionals for different types of eHealth tools. Given GPs and parents play a central role in the management of long-term childhood cancer survivorship care, their opinions and likely uptake of eHealth tools must also be considered. In this study we therefore aimed to understand:

1. The acceptability of eHealth tools for the management of survivorship care for survivors of childhood cancer, parents of young survivors, and GPs caring for childhood cancer survivors.
2. Survivors’, parents of survivors’ and GPs’ perceived benefits of and barriers to using eHealth tools for the management of survivorship care.

METHOD

This study was approved by the ethics authorities at participating hospitals and endorsed by the Australian and New Zealand Children’s Haematology Oncology Group (ANZCHOG).

We followed the consolidated criteria for report qualitative data research (COREQ) guidelines in reporting this study.²⁰

We collected data as a part of the ANZCHOG Survivorship Study, in two stages.²¹ Stage 1 involved childhood cancer survivors and parents completing a questionnaire, in which they could opt-in to complete a telephone interview and nominate their GP for Stage 2. In Stage 2 we invited nominated GPs to a telephone interview. Interviews were conducted one-on-one by trained female clinical psychologists and researchers. Interviewers had no prior relationship with interviewees and were independent to the research topic. We audio-recorded interviews, which were professionally transcribed verbatim. Participants did not review their transcripts. Interviews took 30-60 minutes

Interviewers described to participants a hypothetical eHealth tool (Table 1), defined as a free website or application f to help survivors manage their survivorship care by: i) storing all of their cancer-related information (e.g. treatment(s), screenings); ii) synchronising to their calendars for personalised surveillance reminders; and iii) providing access to tailored survivorship information based on their cancer/treatment(s). The definition and features of the prospective eHealth tool stemmed from previously developed, validated and tested paediatric oncology eHealth tools .²²

After describing the eHealth tool, we asked participants about 1) their willingness to use an eHealth tool such as the one described, 2) the perceived impact that an eHealth tool would have on their confidence to manage survivorship care, and 3) any concerns they may have about using an eHealth tool. We collected survivor and parent demographics using their paper or online questionnaires, whilst GPs' demographics and practice-related data were collected during their interview.

Participant selection

We identified eligible survivors using the electronic hospital medical records of recruiting hospitals. Survivors invited to the study were diagnosed with cancer before 16 years, were more than 5 years post-diagnosis, had finished cancer treatment, and were proficient in English. Parents of survivors under 16 years were invited to participate on behalf of their child. Participants were invited via a letter mailed to their home address.

Data Analysis

We used QSR NVivo12 Pro (QSR International Pty Ltd) for analysis. We analysed data using content analysis to enable the thematic organisation of responses, categorising the data according to broad predetermined themes. This process was guided by the methodology described by Clarke and Braun.²³ Two researchers (MS, MB) double-coded 15% of randomly selected interviews for consistency (99.4% agreement, $k=0.3$ coefficient). Discrepancies were resolved through discussion. Given the sample size and high concordance, one author (MS) coded the remainder.

RESULTS

Sample Characteristics (Table 2)

Of 636 questionnaire respondents, 380 (59.7%) volunteered to participate in the telephone interview. We interviewed 60 participants (31 survivors and 29 parents). As survivors and parents reported similar responses to eHealth tools during their interviews, their data was grouped together to report the findings.

Of the 160 GPs nominated by survivors and invited to participate via mail, 74 agreed to participate (46%) in an interview, of whom 51 were interviewed before we achieved data saturation. Of the GP sample, 29 (56.8%) were male and 33 (65%) worked in major cities. They reported practising for an average of 28.2 years (11.6, 8-60) and having seen an average

of 2.2 (2.1, 1-11) childhood cancer survivors in their practice. GPs were reimbursed with a \$50 gift card for their time.

Acceptability of eHealth tools to survivors and GPs (Figure 1)

Eighty-five percent of (n=51) survivors and parents agreed it would be acceptable to use an eHealth tool to manage their survivorship care; 7% survivors and parents were hesitant (n=4), and 8% were not interested in using an eHealth tool for their care management (n=5).

Seventy five percent of GPs (n=38) agreed that an eHealth tool would be acceptable and 25% (n=13) did not believe an eHealth tool would be appropriate for the management of care for survivors.

Perceived benefits for survivors/parents (Figure 1 and Table 3)

Among survivors and parents, perceived benefits associated with an eHealth tool included increased confidence in their ability to manage their care (63% ;n=38), greater confidence in their healthcare providers (55%; n=33), and increased convenience navigating healthcare systems (30%; n=18).

Potential to increase confidence in self-management

Participants reported that an eHealth tool could increase their own confidence to manage their or their child's survivorship care. Many noted that they "*simply can't remember everything that [they] need to do*" (father of Rhabdomyosarcoma survivor, aged 14) or were unsure of their cancer-related surveillance requirements. Participants noted that an eHealth tool might provide information to guide their care management. They also indicated that an eHealth tool would increase their confidence because the personalised surveillance alerts and schedules could ensure that they were "*doing the right thing*" (mother of Acute Lymphoblastic Leukaemia [ALL] survivor, aged 14).

Potential to increase confidence in health providers

It appeared that an eHealth tool could help to increase their faith in their healthcare provider. Survivors and parents asserted the belief that most GPs have limited knowledge of childhood cancer survivorship and were “*by definition ‘generalists’ ... not ‘specialists’*” (Hodgkin's lymphoma survivor, aged 32). Participants reported that providing patient-specific care recommendations to GPs and “*giving [GPs] more information is going to make me more confident*” (mother of Neuroblastoma survivor, aged 8).

Potential to improve healthcare system navigation

Survivors and parents valued the potential for an eHealth tool to store information about their cancer history. They noted that easy and convenient access to this information could significantly improve their information needs. Specifically, collating cancer-related information in one location as opposed to “*a collection of pieces of paper and scraps*” (mother of ALL survivor, aged 15) could minimise information loss and improve record keeping. Additionally, participants valued being able to easily transport their cancer history and show it to any healthcare provider.

Perceived benefits for GPs

GPs cited three key potential benefits of using an eHealth tool, including improved convenience to survivors information (19%, n=10), being able to provide better support for their survivor patients (17%, n=9), and increased patient responsibility (15%, n=8).

Potential for increased access to survivorship information

GPs cited having more convenient access to survivors' cancer-related information as a potential benefit of an eHealth tool. Specifically, having “*all the information in one place... would be far better than having lots of books and brochures*” (Male GP, 46 years practising). GPs noted that, as they rarely treat childhood cancer survivors, having increased access to

clinically relevant information, in one place, would be helpful when co-ordinating a patient's survivorship care, improving their efficiency

Potential for greater patient support

GPs indicated that an eHealth tool could provide greater support for their survivor patients because of the portability of information “*so if they move... there's an ongoing... guide to be used*” (female GP, 32 years practising). One female GP (12 years practising) recognised that “*unfortunately people don't always keep hold of [care plans]*” and an eHealth tool could provide increased support “*because the website can be stored rather than being lost [like] the piece of paper*”. Recognising that survivors are often a mobile population, GPs suggested that an eHealth tool would make “*people [feel] a lot more empowered*” (female GP, 14 years practising) because their survivor patients would have access to cancer-related information and follow-up care recommendations.

Potential to increase patient responsibility

GPs noted that eHealth tools could increase patient responsibility and their ability to “*come in and say, 'hey, doc, I'm due for this, or due for that'*” (male GP, 29 years practising). As a result, GPs' expected that an eHealth tool would increase awareness, responsibility, and self-management skills in their survivor patients. They cited that an increase in patient self-management skills could reduce GPs' high, and burdensome, level of survivor responsibility and create a shared and more equal level of care between the two.

Survivor and parent reported concerns (Figure 2 and Table 4)

Survivors and parents raised several potential concerns with using eHealth tools, including worries about the privacy of their information (43%, n=26), difficulty using technology (11%, n=7), and difficulty accessing eHealth tools (5%, n=3).

Privacy

The most frequently reported concern regarding eHealth tools was the level of security surrounding their information, highlighting the importance for “*secure access so the information couldn’t be corrupted ... or used by someone else that wasn’t supposed to*” (father of solid tumour survivor, aged 8). They noted that storing personal information online posed a risk of “*the information being hacked*” (father of Acute Lymphoblastic Leukaemia [ALL] survivor, aged 9) or viewed by unapproved individuals (e.g. insurance companies or future employers). Participants also expressed concern about their information being viewed by non-consensual individuals if there was an eHealth tool fault, or they had their device stolen.

Difficulty using technology

Some participants reported hesitation towards using eHealth tools due to their limited knowledge of technology, expressing themselves as “*not a computer person*” (father of Langerhans cell histiocytosis survivor, aged 10) or “*not ... super-duper [technologically] savvy*” (mother of ALL survivor, aged 16). However, some respondents were open to developing their technological skills and knowledge to learn how to use an eHealth tool.

Difficulty accessing technology

Survivors and parents reported concerns about the accessibility of eHealth tools, for example lack of access to the internet or a compatible device. Accessibility was particularly concerning for survivors living rurally/remotely as “*in rural areas we don’t always have internet*” (mother of Sacrococcygeal Teratoma survivor, aged 11).

GP reported concerns

GPs identified the following concerns towards eHealth tools use: a preference for letters (54%, n=27), lack of time (13%, n=7), and privacy/security concerns (7%, n=4).

Preference for written letters

Many GPs (54%) preferred receive patient-information as written letters, citing comfort in knowing where the patient information is stored. Conversely, they described eHealth tools as “*just not as easy to read as the usual mail*” (male GP, practising 24 years). GPs described a preference for existing practices stating that “*if it ain't (sic) broke, don't fix it*” (female GP, practising 24 years) noting the additional burden of having to remember a username and password.

Lack of time

GPs expressed a hesitation towards eHealth tools due to “*the time to use them.*” (female GP, 19 years practising), including learning how to use an eHealth tool, or adding information to it. Most GPs “*try to keep [consults] to 15 minutes...If [they're] looking after... someone with at least three system diseases... updating history is difficult let alone doing other things*” (male GP, 43 years practising). GPs suggested that eHealth tools would become undesirable if they consumed too much of their limited time.

Privacy/security concerns

GPs cited concerns regarding the privacy and security of patient data. GPs indicated that if they shared patient information between multiple platforms, it increased the risk of their patients' data being ‘hacked’ or viewed by non-consensual parties. GPs recognised the importance of therefore gaining patient consent and informing patients when, and in what way, their information would be shared.

DISCUSSION

This study assessed the acceptability of eHealth tools, perceived benefits, and barriers, to use among childhood cancer survivors, their parents, and their GPs. Most survivors/parents (85%), and GPs (75%) endorsed using an eHealth tool for cancer survivorship care management, echoing previous research in adult cancer survivors.^{18,19} Survivors/parents perceived benefits included increased confidence in their ability, and their GPs' ability, to manage their survivorship concerns, whilst barriers related mainly to their ability to use or access eHealth tools. GPs' perceived eHealth tool benefits included increased support for their patients and greater patient-initiated responsibility, however they indicated a preference for letters and a lack of time to learn/navigate a new system. All participants agreed that eHealth tools would make survivorship care management convenient but expressed concerns about the privacy/security of patient information.

Survivors' and parents' acceptance of eHealth tools may relate to its ability to meet more of their follow-up care needs. Survivors limited knowledge of their cancer history and follow-up,^{9,24} can lead to sub-optimal follow-up care engagement.²¹ Survivors/parents in our study valued easy access to personalised late effects information and follow-up recommendation reminders, consistent with other literature^{18,19} Frequent self-monitoring and convenient access to past cancer-related information aligns with the survivorship trajectory. Survivors in early survivorship often move away from their family GP or treating oncology centre.²⁵ Convenient access to survivors' cancer-related information may provide survivors with the flexibility to visit any GP or healthcare professional to receive their recommended follow-up care. A storage feature could accommodate survivors' needs by having convenient, portable, and personalised information with them when needed.

Participants suggested that eHealth tools could improve cancer care management. GPs report feeling more capable of providing care to survivors when provided with prescriptive

and patient-specific information.¹³ Participants also suggested eHealth tools could improve patients' responsibility for their care, potentially fostering a partnership between survivor and GP. Previous research indicates that both GPs and survivors advocate for a shared decision making model in their survivorship care management, which could be facilitate by eHealth tools.²⁶

Some participants reported being hesitance towards eHealth tools due to privacy/security concerns, aligning with adult cancer literature suggests mobile healthcare apps increase vulnerability to being hacked and medical record being revealed to non-consensual parties.²⁷ This reported barrier calls for potential developers to consider using formal privacy impact assessments and certifications as well as higher-level security measures, to provide users more confidence t.

The reported barriers demonstrated that eHealth tools should ideally be developed in collaboration with patient and provider users, who have potentially differing concerns (e.g. access to technology, or lack of time). Our findings highlight important features for the development of eHealth tools including an intuitive, easy-to-use interface, requiring little to no training, and capacity to store and share letters and upload patient information. It is also important that decision makers consider potential implementation challenges which could impact the feasibility of eHealth tool use in this population. eHealth software requires regular maintenance to keep pace with rapidly changing technologies and consumer expectations, which can be costly. Future research should more closely examine challenges of eHealth tool development, maintenance/ sustainability, and if these costs outweigh the potential eHealth tool benefits (e.g. increased surveillance).

Strengths/Limitations

This study is the first to demonstrate that survivors, parents, and GPs find eHealth tools acceptable/desirable, reflecting increasing acceptance of the digitalisation of healthcare services and expectations for accessible and personalised health information.^{28,29} Yet, the eHealth hypothetical tool proposed to participants in this study may not reflect all technology, possibly resulting in different perceived benefits and concerns. Participating GPs had been practising for nearly three decades on average, potentially influencing their desire to implement a new system into their practice.

Conclusions

eHealth tools may reduce some of the known challenges in childhood cancer survivorship, including limited cancer survivorship-specific knowledge, low confidence in care management, and disengagement from follow-up care, by providing high-quality childhood cancer follow-up care information. Our qualitative study demonstrates that survivors, parents, and their GPs generally endorse eHealth tools for the management of survivors' follow-up care. Participants identified concerns about eHealth tools that are critical to consider in the development of any new or existing tool.

REFERENCES

1. Smith MA, Seibel NL, Altekruse SF, et al. Outcomes for children and adolescents with cancer: challenges for the twenty-first century. *Journal of clinical oncology*. 2010;28(15):2625.
2. Bhakta N, Liu Q, Ness KK, et al. The cumulative burden of surviving childhood cancer: an initial report from the St Jude Lifetime Cohort Study (SJLIFE). *The Lancet*. 2017;390(10112):2569-2582.
3. Robison LL, Hudson MM. Survivors of childhood and adolescent cancer: life-long risks and responsibilities. *Nature Reviews Cancer*. 2014;14(1):61.
4. Landier W, Bhatia S, Eshelman DA, et al. Development of risk-based guidelines for pediatric cancer survivors: the Children's Oncology Group long-term follow-up guidelines from the Children's Oncology Group Late Effects Committee and Nursing Discipline. *Journal of Clinical Oncology*. 2004;22(24):4979-4990.
5. Schwartz CL. Long-term survivors of childhood cancer: the late effects of therapy. *The Oncologist*. 1999;4(1):45-54.
6. Nathan PC, Agha M, Pole JD, et al. Predictors of attendance at specialized survivor clinics in a population-based cohort of adult survivors of childhood cancer. *Journal of Cancer Survivorship*. 2016;10(4):611-618.
7. Signorelli C, Wakefield CE, McLoone JK, et al. Models of childhood cancer survivorship care in Australia and New Zealand: Strengths and challenges. *Asia-Pacific Journal of Clinical Oncology*. 2017;13(6):407-415.
8. Berg CJ, Stratton E, Esiashvili N, Mertens A. Young adult cancer survivors' experience with cancer treatment and follow-up care and perceptions of barriers to engaging in recommended care. *Journal of Cancer Education*. 2016;31(3):430-442.

9. Carretier J, Boyle H, Duval S, et al. A review of health behaviors in childhood and adolescent cancer survivors: toward prevention of second primary cancer. *Journal of adolescent and young adult oncology*. 2016;5(2):78-90.
10. Signorelli C, Wakefield CE, Fardell JE, et al. The impact of long-term follow-up care for childhood cancer survivors: A systematic review. *Critical reviews in oncology/hematology*. 2017;114:131-138.
11. Streefkerk N, Heins MJ, Teepen JC, et al. The involvement of primary care physicians in care for childhood cancer survivors. *Pediatric blood & cancer*. 2019;66(8):e27774.
12. Nathan PC, Daugherty CK, Wroblewski KE, et al. Family physician preferences and knowledge gaps regarding the care of adolescent and young adult survivors of childhood cancer. *Journal of Cancer Survivorship*. 2013;7(3):275-282.
13. Signorelli C, Wakefield CE, Fardell JE, et al. The role of primary care physicians in childhood cancer survivorship care: multiperspective interviews. *The oncologist*. 2019;24(5):710-719.
14. Horowitz ME, Fordis M, Krause S, McKellar J, Poplack DG. Passport for care: implementing the survivorship care plan. *Journal of oncology practice*. 2009;5(3):110-112.
15. Yan AP, Chen Y, Hudson MM, et al. Impact of survivorship care plans (SCPs) on adherence to surveillance for second malignant neoplasms (SMNs) and cardiac dysfunction in the Childhood Cancer Survivor Study (CCSS). In: American Society of Clinical Oncology; 2018.
16. Hill RE, Wakefield CE, Cohn RJ, et al. Survivorship Care Plans in Cancer: A Meta-Analysis and Systematic Review of Care Plan Outcomes. *The oncologist*. 2020;25(2):e351.

17. Borrelli B, Ritterband LM. Special issue on eHealth and mHealth: Challenges and future directions for assessment, treatment, and dissemination. *Health Psychology*. 2015;34(S):1205.
18. Ramsey WA, Heidelberg RE, Gilbert AM, Heneghan MB, Badawy SM, Alberts NM. eHealth and mHealth interventions in pediatric cancer: A systematic review of interventions across the cancer continuum. *Psycho-Oncology*. 2019.
19. Chan RJ, Howell D, Lustberg MB, et al. Advances and future directions in the use of mobile health in supportive cancer care: proceedings of the 2019 MASCC Annual Meeting symposium. *Supportive Care in Cancer: Official Journal of the Multinational Association of Supportive Care in Cancer*. 2020.
20. Tong A, Sainsbury P, Craig J. Consolidated criteria for reporting qualitative research (COREQ): a 32-item checklist for interviews and focus groups. *International journal for quality in health care*. 2007;19(6):349-357.
21. Vetsch J, Fardell JE, Wakefield CE, et al. “Forewarned and forearmed”: Long-term childhood cancer survivors’ and parents’ information needs and implications for survivorship models of care. *Patient education and counseling*. 2017;100(2):355-363.
22. Mertens A, Williamson R, Gilleland-Marchak J, Meacham LR, Cherven B, Edwards P. Evaluation of the adoption, acceptance, and satisfaction of SurvivorLink as a communication tool for pediatric cancer survivors. In: American Society of Clinical Oncology; 2016.
23. Braun V, Clarke V. *Successful qualitative research: A practical guide for beginners*. sage; 2013.
24. Syed IA, Klassen AF, Barr R, et al. Factors associated with childhood cancer survivors’ knowledge about their diagnosis, treatment, and risk for late effects. *Journal of Cancer Survivorship*. 2016;10(2):363-374.

25. Klosky JL, Cash DK, Buscemi J, et al. Factors influencing long-term follow-up clinic attendance among survivors of childhood cancer. *Journal of Cancer Survivorship*. 2008;2(4):225.
26. Ducassou S, Chipi M, Pouyade A, et al. Impact of shared care program in follow-up of childhood cancer survivors: An intervention study. *Pediatric blood & cancer*. 2017;64(11):e26541.
27. Kim H, Goldsmith JV, Sengupta S, et al. Mobile health application and e-health literacy: opportunities and concerns for cancer patients and caregivers. *Journal of Cancer Education*. 2019;34(1):3-8.
28. Dallery J, Kurti A, Erb P. A new frontier: Integrating behavioral and digital technology to promote health behavior. *The Behavior Analyst*. 2015;38(1):19-49.
29. O'Reilly-Shah V, Easton G, Gillespie S. Assessing the global reach and value of a provider-facing healthcare app using large-scale analytics. *BMJ global health*. 2017;2(3):e000299.

FIGURE LEGENDS

Table 1. Interview schedule for survivors and parents, and General Practitioners

Table 2. Clinical and demographic characteristics of participating childhood cancer survivors and parents of young survivors and GPs

Table 3. Survivors, parents, and GPs reported potential benefits relating to eHealth tools

Table 4. Survivors, parents, and GPs reported concerns relating to eHealth tools

Figure 1. Survivors', parents' and GPs' willingness to use eHealth tools for survivorship care coordinaiton

Figure 2. Survivors', parents' and GPs' perceived benefits and barriers for eHealth use in survivorship care management.

Table 1. Interview schedule for survivors and parents, and General Practitioners

<p><i>Survivors and Parents</i></p> <p>We are thinking about developing a free web or app product for childhood cancer survivors to help them manage their care. This website or app would:</p> <ul style="list-style-type: none"> - store all your information about your past treatments, tests, and screenings, - indicate a schedule, or lay out a plan, regarding which tests and screenings you should do in the future and when, synced to a calendar. - it could send you email reminders when something was due. - it could provide tailored information based on your cancer and treatments <p>Questions:</p> <ol style="list-style-type: none"> 1. Do you think you would use a product like this? [Probe: Why/why not?] 2. Would you feel more or less confident that you were managing your (child's) health well if you had a tool like this? 3. Would you feel more or less confident that your doctor(s) was able to manage your (child's) health well if they could use this tool? 4. Are there any other features you think would be important to include? 5. Are there any issues that would concern you?
<p><i>General Practitioners</i></p> <p>Survivors often have complex care pathways, including multiple specialist follow-up appointments and screening schedules, and they must manage these alone or with their GPs help. Our team are currently developing a free web or app product for childhood cancer survivors to help them manage their care. This website or app would:</p> <ul style="list-style-type: none"> - store patient information about their past treatments, tests, and screenings, - indicate a schedule, or lay out a plan, regarding which tests and screenings are needed in the future and when, synced to a calendar. - it could send email reminders to the patient or doctor when something was due. - it could provide tailored information based on the patients cancer and treatments <p>Questions</p> <ol style="list-style-type: none"> 1. Do you believe such e-tools may help survivors manage their care? [Probe: Why/why not?] 2. With your patient's permission, if you also had access to the information stored in this application, including their survivorship care plan, would you use this tool? [Probe: Why/Why not? Benefits/barriers?]

Table 2. Clinical and demographic characteristics of participating childhood cancer survivors and parents of young survivors and GPs

Characteristic	Survivors of childhood cancer <i>n</i> = 31 (51.7%)	Parents of young survivors <i>n</i> = 29 (48.3%)	GPs of childhood cancer survivors
Survivor sex, <i>n</i> (%)			
Male	11 (35.5)	20 (69.0)	29 (56.8)
Female	20 (64.5)	9 (31.0)	22 (43.2)
Relationship to survivor, <i>n</i> (%)			
Father	-	5 (8.3)	
Mother	-	24 (40)	
Ethnic background, <i>n</i> (%)			
Australian	27 (87.1)	22 (75.8)	
European	1 (3.2)	7 (24.1)	
Asian	1 (3.2)	-	
Other	2 (6.4)	-	
Area of residence, <i>n</i> (%) ^a			
Major City	27 (87.1)	24 (82.8)	33 (64.7)
Inner/outer regional	4 (12.9)	5 (17.2)	18 (35.3)
Diagnosis, <i>n</i> (%)			
Leukaemia	14 (45.2)	15 (51.7)	
Lymphoma	6 (19.4)	-	
Brain	1 (3.2)	2 (6.9)	
Other	10 (32.3)	12 (41.4)	
Treatment received, <i>n</i> (%)			
Surgery	12 (38.7)	9 (31.0)	
Chemotherapy	31 (100.0)	28 (96.6)	
Radiation	15 (48.4)	13 (44.8)	
Bone marrow transplant	4 (12.9)	6 (20.7)	
Age of survivor in years, mean (SD)	27.0 (6.3)	12.6 (2.3)	
Range	18 – 45	8 – 16	
Time since diagnosis in years, mean (SD)	18.9 (8.9)	9.55 (1.7)	
Range	6-42	7-13	
Time since treatment completion in years, mean (SD)	16.2 (8.6)	8.0 (2.0)	
Range	6-41	4-12	
Years Practising (SD)			28.2 (11.6)
Range			8-60
Number of survivor patients (SD)			2.2 (2.1)
Range			1-11

Abbreviations: **N:** Number of participants; **SD:** Standard deviation, —: not assessed or not applicable, **GP:** General Practitioner,

Numbers and percentages may not add up due to missing values and rounding errors.

^a According to Area of Remoteness Index Australia (ARIA) classifications.

Table 3. Survivors, parents, and GPs reported perceived benefits relating to eHealth tools

Theme	Illustrative quotes
SURVIVOR AND PARENTS PERCEIVED BENEFITS	
Potential increase confidence in self-management (N=38, 63.3%)	<p>“More confident...because I would know exactly what I would have to do, especially if they were reminding you as well.” (ALL survivor, aged 30)</p> <p>“A lot more confident [in myself]... only because I, I don't feel like sick or anything, but I just feel like, if there was anything else out there that I could be having, like to make sure that like I was like doing the best I can for my health.” (Female Acute Lymphoblastic Leukaemia survivor, aged 24)</p> <p>“I think I would feel more confident, because, yeah, based on how you described it, it sounds pretty useful and comprehensive.” (Male Acute Lymphoblastic Leukaemia survivor, aged 21)</p> <p>“more confident ... because it would be like more information to help me around what I need to do and stuff, instead of guessing” (Male Acute Lymphoblastic Leukaemia survivor, aged 27)</p>
Potential increased confidence in health providers (N=33, 55.0%)	<p>“I'd be more confident because, if [GP's] were a little unsure, they could go in, have a look, and sort of do their own research too and actually get a better idea of [my cancer].” (Brain cancer survivor, aged 25).</p> <p>“Yeah I'd feel more confident about that because at the moment the doctor relies on me to explain what I had and the treatment I had and I really don't know much about the treatment I had or about the nature of the cancer that I had” (Male Acute myeloid leukaemia, aged 29)</p> <p>“Yes I think if [GPs] had your entire history of course their care is going to be better.” (Female soft tissue Sarcoma survivor, aged 24)</p> <p>“I think they would feel more confident as well, because GPs treat people for such a broad range of things and I think sometimes we expect them to know everything and they don't and they can't” (Female Acute Lymphoblastic Leukaemia survivor, aged 29)</p>
Potential to improve healthcare system navigation (N=18, 30.0%)	<p>“You could go [to your doctor and say] my last test was this and I had it here and this was the result and if you want to speak to such-and-such doctor, that's who did the test” (ALL survivor, aged 29).</p> <p>“Because [the information] is there any time” (Father of a Wilm's tumour survivor, aged 10)</p> <p>“Just having, the information in front of you all the time.” (Female Acute Lymphoblastic Leukaemia survivor, aged 34)</p> <p>“If I was going to see my adult endocrinologist for the first time, I would take that along, I would show him that and sort of give him that. If I was having my annual check-up with the GP or something like that, again, I would gladly take that and show that as well.” (Female Brain Cancer survivor, aged 25)</p> <p>“It would be good because then I guess I'd have it all at my fingertips, and I wouldn't have to wait that whole year to see my haematologists. If I had a question that might be answered by simply running to the app or online about my treatment or anything else.” (Female Hodgkin's Lymphoma survivor, aged 25)</p>
GP'S PERCEIVED BENEFITS	
Potential for	“if I had [my patient] in consultation and I needed to see something or

increased convenient access to survivorship information (N=10, 19.6%)	<p>check something up, that's where it would be useful" (female GP, 19 years practising).</p> <p>"I guess now that I think about it I encounter childhood cancer survivors ... so it's just helpful when you come across problems or issues that you see infrequently just to have that extra reminder." (Male GP, 12 years practising)</p> <p>"Yeah. I think that it would be useful to have information available, that we could use in his care" (Male GP, 35 years practising)</p> <p>"would think authentic and concise, um [yep] guidelines on ... what I need to watch out for" (Male GP, 35 years practising)</p> <p>"Well, just information, that's all background information." (Male GP, 60 years practising)</p>
Potential for greater patient support (N=9, 17.6%)	<p>"if [patient's] move countries, they move state, it's – there's uh, uh, an ongoing, um, ah, ongoing, ah, guide to be used" (Female GP, 60 years practising)</p> <p>"I think that the electronic stuff, you could oblige your patients and the doctors who are looking after them to use electronic health records... so that she could use that at any GP really. If it's an app that she can open up on her own phone that says this, then that's fine as well." (Male GP, 43 years practising)</p> <p>"I think the use of modern technology and social media or links like that with reminders or calendar apps or all sorts of things, or information and links, is really helping the patients in all levels of care." (Male GP, 29 years practising)</p>
Potential to increase patient responsibility (N=8, 15.6%)	<p>"I would be very happy if a patient came along with a folder full of information, because, you know, it makes your life easy" (female GP, 14 years practising).</p> <p>"nowadays, with technology, and you know, people are a lot more empowered with their own health [yeah], that then I think helps if they depend on the person as to how active they are, but [yeah] I think it would be popular" (Female GP, 14 years practising)</p> <p>"I think that's good for them, so they don't get complacent, so they can keep a regular eye." (Male GP, 23 years practising)</p>

Table 4. Survivors, parents, and GPs reported concerns relating to eHealth tools

Theme	Illustrative quote
SURVIVOR AND PARENTS PERCEIVED CONCERNS	
Privacy (N=26, 43.3%)	<p>“if someone picks up the app on my phone they can then get access to all my stuff” (Germ Cell Tumour survivor, aged 24).</p> <p>“I guess with anything I guess the confidentiality you wouldn’t want anyone anybody tapping ... more or less the privacy issue” (Mother of a Rhabdomyosarcoma survivor, aged 15)</p> <p>“...just the security. So personalised information not getting out there. Um, only me having access to it.” (Male Acute myeloid leukaemia survivor, aged 29)</p> <p>“I’d probably have it password protected on each person’s phone because I guess it’s quite personal information.” (Male Hodgkin's lymphoma survivor, aged 25)</p> <p>“Oh, I mean, only the obvious which would be confidentiality and knowing that random people don’t have access to information about [child]” (Mother of a Acute Lymphoblastic Leukaemia survivor, aged 15)</p>
Difficulty using technology (N=7, 11.6%)	<p>“I am not tech savvy... if it was there in front of me, like for me to go like I’m not very good with stuff like that” (Father of Wilm’s tumour survivor, aged 10)</p> <p>“Personally I still like to have paper copies, or um a stored file that’s not on my phone ... I would still want there to be a person behind it, rather than just uh, completely automated.” (Male Acute myeloid leukaemia survivor, aged 29)</p> <p>“I’m really not good with technology” (Female Wilm’s tumour survivor, aged 45)</p> <p>“I could give it a shot. I missed out on the computer stuff at school.” (Female, Hodgkin’s Lymphoma survivor, aged 25)</p>
Difficulty accessing technology (N= 3, 5%)	<p>“Probably the phone, my phone’s not an iPhone so...I’ve only got the HTC and isn’t it Android or something like that? So I’d probably use the website more than the phone” (Mother of a solid tumour survivor, aged 11)</p> <p>“I do minimal even though I’m on my phone and stuff, got a smartphone and stuff, but I do what I need to do and kind of that’s it.” (Father of Wilm’s Tumour survivor, aged 10)</p>
GP’S PERCEIVED CONCERNS	
Preference for written letters (N= 27, 45%)	<p>“I don't know, I'd just like it all written down.” (Female GP, 30 years practising)</p> <p>“I normally prefer in a paper version so that I can take it around with me and read it whenever I like.” (Male GP, 39 years practising)</p> <p>“I'm of an age where letters are still important for me, because I can put them in the in basket and not lose them. Emails tend to drift out over the horizon and blow off the screen.” (Male GP, 43 years practising)</p> <p>“Yeah, I think a letter is best.” (Female GP, 28 years practising)</p> <p>“I think letter would be by far and away the most useful. A letters force you to read them, electronic things you, whilst you read them and scan them you can then park them and they don’t seem to stick in my memory so well” (Male GP, 43 years practising)</p>

	<p>"I would never use [an eHealth tool] ...because you're going to ask me to have a login and a password and I'm not going remember one." (male GP, 12 years practising).</p>
<p>Lack of time (N=7, 13.7%)</p>	<p>"Look, to be quite frank I don't have time for a lot of the up to date technology and stuff and if I - I probably wouldn't have time to use it" (Female GP, 24 years practising)</p> <p>"I think the biggest issue is time." (Male GP, 35 years practising)</p> <p>"well I guess in terms of barrier such as time" (Female GP, 19 years practising)</p>
<p>Privacy and security concerns (N=4, 7.8%)</p>	<p>"We have a problem with this secure messaging stuff at a national level" (Male GP, 37 years practising)</p> <p>"I'm not sure how the privacy issues would relate to it but I would be thinking that the patient would get consent and they give consent to me to be able to access it as well" (Male GP, 12 years practising)</p>