

**Table 1: Baseline characteristics**

	All Patients Median (IQR) n=120	Group 1 Median (IQR) n=60	Group 2 Median (IQR) n=60	p value
<i>Preoperative</i>				
Age (years old)	67 (51-85)	66.5 (51-80)	67 (52-85)	0.692
Preoperative PSA (ng/dL)	4.8 (0.7-43)	5 (0.8-12.2)	4.6 (0.7-43)	0.944
Prostate volume (mL)	95 (35-266)	95 (35-266)	91 (48-242)	0.935
ASA score (%)				0.681
I	20 (16.7%)	13 (21.7%)	7 (11.7%)	
II	75 (62.5%)	33 (55%)	42 (70%)	
III	25 (20.8%)	14 (23.3)	11 (18.3%)	
IV	-	-	-	
Antithrombotic	21 (17.5%)	8 (13.3%)	13 (21.7%)	0.230
Previous prostate biopsy	31 (25.8%)	12 (20%)	19 (31.7%)	0.144
Q max	7.8 (3-20)	7.9 (4-19)	7.8 (3-20)	0.809
IPSS	24 (10-32)	24 (13-32)	24 (10-31)	0.418
PVR	100 (70-140)	100 (80-130)	105 (70-140)	0.352
QoL		4 (3-6)	4 (3-6)	0.872

IQR: Inter Quantile Range, PSA: Prostate specific antigen, ASA: American Society of Anesthesiologists, Q max: Maximum flow rate, IPSS: International prostate symptoms score, PVR: Postvoiding residual volume, and QoL: Quality of life.