

# PREVALENCE OF HAMSTRING TIGHTNESS IN SCHOOL CHILDREN IN THE UNITED ARAB EMIRATES.

## ABSTRACT

## BACKGROUND:

Back pain amongst children is on the rise exponentially. Hamstring tightness is known to have cause/effect relationship with low back pain. Presence of hamstring tightness is also associated with increase in risk of muscular strains. The purpose of this study was to assess the prevalence of hamstring tightness in school going children in the United Arab Emirates. To our knowledge, this is the first such reported study in the Middle East.

## METHODS:

All children of a school based in Dubai, UAE from grade 2 to grade 10 aged 7-15 years were assessed by a team of 5 physiotherapists. Hamstring tightness was evaluated by standard active knee extension [AKE] test in the school medical department. A total of 317 children were assessed [634 limbs] by two observers and any inability to actively extend the knee completely while the hip being at 90-degree flexion was considered as hamstring tightness.

## RESULTS:

Out of 317 subjects, 287 [90.53%] were found to have at least some hamstring tightness. There was a statistically significant difference between genders: boys showing more prevalence. All age groups across the school showed similar pattern. This may be the highest ever recorded prevalence of hamstring tightness in school children in any reported study in English literature.

## CONCLUSION:

An alarming proportion of children in a Dubai based school have hamstring tightness. Wider studies with larger numbers should be undertaken so as to ascertain if this is in some way related to lack of active lifestyle in the UAE where gadget-dependency of younger generation as well as prolonged sitting might be a causative factor. Whether such high prevalence is pathological or physiological also needs to be addressed. School authorities and parents in UAE should engage the children in regular stretching programs.

## 29 INTRODUCTION

30 The musculoskeletal health of the pediatric age group doesn't get enough attention and  
31 recognition worldwide [1]. Various conditions and disorders (e.g., musculoskeletal deformities,  
32 congenital/developmental syndromes, mental health, obesity) that begin during childhood may have a  
33 lot of implications to a child's daily function, quality of life, family relations and psychological  
34 wellness [1,2]. The effects of these health issues may stay and be seen in adulthood [3]. However, back  
35 pain, which has not been seen frequently in children before, has been increasingly reported recently [4].  
36 The stated percentage has risen from 2-11% to 27-51% in recent studies [5-8]. Other reports have noted  
37 a lifetime prevalence of 70-80% in patients up to 20 years of age [9]. Now, various reasons are being  
38 considered as to being the cause of childhood low back pain. Some of them are, to heavy school bag  
39 use, reduced physical activity, too much video-game playing and/or television viewing, improper sleep,  
40 and obesity [8,10,11-13]. Hamstring tightness has been reported to be the cause of posterior pelvic  
41 tilting, reduced lumbar lordosis and exacerbation of existing pain in patients with low back pain [14]. A  
42 number of studies have shown that there lies a strong link between hamstring flexibility and low back  
43 pain [15.]

44 Hamstring muscles play an important role in managing the stride length during running; if the  
45 stride length is longer, then fewer contractions cycles are needed to cover the same distance. This  
46 equates to a conservation of energy, allowing for greater all longer performance prior to fatigue and a  
47 reduction in chance of injury [16,17].

48 Inability to extend the knee completely when the hip is flexed, accompanied by discomfort or  
49 pain along the posterior thigh and/or knee is considered as hamstring tightness [18]. Tight hamstring  
50 muscles increase the patellofemoral compressive force because of the increased passive resistance  
51 during the swing phase of ambulation and running [19].

52 This study aims to find the prevalence of hamstring tightness amongst school children in United  
53 Arab Emirates where such statistics are not available as yet.

54

55

56

57

## 58    **METHODS**

59            All grade 2 to grade 10 students of a school based in Dubai were included in this study.  
60    Permission from the relevant authorities was obtained prior to the assessment. The subjects as well as  
61    their parents were briefed about the Active Knee Extension (AKE) test [16] and the outcome. AKE test  
62    remains the gold-standard for hamstring flexibility assessment and has high intra-rater and inter-rater  
63    reliability.

64            A total of 317 subjects between the age of 7 to 15 of grades 2 to 10<sup>th</sup> were assessed. The average  
65    age was 9.2 years. The subjects included 181 boys and 136 girls. The AKE test was conducted by five  
66    licensed physiotherapists practicing in the UAE. The team comprised 2 male and 3 female  
67    physiotherapists. Each student was assessed by at least 2 examiners to confirm the findings. There was  
68    an administrative staff included to record the details of the participants. The subjects were called to the  
69    setup by the increasing order of the class grades. Each student was explained about the standard AKE  
70    test and the test was conducted as per described positioning of hips and knees bilaterally. Children of  
71    all grades had their data recorded as positive and negative and compared at the end of the study. Any  
72    inability to actively extend the knee completely while the hip being flexed at 90 degree flexion was  
73    considered as hamstring tightness. Results were compared between sexes by Chi Square test and any  
74    statistically significant differences were identified.

### 75    **Active Knee Extension Test [16]**

76            Subjects were instructed to assume a supine position on the examination table and both the hip  
77    and knee flexed 90° of one lower limb, the other limb being fixed to table in full extension (Fig.1). The  
78    subject then grasped behind the examining knee with both hands to stabilize the hips at 90 degree of  
79    flexion. Marking was done to identify lateral malleolus and lateral femoral condyle. The subject  
80    actively extended the knee in turn as much as possible. Then based on that, the subject's hamstring  
81    flexibility was assessed. A subject who could completely extend the knee was considered not to have  
82    any hamstring tightness. Any shortfall in making the knee extend to 180° was considered tightness.  
83    Each child was examined by two therapists and concurrence of hamstring being tight was established.

84    **Exclusion criteria** Only those subjects who had no previous history of congenital or developmental  
85    disorders or trauma were included.

86

87  
88  
89  
90  
91  
92  
93  
94  
95  
  
96  
97  
98  
  
99  
  
100  
  
101  
  
102  
  
103  
  
104  
  
105  
  
106  
  
107  
  
108  
  
109  
  
110  
  
111

**RESULTS:**

Out of the 317 subjects, 287 had hamstring tightness while 30 did not (Fig 3). Among those who presented with tightness 171 subjects were males and 116 females. Among those who did not have tightness, 10 were male and 20 female. Amongst all grades, the maximum number of children (104) belonged to grade 2 where 89.4% showed hamstring tightness. All [100%] children of grade 5 and grade 9 showed positive results for hamstring tightness although the number of students were only 18 and 7 respectively. On the other hand, out of 23 children of grade 8, only 78.2% showed hamstring tightness. Grade wise comparison is illustrated in Fig 2.

The outcome showed an average of 90.53% having hamstring tightness. Hamstring tightness was more among the males being 95% and lesser in females with average 85.2% which is statistically significant  $p < .05$  by chi square test.

112

## 113 **DISCUSSION:**

114 Over the years it has become clear that LBP occurs not only in adults but also in the young  
115 [20,21]. Some previously reported studies provide linkage of hamstring tightness with low back pain.  
116 Hamstring tightness causes pelvic tilt thereby changing the biomechanics of lumbar region and causing  
117 back pain [23]. Stiffness of one muscle group can cause compensatory movement at an adjoining joint  
118 that is controlled by muscles or joints with less stiffness [24]. Appropriate and everyday stretching of  
119 the hamstring muscles should be taught to all age groups, especially before age 30 as tightness  
120 increases more at that time [25].

121 The above study was conducted on seemingly healthy group of children by AKE test that has an  
122 excellent inter-rater and intra-rater reliability [26]. It was alarming to see a significant proportion of  
123 school going children having hamstring tightness in this study who showed at least some degree of  
124 hamstring tightness. This leads to a concern about its etiology. The reasons for this can be  
125 multitudinous, which include growth spurts, prolonged sitting while spending more time on studying,  
126 video gaming, digital gadget usage and viewing television rather than going outdoors for play or other  
127 activities. The hot climate of the country can be a reason due to which the children avoid spending  
128 more time in the outdoors. Even the homework and class assignments and most of the projects are done  
129 on computers which only adds to hours in front of the computer and other devices. Increased sitting and  
130 inactivity causes a certain fixture in posture due to which the hamstring muscle may develop tightness  
131 and recent research points out towards this association [27]. Additionally, sudden growth spurts are  
132 also known to cause the stretching of the muscle leading to reduced flexibility [28].

133 This indeed is an issue which needs immediate attention and also measures to curb or get it in  
134 control. If not, this can lead to bigger health issues in early adulthood. It would be interesting for the  
135 global community to address if this is physiological or pathological by conducting such studies on a  
136 larger scale involving many schools within their countries.

137 The study involved subjects as young as age 7, and the presence of hamstring tightness in them  
138 was indeed a concerning issue. School teachers, especially physical education teachers can help in this  
139 condition. Physiotherapists should also include routine screening of hamstring tightness and advice  
140 stretching exercises into the treatment program of patients suffering from musculoskeletal disorders of  
141 the lower limb and the lower back.

Limitations of this study include the fact that every grade didn't have equal number of students and the senior grades has fewer subjects available thus making this a non-homogenous mix of available subjects. Also, we did not measure and categorize the degree of hamstring tightness as there is no established parameters of grading of hamstring tightness available in the literature. We considered any degree of loss of full extension of knee in the AKE test to represent hamstring tightness whereas some authors consider 20° flexion in adults to be normal [29]. Further, this study didn't aim at recording any improvements in degree of hamstring tightness.

## **CONCLUSION:**

As far as the authors are aware, this is the first study conducted in UAE and the Middle East to assess prevalence of hamstring tightness in school children. The result of the above study showed that the hamstring tightness among school going children is highly prevalent in the UAE. The participants of the study were normal healthy children who were not narrowed down through any category, but were selected from all grades available on the day for evaluation. Thus, such a high percentage of positive results in them can only speak for the rest of the pediatric population. Although the sample in this study wasn't equally distributed amongst ages, most of the age groups were found having hamstring tightness and this shows that the problem is not age specific. Females showed statistically significant lesser percentage of hamstring tightness than male participants which is in line with other researchers' findings [30]. Simple methods of hamstring stretching techniques can be applied by school and parents to prevent and treat hamstring tightness. Also, increased outdoor activity whilst reducing time children spend on sitting may be helpful. A study involving large number of children would be more beneficial to get an even better understanding of the prevalence of hamstring tightness in children as such data is scarcely available.

## **CONFLICT OF INTEREST: NONE**

## REFERENCES

1. Shudy M, De Almeida ML, Ly S et al. Pediatrics 2006 Dec;118 Suppl 3:S203-18.
2. Rennick JE, Johnston CC, Lambert SD et al. Measuring psychological outcome following paediatric intensive care unit hospitalization: psychometric analysis of the Children's Critical Illness Impact Scale. Paediatric Crit Care Med 2011;12:635-642.
3. O'Connor EA, Evans CV, Burda BU, Walsh ES, Eder M, Lozano P. Screening for obesity and intervention for weight management in children and adolescents: evidence report and systematic review for the US Preventive Services Task Force. JAMA 2017;317:2427-2444.
4. Jennifer Hwang, Philip K Louie, Frank M. Phillips, Howard S. An, Dino Samartzis. Low back pain in children: a rising concern. Eur Spine Journal 2018;28:211-213.
5. Taimela S, Kujala UM, Salminen JJ, Viljanen T. The prevalence of low back pain among children and adolescents. A nationwide, cohort-based questionnaire survey in Finland. Spine (Phila Pa 1976) 1997 May 15;22(10):1132-1136.
6. Swain MS, Henschke N, Kamper SJ, Gobina I, Ottova-Jordan V, Maher CG. An international survey of pain in adolescents. BMC Public Health 2014;14:447
7. King S, Chambers CT, Huguet A, MacNevin RC, McGrath PJ, Parker L, MacDonald AJ. The epidemiology of chronic pain in children and adolescents revisited: a systematic review. Pain 2011;152:2729-2738.
8. Balague F, Troussier B, Salminen JJ. Non specific low back pain in children and adolescents: risk factors. Eur Spine J 1999; 8:429-438.
9. MacDonald J, Stuart E, Rodenberg R. Musculoskeletal low back pain in school-aged children: a review. JAMA Pediatr 2017;171:280-287.
10. Gunzburg R, Balague F, Nordin M et al. Low back pain in population of school children. Eur Spine J 1999;8(6):439-443.
11. Samartzis D, Karppinen J, Mok F, Fong DY, Luk KD, Cheung KM. A population-based study and diminished functional status. J Bone Joint Surg Am 2011;93:662-670.
12. Negrini S, Carabalona R. Backpacks on! School children's perceptions of load, associations with

197 back pain and factors determining the load. *Spine* 1976;27:187–195.

198 13 Szita J, Boja S, Szilagyi A, Somhegyi A, Varga PP, Lazary A. Risk factors of non-specific spinal  
199 pain in childhood. *Eur Spine J* 2018;27:1119–1126.

200 14. Wendell L. Flexibility and Muscular Strength. *Journal of Physical Education, Recreation & Dance*.  
201 1988;59:37-40.

202 15. Abeera H, Waqar A, Muhammad NB, Hasneeza H. Relationship between hamstring flexibility  
203 disability related to low back pain. *Int J Rehabil Sci* 2018;7(1):20-23.

204 16. Baltaci G, Un N, Tunay V, et al.: Comparison of three different sit and reach tests for measurement  
205 of hamstring flexibility in female university students. *Br J Sports Med* 2003;37:59–61.

206 17. Barlow.A , Clarke.R, Johnson N, et al. Effect of massage of the hamstring muscle group on  
207 performance of the sit and reach test. *Br J Sports Med* 2004;38:349-351.

208 18. NR Shakya, S Manandhar. Prevalence of hamstring tightness among undergraduate Physiotherapy  
209 students of Nepal using passive knee extension angle test. *Int Journal Scientific and Research*  
210 *Publication* 2018;8(1):182-187.

211 19. Akinpelu AO, Bakare U, Adegoke BOA. Influence of age on hamstring tightness in apparently  
212 healthy Nigerians. *Journal of The Nigeria Society of Physiotherapy* 2005;15(2):35-41.

213 20. Balagué F, Nordin M. Back pain in children and teenagers. *Baillière's Clin Rheumatol* 1992;6:575-  
214 593.

215 21. Szpalski, M., Gunzburg, R., Balagué, F. et al. *Eur Spine J* 2002;11:459-464.

216 22. Fieldman H: Relative contribution of the back and hamstring muscles in the performance of the toe  
217 touch test after selected extensibility exercises. *Res Q Am Assoc Health Phys Educ* 1967;139:518-523.

218 23. Janda V, Frank C, Liebenson C, Evaluation of muscular imbalance. *Rehabilitation of the spine:a*  
219 *practitioner's manual*.1996;6:97-112[30].

220 24. José M. Muyor, Pedro A. López-Miñarro, Fernando Alacid. The relationship between hamstring  
221 muscle extensibility and spinal postures varies with the degree of knee extension. *Journal of Applied*  
222 *Biomechanics* 2013;29:.678–686.

223 25. Loutsch RA, Baker RT, May JM, Nasypany AM. Reactive neuromuscular training results in  
 224 immediate and long term improvements in measures of hamstring flexibility: A case Report. *Int J*  
 225 *Sports Phys Ther* 2015;10:371–377.

226 26. Mohamad Shariff A Hamid, Mohamed Razif Mohamed Ali, Ashril Yusof. Interrater and Intrarater  
 227 Reliability of the Active Knee Extension (AKE) Test among Healthy Adults. *J Phys Ther Sci*  
 228 2013;25(8):957–961.

229 27. Ghulam F, Muhammad MQ, Jawad UH, Ayesha B. Extended sitting can cause hamstring tightness.  
 230 *Saudi J Sports Med* 2017;17(2):110-114.

231 28. Micheli LJ, Stratford PW, Tarnopolsky M, Lotter A. Is adolescent growth associated with changes  
 232 in flexibility? *Clin J Sport Med* 2000;10(1):76.

233 29. Hansberger BL, Loutsch R, Hancock C, Bonser R, Zeigel A, Baker RT. Evaluating the relationship  
 234 between clinical assessments of apparent hamstring tightness: a correlational analysis. *Int J Sports Phys*  
 235 *Ther.* 2019 Apr;14(2):253-263.

236 30. Marshall PW, Siegler JC. Lower hamstring extensibility in men compared to women is explained  
 237 by differences in stretch tolerance. *BMC Musculoskelet Disord* 2014;15:223.

238

239

240

241

242

243

244 Fig 1.



245

246 TEST STARTING POSITION

AKE



247

248 AKE TEST FINAL POSITION

249

## Grade Wise Comparison





Fig 2. Grade of school wise comparison of presence or absence of hamstring tightness.



254

255

256 Fig 3. Overview of hamstring tightness in 317 school children