

**Table 4:** Comparison of final symptoms/affected systems.

<b>Symptoms/diagnosis</b>	<b>2019 (n = 11704)</b>	<b>2020 (n = 11497)</b>	<b>P-value<sup>a</sup></b>
<b>Suspected infection of COVID-19</b>	-	1867 (16.2%)	-
<b>Infected with COVID-19</b>	-	1554 (13.5%)	-
<b>Fever</b>	750 (6.4%)	357 (3.1%)	<b>0.000</b>
<b>Respiratory system</b>	912 (7.8%)	400 (3.5%)	<b>0.000</b>
<b>Cardiovascular system</b>	713 (6.1%)	372 (3.2%)	<b>0.000</b>
<b>Gastrointestinal system</b>	1042 (8.9%)	866 (7.5%)	<b>0.000</b>
<b>Neurological system</b>	774 (6.6%)	568 (4.9%)	<b>0.000</b>
<b>Hematological system</b>	145 (1.2%)	143 (1.2%)	0.973
<b>Psychiatric causes</b>	717 (6.1%)	457 (4.0%)	<b>0.000</b>
<b>Traumatic causes</b>	4585 (39.2%)	3306 (28.8%)	<b>0.000</b>
<b>Poisoning</b>	450 (3.8%)	300 (2.6%)	<b>0.000</b>
<b>Other causes</b>	1548 (13.2%)	1266 (11.0%)	<b>0.000</b>
<b>Arrest</b>	68 (0.6%)	41 (0.4%)	<b>0.012</b>

a: Chi square test (n, %).