




Species	Image	Feeding strategy	Dietary preference	Habitat	DVM	References
Order: Calanoida Family: Temoridae <i>Temora stylifera</i> total length 0.956 ± 0.048 mm		filter feeder	tendency towards herbivory small algae, heterotrophic flagellates, dinoflagellates, diatoms, cyanobacteria	restricted to epipelagic zone (0-200 m)	no DVM	Benedetti <i>et al.</i> 2016 Zamora-Terol <i>et al.</i> 2020
Order: Cyclopoida Family: Oithonidae <i>Oithona nana</i> total length 0.506 ± 0.003 mm		active ambush feeder	omnivore small algae, nauplii and copepodites, autotrophic and heterotrophic microflagellates, dinoflagellates, and ciliates	restricted to the epipelagic zone (0-200 m)	weak (occurs within tens of meters)	Benedetti <i>et al.</i> 2016 Temperoni <i>et al.</i> 2010
Order: Calanoida Family: Centropagidae <i>Centropages ponticus</i> total length 0.845 ± 0.038 mm		mixed feeder	omnivore feeds on a wide spectrum of prey from small algae to yolk-sac fish larvae, ciliates or dinoflagellates, cyanobacteria	epipelagic zone (0-200 m) affiliated with neritic environments	NA	Benedetti <i>et al.</i> 2016 Calbet <i>et al.</i> 2007 Zamora-Terol <i>et al.</i> 2020