


Helium and carbon isotope characteristics of Andean Convergent Margin geothermal fluids

P.H. Barry¹, D.V. Bekaert¹, A.M. Seltzer¹, J. Curtice¹, J. M. de Moor², G.L. Jessen³, D. Giovannelli^{4,1}, M. Schrenk⁵, J. Buongiorno⁶, A. Chiodi⁷, C.J. Ramírez⁸, T.J. Rogers⁹, K.G. Lloyd⁹



¹Marine Chemistry and Geochemistry Department, Woods Hole Oceanographic Institution, Woods Hole, MA, USA, ²Observatorio Volcanológico y Sismológico de Costa Rica (OVSICORI), Universidad Nacional, Costa Rica, ³Instituto de Ciencias Marinas y Limnológicas, Universidad Austral de Chile, Valdivia, Chile, ⁴Department of Biology, University of Naples "Federico II", Naples, Italy, ⁵Department of Earth and Environmental Sciences, Michigan State University, MI, USA, ⁶Division of Natural Sciences, Maryville College, Maryville, TN, USA, ⁷Instituto de Bio Y Geociencias Del Noroeste Argentino, Salta, Argentina, ⁸Servicio Geológico Ambiental (SeGeoAm) Heredia, Costa Rica, ⁹Department of Microbiology, University of Tennessee, TN, USA

Background:
Subduction zones are the interface between Earth's interior (crust and mantle) and exterior (atmosphere and oceans), where carbon and other volatiles are actively cycled between terrestrial reservoirs by plate tectonics (e.g., Bekaert et al., 2021).

Why helium and carbon?
Helium is a sensitive tracer of mantle inputs and can be used to deconvolute mantle and crustal volatile pathways in arc settings. Carbon is highly reactive and isotope fractionation patterns can be used to understand the dominant processes in the arc system (e.g., Barry et al., 2019).

Driving Research Question:
How do tectonic scale features (i.e., sediment thickness of incoming slab, slab dip, crustal thickness) affect volcanic outputs and the overall volatile budget?




Fig. 1 - Schematic of a convergent margin showing the types of samples to be collected: A) plume fluxes, B) volcanic gases, C) forearc fluids, and leveraged IODP data: D) submarine forearc and E) downgoing slab.




Fig. 2 - The Andean Convergent Margin (ACM). Red circle = volcanoes. Map is from Tilling et al., 2009.






Fig. 3 - Along-arc variations in tectonic-scale phenomena. Volcanoes = grey dots. Blue box is our CVZ study area and red box is our SVZ study area.

Panel A) Offshore sediment thickness, with contour lines every 100 m and bold contour lines every 500 m.
Panel B) Depth of the slab with contour lines every 20 km and bold contour lines every 50 km. Convergence rate (70km/Ma) is also shown.
Panel C) Crustal thickness along the ACM, whereby dot size indicates crustal thickness.



Summary:

- 1) We report new He and C isotope and abundance data for 33 deeply-sourced fluid and gas samples in the CVZ of Argentina and the SVZ of Chile.
- 2) Air-corrected He isotopes from the CVZ range from 0.11 to 2.58 R_A (n=17), SVZ range from 0.65 to 5.03 R_A (n=16). New data are plotted alongside published data in figs.
- 3) Lowest He isotopes are in the CVZ = more significant crustal (radiogenic) contributions. The crustal thickness in the CVZ is up to 70 km, significantly more than in the SVZ (35-45 km). Overriding crustal thickness exerts a primary control on the extent of fluid-crust interaction.
- 4) Carbon isotopes vary between -12.0 to -1.2‰ (vs. PDVB) in the CVZ and CO₂/³He values vary over 3 orders of magnitude (9.0 × 10⁸ to 1.1 × 10¹²). In the SVZ, carbon isotopes (δ¹³CO₂) vary between -14.0 to -8.7‰ (vs. PDVB) and CO₂/³He values vary by nearly 3 orders of magnitude (4.7 × 10⁷ to 3.9 × 10¹⁰). Lower δ¹³CO₂ and CO₂/³He values in the SVZ are consistent with calcite precipitation in shallow-level (upper 5 km) hydrothermal systems (e.g., Ray et al., 2009).
- 5) Offshore sediment thickness, and thus delivery of isotopically light C is higher in the SVZ and could explain some of the lighter C isotopes. No correlations is detected with slab angle.

Open Questions:

- 1) Why is SVZ seemingly more affected by removal of carbon? 2) What is the role of biology in the SVZ? 3) What is the role of organic soil carbon in the SVZ? Adding fluid residence time information (tritium and ¹⁴C) to future studies will help differentiate between deep and surface processes and enable flux calculations.

Fig. 4 - Latitudinal variations in He-isotopes in fluids and gases. The CVZ has significantly lower He isotopes, interpreted to reflect interaction with thicker (i.e., more radiogenic) overriding crust vs. the SVZ, which retains a more pristine mantle signal. Solid symbols are data from this study and crosses are published data.

Fig. 5 - Latitudinal variations in CO₂/³He. All CVZ samples have gained excess CO₂ relative to mantle ³He, whereas SVZ values are typically lower and more variable.





Fig. 6 - CO₂/³He vs. ³He/4He. Higher CO₂/³He and lower ³He/4He in the CVZ are attributed to crustal assimilation, whereas SVZ are more mantle like.

Fig. 7 - CO₂/³He vs. C isotopes. Almost all CVZ samples are above mantle values, suggesting crustal assimilation. SVZ values are more disparate, potentially suggesting C removal (due to calcite ppt or biological processes?; Barry et al., 2019) or a light subduction signature associated with organic sediments in the slab.