

# Differentiating EDRs from the Background Magnetopause Current Sheet: A Statistical Study

J. M. H. Beedle<sup>1,2</sup>, D. J. Gershman<sup>2</sup>, V. M. Uritsky<sup>1,2</sup>, T. D. Phan<sup>3</sup>, B. L. Giles<sup>2</sup>

<sup>1</sup>Department of Physics, The Catholic University of America, Washington D.C. USA

<sup>2</sup>NASA Goddard Space Flight Center, Greenbelt, MD, USA

<sup>3</sup>Space Sciences Laboratory, University of California, Berkeley, CA, USA

## Key Points:

- EDR crossings show current densities an order of magnitude higher than non-EDR magnetopause crossings.
- EDR crossings contain significant current components parallel to the local magnetic field.
- Ion velocity along the magnetopause is highly correlated with an event's location - indicating the presence of magnetosheath flows.

---

Corresponding author: Jason M. H. Beedle, [beedle@cua.edu](mailto:beedle@cua.edu)

## Abstract

The solar wind is a continuous outflow of charged particles from the Sun's atmosphere into the solar system. At Earth, the solar wind's outward pressure is balanced by the Earth's magnetic field in a boundary layer known as the magnetopause. Plasma density and temperature differences across the boundary layer generate the Chapman-Ferraro current which supports the magnetopause. Along the dayside magnetopause, magnetic reconnection can occur in electron diffusion regions (EDRs) embedded into the larger ion diffusion regions (IDRs). These diffusion regions form when opposing magnetic field lines in the solar wind and Earth's magnetic field merge, releasing magnetic energy into the surrounding plasma. While previous studies have given us a general understanding of the structure of these diffusion regions, we still do not have a good grasp of how they are statistically differentiated from the non-diffusion region magnetopause. By investigating 251 magnetopause crossings from NASA's Magnetospheric Multiscale (MMS) Mission, we demonstrate that EDR magnetopause crossings show current densities an order of magnitude higher than non-EDR magnetopause crossings - crossings that either passed through the reconnection exhausts or through the non-reconnecting magnetopause. Significant current signatures parallel to the local magnetic field in EDR crossings are also identified, which is in contrast to the dominantly perpendicular current found in the non-EDR magnetopause. Additionally, we show that the ion velocity along the magnetopause is highly correlated with a crossing's location, indicating the presence of magnetosheath flows inside the magnetopause current sheet.

## Plain Language Summary

The magnetopause is a dynamic boundary layer created through the interaction of the solar wind with Earth's magnetic field. This boundary is supported by a current sheet and acts as the "entry gate" of the solar wind's energy into the magnetosphere through a process called magnetic reconnection where energy previously stored in the magnetic field is released into the surrounding magnetopause plasma. The reconnection process is initiated in localized diffusion regions, which form in the magnetopause's current sheet during specific solar wind conditions. In this paper, we clarify what makes the diffusion regions stand out from the background magnetopause current sheet by utilizing data from NASA's Magnetospheric Multiscale mission. Our analysis reveals that the diffusion regions have stronger currents than the background magnetopause and that a significant portion of this current becomes parallel to the local magnetic field.

## 1 Introduction

The magnetopause is a boundary layer created through the balancing of the solar wind's dynamic pressure with Earth's magnetic field. Across this boundary layer, pressure gradients generate a current sheet, named the Chapman-Ferraro (CF) current, that supports the magnetopause (e.g. Chapman and Ferraro (1931)). This current sheet is a large scale, mainly ion driven current generated from ion density and temperature gradients - e.g. Beedle et al. (2022) and references therein.

Along the dayside magnetopause current sheet, magnetic reconnection occurs when opposing field lines in the solar wind and Earth's magnetic field are driven together by plasma flows, causing the magnetopause boundary to thin. As the current sheet compresses, there becomes a small-scale region where the frozen-in condition in the plasma is violated, allowing the magnetic field to become disassociated from the plasma and diffuse through it, break, and then reform, changing the local magnetic topology (e.g. Vasyliunas (1975); Hesse and Cassak (2020)). This process occurs in what are known as diffusion regions. Specifically, there are two distinct regions: an ion diffusion region (IDR) and an electron diffusion region (EDR). The IDR is the larger outer region where ions first

dissociate from the magnetic field while the electrons remain frozen-in. This hybrid configuration with magnetized electrons and free ions then creates the Hall currents and their associated quadrupole Hall magnetic field in the IDR (e.g. Sonnerup (1979); Oieroset et al. (2001); Mozer et al. (2002)). The EDR is the smaller inner diffusion region, embedded in the larger IDR, where both the electrons and ions are decoupled from the magnetic field, which allows magnetic reconnection to take place - e.g. Vasyliunas (1975), Burch et al. (2016).

The process of magnetic reconnection leads to the magnetopause acting as the "entry gate" of the solar wind's energy into the Earth's magnetosphere. Thus a thorough grasp of this process and its impact on the magnetopause's current sheet is vital to understanding the energy transfer into the terrestrial space weather system. Because of this significance, numerous missions (ISEE, AMPTE, Polar, Cluster, Themis, MMS, etc) devoted their resources to gaining insights into the magnetopause current sheet and dayside magnetopause reconnection. A number of studies [Burch et al. (2016), Lavraud et al. (2016), Norgren et al. (2016), Phan et al. (2016), Burch and Phan (2016), Chen et al. (2016), Chen et al. (2017), etc.] have focused on individual dayside magnetopause EDR events using the MMS spacecraft to study common elements of EDR crossings including ion jets and jet reversals around the diffusion regions, plasma inflows, non-gyrotropic crescent shaped electron outflows, intense currents, and strong heating. Other studies [Rager et al. (2018), Webster et al. (2018), Shuster et al. (2019), Genestreti et al. (2020), Shuster et al. (2021), etc.] have addressed these generalized characteristics of EDR events in more detail and confirmed the prevalence of crescent-shaped electron velocity phase space densities, ohmic heating of the plasma, as well as the role of electron scale currents to the EDR regions.

While previous studies provide a general understanding of the diffusion regions' structure, there has not yet been a statistical study of the characteristic differences between magnetopause crossings with and without active signatures of reconnection. To begin answering this question, our study analyzes data from NASA's MMS Mission during EDR and non-EDR magnetopause crossings where these non-EDR events could either be encounters of the reconnection exhausts downstream of the diffusion region, or non-reconnecting magnetopause crossings. In Section 2, we describe the methods we used to accomplish this analysis as well as the findings from comparing the EDR and non-EDR magnetopause crossings. Section 3 covers an in-depth analysis of our results, with observations about the magnetopause's current structure and ion velocities measured during these EDR events. Section 4 introduces a brief discussion of our findings, and Section 5 provides a summary of the main results of this study.

## 2 Observations

### 2.1 MMS Data and Current Calculations

For this study, we utilized data from NASA's MMS mission, which is a mission comprised of four spacecraft that travel in a tetrahedron pattern through the magnetosphere. MMS's Fast Plasma Investigation (FPI) (Pollock et al., 2016) and Fluxgate Magnetometer (Russell et al., 2016) instruments enable four simultaneous measurements of plasma properties and magnetic field conditions, respectively, across MMS's constellation. Using the data from these two instruments, we analyzed the magnetopause current system through the following currents.

The first is called the curlometer current, or  $\mathbf{J}_{curl}$ , which was calculated using Dunlop et al. (1988)'s curlometer method to approximate gradients in the magnetic field, yielding Ampere's law in the MHD approximation:

$$\mathbf{J}_{curl} = \frac{\nabla \times \mathbf{B}}{\mu_0}, \quad (1)$$

where  $\mathbf{B}$  is the magnetic field and  $\mu_0$  is the permeability of free space.  $\mathbf{J}_{curl}$  represents the current consistent with magnetic field perturbations and is thus a proxy for the total current density encountered by MMS during a magnetopause current sheet crossing as it contains current components parallel and perpendicular to the magnetic field including both ion and electron contributions. While  $\mathbf{J}_{curl}$  has been found to be less sensitive to small current structures than the current density calculated from plasma moments, also known as the FPI current, we decided to focus on using  $\mathbf{J}_{curl}$  as the average MMS separation during our time frame (2015–2018) was between 10 to 60 kms, which is sufficiently close to consider the ion dominated CF current as well as any large scale currents in and around the diffusion regions in EDR events. This is in contrast to studies that have focused more heavily on the smaller scale, electron dominated currents of the EDR, which generally use the FPI current (e.g. Lavraud et al. (2016); Phan et al. (2016)). We also considered the components of  $\mathbf{J}_{curl}$  parallel and perpendicular to the locally measured magnetic field  $\mathbf{B}$ :  $\mathbf{J}_{curl\parallel}$  and  $\mathbf{J}_{curl\perp}$ , defined as follows:

$$\mathbf{J}_{curl\parallel} = \left( \frac{\mathbf{B} \cdot \mathbf{J}_{curl}}{|\mathbf{B}|} \right) \hat{\mathbf{B}} \quad , \quad \mathbf{J}_{curl\perp} = \mathbf{J}_{curl} - \mathbf{J}_{curl\parallel}. \quad (2)$$

Note,  $\mathbf{B}$  is averaged across all four MMS spacecraft to match the curlometer method calculations.

Along with  $\mathbf{J}_{curl}$ , we looked at the ion and electron diamagnetic currents:  $\mathbf{J}_{dia\ Total_i}$  and  $\mathbf{J}_{dia\ Total_e}$  and their current components generated from temperature and density gradients, which were approximated using the curlometer method. Both the ion and electron diamagnetic currents and their density and temperature components were defined in the following manner, with their respective densities and temperatures, in the same way as Beedle et al. (2022):

$$\mathbf{J}_{dia\ \nabla N} = \frac{\mathbf{B} \times (k_b \overleftrightarrow{T} \cdot \nabla N)}{|\mathbf{B}|^2} \quad , \quad \mathbf{J}_{dia\ \nabla \cdot \overleftrightarrow{T}} = \frac{\mathbf{B} \times (k_b N \nabla \cdot \overleftrightarrow{T})}{|\mathbf{B}|^2}, \quad (3)$$

where  $\mathbf{B}$  represents the magnetic field,  $k_b$  is Boltzmann's constant,  $\overleftrightarrow{T}$  is the temperature tensor, and  $N$  is the number density. By definition,  $\mathbf{J}_{dia\ Total} = \mathbf{J}_{dia\ \nabla N} + \mathbf{J}_{dia\ \nabla \cdot \overleftrightarrow{T}}$ . Note,  $\mathbf{B}$ ,  $\overleftrightarrow{T}$ , and  $N$  were averaged across all four MMS spacecraft to match with the use of the curlometer method to calculate the gradients. When referencing these components, we will refer to  $\mathbf{J}_{dia\ \nabla N}$  as its current's density component and to  $\mathbf{J}_{dia\ \nabla \cdot \overleftrightarrow{T}}$  as its current's temperature component.

While we considered both parallel and perpendicular components for  $\mathbf{J}_{curl}$ , diamagnetic current is, by definition, perpendicular to the magnetic field, thus the ion and electron diamagnetic currents represent the primary perpendicular components to the magnetic field in the magnetopause.

To summarize, we analyzed the following set of current densities:  $\mathbf{J}_{curl}$ ,  $\mathbf{J}_{dia\ Total_i}$ ,  $\mathbf{J}_{dia\ Total_e}$ ,  $\mathbf{J}_{dia\ \nabla N_i}$ ,  $\mathbf{J}_{dia\ \nabla \cdot T_i}$ ,  $\mathbf{J}_{dia\ \nabla N_e}$ ,  $\mathbf{J}_{dia\ \nabla \cdot T_e}$  during each of the studied magnetopause crossings.

When interpreting these quantities, it is important to note that MHD physics breaks down in the diffusion regions as plasma disassociates from the magnetic field. Because the diamagnetic current equations are defined under MHD conditions, the concept of a diamagnetic current also breaks in the diffusion regions as the plasma must now be described using kinetic theories. Thus, while the current around the diffusion region is still


represented by the diamagnetic, ion dominated CF current, inside the diffusion regions they become kinetic and can no longer be described in the same way. For this reason our results using the diamagnetic current are more likely to contain anomalously large current spikes once the MMS constellation entered the IDR and EDR of that magnetopause crossing event. However, as the EDR itself is still quite small when compared with the current sheet that MMS observes, there are regions where the diamagnetic current is a useful measure. It is also worthy to note that, as the curlometer current relies on the deviations in the magnetic field itself, it is not impacted in the same manner and presents accurate current measurements all throughout the current sheet crossing, be it in the current sheet itself, or the diffusion regions.

Data taken from MMS, as well as the calculated currents, were interpolated to the 30 ms FPI electron time resolution. As our main analysis involves ion and electron diamagnetic currents and the total current as computed from the curlometer method, any sub 150 ms variations in the ion parameters should not impact our results. For all measured quantities that did not use the curlometer method, we averaged over all four MMS spacecraft to create a single data stream. Our calculations and measurements were completed in Cartesian GSE coordinates and then stored in spherical GSE coordinates, in which the  $\phi$  angle is in the primary current direction along the dayside magnetopause as can be seen defined in Figure 1. A more detailed description of these spherical coordinates is provided in Section 3.


## 2.2 Event Selection

To select relevant data for our study, we used EDR crossings provided by Webster et al. (2018), who compiled previously identified EDR events with a set of newly-identified EDR events based on shared characteristics including the occurrence of non-gyrotropic crescent-shape electron distributions, ion jet reversals, and large current densities. Because of this reliance on non-gyrotropic electron distributions to identify EDR events, the Webster et al. (2018) events can only include, at most, a moderate guide field as stronger guide fields tend to obscure this feature. In all, Webster et al. (2018) reported 32 EDR events, 26 of which were included in our study based on their location along the dayside magnetopause as well as the availability of MMS data from all four spacecraft. Four of Webster et al.'s events (A13, B14, B15, and B17) were located outside of the bounds of our definition of the dayside magnetopause (see Figure 1), while two other events (A7 and B26) caused errors with our code because of data outages from one or more MMS spacecraft. The selected 26 EDR events then represented the EDR sample group that we measured the aforementioned current densities and other plasma characteristics over. The locations of these events along the dayside magnetopause are denoted in blue in Figure 1.

Along with these 26 EDR events, we also investigated 225 dayside magnetopause current sheet crossings taken from Paschmann et al. (2018) and Haaland et al. (2020)'s database of MMS magnetopause crossings. These 225 events were previously used in the Beedle et al. (2022) study and include complete, monotonic magnetopause crossings and, to our knowledge, do not include any previously identified EDR events. Thus, these 225 events represent our non-EDR magnetopause crossing sample group that we then compared with the EDR samples. As previously mentioned, these non-EDR crossings could either be a crossing of the reconnection exhausts downstream of the diffusion regions, or a crossing of the non-reconnection magnetopause. Beedle et al. (2022) provides a detailed explanation of the selection criteria for these 225 events. The locations of the 225 magnetopause crossing events are denoted in red in Figure 1. An example of a non-EDR event versus an EDR event is provided in Figure 2.



**Figure 1.** Diagram of our 225 dayside non-EDR (pink) and 26 EDR (blue) magnetopause crossings. We define a spherical coordinate system with  $\phi$  in the  $X_{GSE} - Y_{GSE}$  plane, positively defined from the  $+X_{GSE}$  axis,  $R$  defined as radially outward, and  $\theta$  as the polar angle into the  $+Z_{GSE}$  direction, completing the right-handed coordinate system. The Dayside is defined as being from  $+50^\circ$  to  $-50^\circ$  in  $\phi$ , following the same convention as Beedle et al. (2022).



**Figure 2.** Example magnetopause crossings representing a non-EDR crossing (left) and an EDR event (right). The orange dashed lines represent the magnetopause crossing as identified by our algorithm for each event (see Section 2.3). The example EDR event is from Burch et al. 2016. (a) Magnitude and magnetic field in LMN coordinates determined through MVAB analysis (Sonnerup & Scheible, 1998), (b) and (c) ion and electron omni directional spectrograms, (d) ion number density, (e) ion perpendicular and parallel temperature, (f) ion velocity, (g, h, i, j and k) curlometer, total ion diamagnetic, ion density component, ion temperature component, and total electron diamagnetic current densities respectively in LMN coordinates with magnitudes indicated in black.



### 2.3 Magnetopause Identification

Each of the events in our database was processed by an algorithm to identify their magnetopause crossing times. We used  $\mathbf{J}_{curl}$  to identify the largest current magnitude peak during an event, and then applied a threshold equal to 20% of this peak value to the current density measured during the crossing. This separated the event into current segments, with each weighted based on their duration, average  $|\mathbf{J}_{curl}|$  current density, and the magnetic field magnitude,  $|\mathbf{B}|$ , measured over the segment. The segment with the longest duration, highest average current density, and largest change in  $|\mathbf{B}|$  across the segment was then selected as the primary current sheet crossing for that event, with the start and end times of the current segment then becoming the beginning and end of that event's magnetopause crossing. Two examples of the algorithm's selection method can be seen represented by the vertical orange dashed lines in Figure 2. Note, as this method uses the magnitudes of each value, it is coordinate system invariant.

This method was applied to both the EDR and non-EDR crossings, with the results for the average current density for the 225 non-EDR events matching within error the average current over the magnetopause crossing times identified by the Paschmann et al. (2018) database's minimum variance analysis method as previously reported in Beedle et al. (2022). The performance of our algorithm was also double checked over the 26 EDR events so that the selected magnetopause crossing correctly captured the EDR event as previously identified by their respective papers.

### 3 EDR and Non-EDR Crossing Analysis

Over each of the 26 EDR events and 225 non-EDR magnetopause crossings, we recorded individual current density data and stored the results in spherical GSE coordinates with  $R$  being defined as radially outward,  $\phi$  going from dawn-to-dusk in the  $X_{GSE}$ - $Y_{GSE}$  plane, and  $\theta$  pointing in the  $+Z_{GSE}$  direction (see Figure 1 for a visual depiction). We utilize spherical GSE coordinates instead of LMN coordinates for our statistical survey to be able to compare current density components measured over the EDR and non-EDR events on an equal footing. In previous statistical studies (e.g. Paschmann et al. (2018), Haaland et al. (2020)), MVAB analyses (Sonnerup & Scheible, 1998) were utilized over MMS's burst mode intervals to generate LMN coordinates for their events. This works well for intervals that involve a single, clear magnetopause crossing, but leads to uncertainties when MMS passes over the magnetopause multiple times in quick succession, such as during active solar wind conditions. These crossings are often nonuniform and contain small scale embedded structure. In such cases, the MVAB analysis interval needs to be adjusted in order to capture the appropriate crossing, which leads to some ambiguity, especially when trying to compare the individual current directions measured over many such events. In the aforementioned studies, the magnitude of the current density was reported for each event, which is unaffected by these differences in coordinate determination. For our statistical survey, as we directly compare currents along coordinate directions, we decided to use a global coordinate system that is equally applied to all of our crossings, regardless of the dynamics involved.


Our analysis resulted in 6,332 data points for each current component from the EDR events and 73,865 data points from the non-EDR events. We then analyzed the combined data's mean and median values as well as their standard errors or  $\sigma/\sqrt{N}$  where  $\sigma$  is the standard deviation of the data and  $N$  is the number of data points recorded. This analysis was completed for each of our currents densities ( $\mathbf{J}_{curl}$ ,  $\mathbf{J}_{dia\ Total_i}$ ,  $\mathbf{J}_{dia\ Total_e}$ ,  $\mathbf{J}_{dia\ \nabla N_i}$ ,  $\mathbf{J}_{dia\ \nabla T_i}$ ,  $\mathbf{J}_{dia\ \nabla N_e}$ ,  $\mathbf{J}_{dia\ \nabla T_e}$ ) in their component directions ( $\hat{R}$ ,  $\hat{\phi}$ ,  $\hat{\theta}$ ), as shown in Table 1 below. Additionally, we compiled the current data into probability distributions, which are shown in Figure 3 for  $J_{curl}$  and its parallel and perpendicular components in the  $\hat{R}$ ,  $\hat{\phi}$ ,  $\hat{\theta}$  directions (9 panels in total) with the EDR data points represented in the blue distributions, while the non-EDR data points are represented in the pink distribu-




tions. Likewise, Figure 4 shows the results for the ion diamagnetic current and its density and temperature components. The electron diamagnetic current and its components are provided in Figure 5. Figure 6 then shows the probability density histograms of ion and electron velocity measurements over the EDR and non-EDR crossings. Each distribution figure includes labels that show the total number of points  $N$  as well as the mean and median values of their respective distributions, which are also shown in Table 1.

**Table 1.** Comparison of current densities obtained during the 225 non-EDR and 26 EDR crossings with the following format: mean (median)  $\pm$  standard error of the current densities as measured in spherical GSE coordinates ( $\hat{R}$ ,  $\hat{\phi}$ ,  $\hat{\theta}$ ). The mean and median values are computed and presented in the same way as those shown in Figures 3 - 5. The EDR/Non-EDR ratio was also computed based on these mean and median values.


Current	Non-EDR ( $nA/m^2$ )	EDR ( $nA/m^2$ )	EDR / Non-EDR
$J_R \text{ curl}$	4.96 (2.40) $\pm$ 0.39	30.1 (18.5) $\pm$ 2.53	6.1 (7.7)
$J_R \text{ curl}_\perp$	-0.04 (-0.60) $\pm$ 0.33	14.1 (-1.70) $\pm$ 2.19	350 (2.8)
$J_R \text{ curl}_\parallel$	5.00 (1.00) $\pm$ 0.20	16.0 (6.70) $\pm$ 1.59	3.2 (6.7)
$J_R \text{ dia Total}_i$	-1.21 (-0.70) $\pm$ 0.70	-26.6 (-23.1) $\pm$ 3.54	22 (33)
$J_R \text{ dia } \nabla N_i$	-7.19 (-1.20) $\pm$ 0.71	-0.88 (-18.4) $\pm$ 2.77	0.12 (15)
$J_R \text{ dia } \nabla \cdot T_i$	5.98 (-0.30) $\pm$ 0.71	-25.7 (-5.00) $\pm$ 3.15	4.3 (17)
$J_R \text{ dia Total}_e$	-2.23 (-1.00) $\pm$ 0.10	13.2 (2.20) $\pm$ 0.91	5.9 (2.2)
$J_R \text{ dia } \nabla N_e$	-2.41 (-1.00) $\pm$ 0.08	4.99 (-0.10) $\pm$ 0.48	2.1 (0.1)
$J_R \text{ dia } \nabla \cdot T_e$	0.18 (0.00) $\pm$ 0.07	8.25 (1.50) $\pm$ 0.74	49 (NA)
$J_\phi \text{ curl}$	89.5 (68.4) $\pm$ 0.52	349 (324) $\pm$ 3.60	3.9 (4.7)
$J_\phi \text{ curl}_\perp$	56.8 (42.6) $\pm$ 0.36	201 (161) $\pm$ 3.06	3.5 (3.8)
$J_\phi \text{ curl}_\parallel$	32.7 (7.4) $\pm$ 0.39	148 (85.3) $\pm$ 2.69	4.5 (11.5)
$J_\phi \text{ dia Total}_i$	57.6 (44.4) $\pm$ 0.67	210 (129) $\pm$ 4.93	3.6 (2.9)
$J_\phi \text{ dia } \nabla N_i$	80.6 (60.8) $\pm$ 0.65	252 (136) $\pm$ 5.59	3.1 (2.2)
$J_\phi \text{ dia } \nabla \cdot T_i$	-23.0 (-15.6) $\pm$ 0.63	-42.5 (-28.7) $\pm$ 3.69	1.8 (1.8)
$J_\phi \text{ dia Total}_e$	6.24 (3.80) $\pm$ 0.11	28.1 (20.0) $\pm$ 1.08	4.5 (5.3)
$J_\phi \text{ dia } \nabla N_e$	5.28 (3.80) $\pm$ 0.08	18.3 (12.3) $\pm$ 0.60	3.5 (3.2)
$J_\phi \text{ dia } \nabla \cdot T_e$	0.96 (0.00) $\pm$ 0.06	9.82 (4.70) $\pm$ 0.86	10 (NA)
$J_\theta \text{ curl}$	8.90 (9.50) $\pm$ 0.57	-38.0 (-33.1) $\pm$ 4.15	4.3 (4.1)
$J_\theta \text{ curl}_\perp$	1.99 (1.80) $\pm$ 0.28	-59.4 (-45.6) $\pm$ 2.43	30 (25)
$J_\theta \text{ curl}_\parallel$	6.92 (5.30) $\pm$ 0.53	21.4 (28.0) $\pm$ 3.98	3.1 (5.3)
$J_\theta \text{ dia Total}_i$	-6.13 (1.70) $\pm$ 0.63	-31.1 (-16.4) $\pm$ 4.57	5.1 (9.6)
$J_\theta \text{ dia } \nabla N_i$	-0.85 (2.60) $\pm$ 0.55	-24.0 (-22.8) $\pm$ 4.16	28 (8.8)
$J_\theta \text{ dia } \nabla \cdot T_i$	-5.28 (-1.70) $\pm$ 0.64	-7.08 (17.0) $\pm$ 4.30	1.3 (10)
$J_\theta \text{ dia Total}_e$	1.94 (0.50) $\pm$ 0.10	-13.2 (-6.80) $\pm$ 0.92	6.8 (14)
$J_\theta \text{ dia } \nabla N_e$	1.48 (0.40) $\pm$ 0.07	-10.8 (-5.60) $\pm$ 0.48	7.3 (14)
$J_\theta \text{ dia } \nabla \cdot T_e$	0.46 (0.00) $\pm$ 0.06	-2.34 (-1.80) $\pm$ 0.70	5.1 (NA)




**Figure 3.** Probability distribution histograms of the curlometer current and its parallel and perpendicular components to the local magnetic field from the 26 EDR crossings (blue) and 225 non-EDR magnetopause crossings (pink) measured across the three global coordinates ( $\hat{R}, \hat{\phi}, \hat{\theta}$ ). The EDR events gave us 6,332 data points in total, while the non-EDR crossings gave us 73,865 data points. Note that the vertical axis in each plot is normalized, with the same scale used for each subplot for the vertical and horizontal axes respectively. The bins used are also the same for each subplot's distributions. The sample mean and median values are provided in the top right of each subplot.



**Figure 4.** Probability distribution histograms of the ion diamagnetic current and its current components - the density component and the temperature component - in EDR crossings (blue) and non-EDR crossings (pink) over the spherical  $\hat{R}$ ,  $\hat{\phi}$ , and  $\hat{\theta}$  component directions. The sample mean and median values are provided in the top right of each subplot.



**Figure 5.** Probability distribution histograms of the electron diamagnetic current and its current components - the density component and the temperature component - in EDR crossings (blue) and non-EDR crossings (pink) over the spherical  $\hat{R}$ ,  $\hat{\phi}$ , and  $\hat{\theta}$  component directions. The sample mean and median values are provided in the top right of each subplot.



**Figure 6.** Ion and electron velocity histograms from the 26 EDR crossings (blue) and the 225 non-EDR magnetopause crossings (pink). Note the double peak structure of the EDR ion velocities in the  $\hat{\phi}$  direction as well as the multi-peak structure in the  $\hat{\theta}$  direction.

### 3.1 EDR vs Non-EDR Current Structure

Using the non-EDR events as a baseline for the background CF current sheet in the magnetopause, we can make the following observations about the magnetopause's current structure around and during EDR events from Table 1 and Figures 3 - 6:

1. EDR current densities along the primary,  $\hat{\phi}$ , magnetopause direction are an order of magnitude higher, on average, than a non-EDR crossing.

$\mathbf{J}_{curl}$ ,  $\mathbf{J}_{dia\ Total_i}$ , and  $\mathbf{J}_{dia\ Total_e}$  are all larger by an order of magnitude in the  $\hat{\phi}$  direction. This applies for both the mean and median values of the measured current densities, implying that this enhancement does not just affect our data's outlying points. The amplified current density matches with general expectations of EDR crossings having strong currents because of the EDR's thin, electron-scale current sheets (see e.g. Webster et al. (2018)). There is, however, one interesting outlier to this conclusion;  $\mathbf{J}_{dia\ \nabla \cdot T_i}$ . Not only does the temperature-generated current density,  $\mathbf{J}_{dia\ \nabla \cdot T_i}$ , fail to show this order of magnitude jump, but it actually exhibits the smallest increase of all the average currents, in the  $\hat{\phi}$  direction, with a 1.8x increase from its non-EDR counterpart. This suggests that, while the density-generated current density,  $\mathbf{J}_{dia\ \nabla N_i}$ , does see a substantial boost during these EDR events,  $\mathbf{J}_{dia\ \nabla \cdot T_i}$  does not show a similar reaction. Overall indicating that ions do not see the same level of heating inside of the diffusion regions, and perhaps indicating that ions are largely unaffected by the electron-scale dynamics in the EDR.

2. There are significant  $\hat{\phi}$  directed current signatures parallel to the local magnetic field during EDR events.

$\mathbf{J}_{curl}$  parallel to the local magnetic field along the primary  $\hat{\phi}$  direction becomes significantly enhanced during EDR events. This can be seen in Figure 3 and Table 1, with the parallel component's mean value increasing by 4.5x and its median value increasing by 11.5x. While there is a significant enhancement in the  $\hat{\phi}$  perpendicular current density as well, the perpendicular component's mean value increases by 3.5x and its median by only 3.8x, noticeably less than the parallel component. This difference between the parallel and perpendicular components leads to the overall amount of current parallel to the magnetic field in the primary  $\hat{\phi}$  direction (along the CF current's flow) to increase in EDR events. Specifically, the mean parallel current density accounts for 42% of the mean curlometer current density in EDR events, up slightly from 36% in the non-EDR events. However, looking at the more impacted median values, the median parallel current density accounts for approximately 26% of the median curlometer current density in EDR events, over twice as much as the 11% contribution seen in non-EDR events. This indicates that a large percentage of formerly perpendicular current density - the CF current - in the non-EDR magnetopause becomes parallel to the local magnetic field during EDR events.

3. The ion diamagnetic current density dominates that of the electron current density, but to a lesser extent in EDR events.

From Table 1 and Figures 4 and 5, both non-EDR and EDR events show that the average  $\mathbf{J}_{dia\ Total_i}$  is greater than that of the average  $\mathbf{J}_{dia\ Total_e}$ . Specifically, in non-EDR crossings, the average  $\mathbf{J}_{dia\ Total_i}$  is 9.2x larger than  $\mathbf{J}_{dia\ Total_e}$ . This matches with findings from Beedle et al. (2022) where the non-EDR magnetopause current was found to be ion dominated. During EDR crossings, we still find that the ions dominate, but to a lesser degree. Looking at the average values from Table 1, one can see that, during EDR crossings,  $\mathbf{J}_{dia\ Total_i}$  is 7.5x larger than  $\mathbf{J}_{dia\ Total_e}$ . So, while the ions are still the main contributors, their contribution seems to decrease - primarily because of a stronger electron response in EDR events. Specifically,  $\mathbf{J}_{dia\ Total_e}$  sees a 4.5x average increase in EDR crossings when compared with their non-EDR counterparts, while  $\mathbf{J}_{dia\ Total_i}$  sees a lesser 3.6x average increase. The increasing importance of  $\mathbf{J}_{dia\ Total_e}$  during EDR events matches with the general expectations of an EDR crossing where the electron diffusion region itself is known to be dominated by electron scale currents (e.g. Shuster et al. (2019)). However, our results imply that, while the central electron diffusion region is dominated by these electron currents, the CF current in the magnetopause current sheet itself is still primarily ion dominated.

4.  $\mathbf{J}_{dia\ Total_e}$  is composed of temperature and density components that work together instead of destructively like  $\mathbf{J}_{dia\ Total_i}$ 's components. The enhanced  $\mathbf{J}_{dia\ Total_e}$  found in an EDR event comes primarily from an increase in the temperature component, whose relative contribution increases by an order of magnitude.

Figure 5 and Table 1 show that, in both EDR as well as non-EDR events, the electron temperature and density components work with one other in the  $+\hat{\phi}$  direction. While this is true in both types of crossings, it is significantly more pronounced in EDR events. Additionally, the contribution of  $\mathbf{J}_{dia\ \nabla \cdot \vec{T}_e}$  is measurably enhanced in EDR crossings with a 10x increase in its average value seen in Table 1 as compared to the 3.5x increase in  $\mathbf{J}_{dia\ \nabla N_e}$ . This impressive enhancement to  $\mathbf{J}_{dia\ \nabla \cdot \vec{T}_e}$  is likely a result of electron heating, leading to the formation of a strong electron temperature divergence in the diffusion region. The presence of electron heating has been noted as a key component to providing pressure balance in the EDR (Hesse



& Cassak, 2020). Overall, the  $\phi$  enhancements to both components leads to  $\mathbf{J}_{dia\ Total_e}$ 's net strength increasing by 4.5x.

5. The EDR's ion velocity measurements are characterized by multi-peak probability distributions, while the non-EDR events are described by single peak distributions.

From Figure 6, we can clearly see that the EDR ion velocity in the  $\hat{\phi}$  and  $\hat{\theta}$  directions has multi-peak distributions, which is in contrast with the single peak distributions of the non-EDR events.

Regarding the  $\hat{\theta}$  ion distribution, there is a multi-peak distribution in the data, which likely forms from ion outflows jets in the IDR with the jets extending out from the reconnection site along  $\hat{L}$  in LMN coordinates or along  $\hat{\theta}$  in our spherical coordinates. As MMS flies through the magnetopause, it can encounter both sides of the jets, forming the positive and negative peaks, or the center of the reconnection site, where there is little to no ion movement, forming the central peak near zero.

There is also a clear dual peak in the EDR events'  $V_{i\phi}$  velocity distribution. Interestingly, this double peak can be explained by considering magnetosheath flows around the dayside magnetopause. On the dusk-side of the subsolar point, the magnetosheath plasma flows in the  $+\hat{\phi}$  direction, while on the dawn-side, the sheath plasma flows in the  $-\hat{\phi}$  around the magnetopause. We have found that EDR events on the dusk side of the subsolar point tend to have average  $+V_{i\phi}$  flows across their MP crossings - accounting for the positive peak in Figure 6, while EDR events on the dawn side of the subsolar point tend to have average  $-V_{i\phi}$  flows - accounting for the negative peak in Figure 6. Thus, this matches with the expectation of the aforementioned magnetosheath flows. Performing a linear correlation analysis between the position of MMS along the dayside magnetopause with the average  $V_{i\phi}$  across the MP current sheets gives a correlation of 0.9, which shows how strongly correlated the EDR event's location is with the appearance of these magnetosheath flows. Interestingly, this correlation is even able to be seen in the non-EDR events as the linear correlation between MMS's location and average  $V_{i\phi}$  is 0.78.


Additionally, Figure 6 suggests that the electron velocity in the  $\hat{\phi}$  and  $\hat{R}$  directions tends to be higher than the ion velocity during EDR events. This indicates periods where the current sheet is primarily controlled by electron scale current structures as was previously observed (e.g. Phan et al. (2016)). During the non-EDR crossings, the electron velocity is generally smaller than the ion velocity, indicating that the non-EDR magnetopause current is dominated by the ion current, as previously reported in e.g. Beedle et al. (2022).

## 4 Discussion

### 4.1 Magnetosheath Flows in the Dayside Magnetopause

As stated above (Item 5, Section 3.1), magnetosheath flows dominate the ion velocity running along the magnetopause boundary, or  $V_{i\phi}$ , in both the non-EDR and EDR magnetopause current sheet as is illustrated in Figure 7. This suggests that sheath flows are primarily responsible for the ion velocity along this direction and overshadow the CF current ions in their dawn-to-dusk circulation around the dayside MP, revealing two relevant aspects of the magnetopause current system:

First, the magnetopause current sheet during both active and inactive solar wind conditions, is open to the influence of magnetosheath flows. This indicates that the cur-



**Figure 7.** Left: figure illustrating the sheath flows in the  $+$  and  $-\hat{\phi}$  directions around the subsolar point ( $\phi = 0^\circ$ ) of the magnetopause. Indicated in red is an example position of MMS on its orbit around the dayside magnetopause with its location in  $\phi$ . Right: two linear correlation diagrams of averaged  $V_{i\phi}$  over each MP crossing with MMS's  $\phi$  location along the magnetopause. EDR events are represented in the top diagram while non-EDR events are represented on the bottom. The correlation coefficient of the plotted linear fit (shown in red) is provided in the lower right-hand side of each plot.

rent sheet, even while retaining the structure and flow mechanics of the CF current, is dominated by faster flowing sheath ions, which changes the observed  $V_{i\phi}$  flow. While less correlated, the electron velocity along the magnetopause also seems to be correlated with position, with average  $V_{e\phi}$  for EDR events having a correlation of 0.51 and average  $V_{e\phi}$  for non-EDR events having a correlation of 0.67. As both ion and electron velocities are correlated with position along the magnetopause, this means that these magnetosheath flows are likely bulk flows and should not impact the current structure of the magnetopause itself.

Second, if we directly consider the  $\hat{\theta}$  direction flows, both  $V_{i\theta}$  and  $V_{e\theta}$  do not show any correlation with MMS's  $\phi$  position along the magnetopause. However, we can also consider MMS's location relative to the  $X_{GSE} - Y_{GSE}$  plane, or its  $\theta$  position angle. If we consider a similar correlation analysis with the ion and electron velocities versus MMS's  $\theta$  location, we see the following correlations. Both  $V_{i\theta}$  and  $V_{e\theta}$  for EDR events show a strong linear correlation with MMS's  $\theta$  position with -0.65 and -0.6 respectively. For non-EDR events,  $V_{i\theta}$  and  $V_{e\theta}$  show much lower correlations at -0.27 and -0.22 respectively. This shows the more open nature of the EDR event's magnetopause and also indicates the presence of sheath flows wrapping up and around the dayside magnetopause in the  $\hat{\theta}$  or  $Z_{GSE}$  direction.

## 4.2 Current Structure in EDR Events

As Items 1-4 of Section 3.1 suggest, EDR events depict a more complex and dynamic current structure than the non-EDR magnetopause. While this is generally expected because of the added complexity from filamentary electron-scale current sheets in the EDR (e.g. Phan et al. (2016), Shuster et al. (2019), Shuster et al. (2021)) and electron dominated Hall currents in the IDR (e.g. Sonnerup (1979), Nagai et al. (2001), Mozer et al. (2002)), there are findings that come as a surprise. The most prevalent of these is regarding the increased presence of parallel current in EDR events. Not only is this parallel current stronger than during the non-EDR magnetopause, but also represents an interesting counterpoint to the primarily perpendicular, ion dominated diamagnetic current seen in the background magnetopause current sheet (e.g. Beedle et al. (2022)). As the inner EDR is void of appreciable magnetic field components in the  $\hat{M}$  or  $\hat{\phi}$  direction (for low to no guide field cases), this  $\hat{\phi}$  parallel current indicates that a measurable and significant current in this direction is detected inside the outer IDR, becoming parallel to its  $\hat{M}$  directed Hall magnetic field. This could suggest additional current structure beyond the traditional 2.5D picture of the reconnection plane as is shown in zero-guide field PIC simulations such as those depicted in Shay et al. (2016) etc. These 2.5D structures typically show strong  $J_M$  generated by electron currents in the inner EDR, but whose strength diminishes inside the outer IDR where the Hall magnetic field aligns with its  $M$  direction. This thus predicts an overall weaker parallel current structure than what our data suggests. Further investigation of this parallel current signature's mechanism, and the role that the moderate guide fields in these Webster et al. (2018) events play, is needed however.

## 5 Summary and Conclusions

We used MMS magnetopause crossing data over 26 dayside EDR crossings and 225 non-EDR crossings to characterize differences between the diffusion regions and the background magnetopause current sheet. From this statistical analysis, we found the following:

- EDR crossings show current densities an order of magnitude higher than non-EDR magnetopause crossings, representing the significantly enhanced current sheet during EDR events.

- EDR crossings contain pronounced current components parallel to the local magnetic field. This is in contrast to the primarily perpendicular current density found in the non-EDR current sheet and suggests a large portion of the formerly perpendicular CF current in the non-EDR magnetopause becomes parallel to the local magnetic field during EDR events.
- EDR and non-EDR crossings both show average ion velocities that are highly correlated with a crossing's location along the magnetopause, indicating the presence of magnetosheath flows in the magnetopause current sheet. These flows tend to overshadow the CF current ions in their dawn-to-dusk circulation.

## 6 Open Research

The MMS data used in this study is publicly available at <https://lasp.colorado.edu/mms/sdc/public/dataset> from the FPI, and FIELDS datasets. The averaged MMS crossing data as well as the data used to create the histograms in Figures 3 - 6, from the 225 dayside magnetopause crossings and 26 EDR events, is available through a Harvard Dataverse public database: <https://doi.org/10.7910/DVN/UEDWO9>.

## Acknowledgments

The MMS current sheet database was created by Goetz Paschmann and Stein Haaland, and further developed by the International Space Science Institute Team 442, "Study of the physical processes in magnetopause and magnetosheath current sheets using a large MMS database". We thank them as well as the entire MMS team and instrument leads for the data access and support. We also thank Jim Drake, Li-Jen Chen, and Jason Shuster for our conversations regarding EDR simulations and EDR structure. We also thank the pySPEDAS team for their support and data analysis tools. This research was supported by the NASA Magnetospheric Multiscale Mission in association with NASA contract NNG04EB99C. J. M. H. B. and V. M. U. were supported through the cooperative agreement 80NSSC21M0180.

## References

- Beedle, J. M. H., Gershman, D. J., Uritsky, V. M., Phan, T. D., & Giles, B. L. (2022). A systematic look at the temperature gradient contribution to the dayside magnetopause current. *Geophysical Research Letters*, 49. doi: <https://doi.org/10.1029/2021GL097547>
- Burch, J. L., & Phan, T. D. (2016). Magnetic reconnection at the dayside magnetopause: Advances with mms. *Geophysical Research Letters*, 43(16), 8327-8338. Retrieved from <https://agupubs.onlinelibrary.wiley.com/doi/abs/10.1002/2016GL069787> doi: <https://doi.org/10.1002/2016GL069787>
- Burch, J. L., Torbert, R. B., Phan, T. D., Chen, L. J., Moore, T. E., Ergun, R. E., ... et al. (2016). Electron-scale measurements of magnetic reconnection in space. *Science*, 352. doi: 10.1126/science.aaf2939
- Chapman, S., & Ferraro, V. C. A. (1931). A new theory of magnetic storms. *Terrestrial Magnetism and Atmospheric Electricity*, 36, 77-97.
- Chen, L. J., Hesse, M., Wang, S., Gershman, D., Ergun, R., Pollock, C., ... et al. (2016). Electron energization and mixing observed by mms in the vicinity of an electron diffusion region during magnetopause reconnection. *Geophysical Research Letters*, 43, 6036-6043. doi: [doi:10.1002/2016GL069215](https://doi.org/10.1002/2016GL069215)
- Chen, L. J., Hesse, M., Wang, S., Gershman, D., Ergun, R. E., Burch, J., ... Avanov, L. (2017). Electron diffusion region during magnetopause reconnection with an intermediate guide field: Magnetospheric multiscale observations. *J.*

- Geophys. Res. Space Physics*, 112, 5235–5246. doi: doi:10.1002/2017JA024004
- Dunlop, M. W., Southwood, D. J., Glassmeier, K.-H., & Neubauer, F. M. (1988). Analysis of multipoint magnetometer data. *Advanced Space Research*, 8, 273–277.
- Genestreti, K. J., Liu, Y.-H., Phan, T.-D., Denton, R. E., Torbert, R. B., Burch, J. L., ... Eriksson, S. (2020). Multiscale coupling during magnetopause reconnection: Interface between the electron and ion diffusion regions. *Journal of Geophysical Research: Space Physics*, 125(10), e2020JA027985. Retrieved from <https://agupubs.onlinelibrary.wiley.com/doi/abs/10.1029/2020JA027985> (e2020JA027985 10.1029/2020JA027985) doi: <https://doi.org/10.1029/2020JA027985>
- Haaland, S., Paschmann, G., Øieroset, M., Phan, T., Hasegawa, H., Fuselier, S. A., ... Burch, J. (2020). Characteristics of the flank magnetopause: Mms results. *Journal of Geophysical Research: Space Physics*, 125. doi: <https://doi.org/10.1029/2019JA027623>
- Hesse, M., & Cassak, P. A. (2020). Magnetic reconnection in the space sciences: Past, present, and future. *Journal of Geophysical Research: Space Physics*, 125. doi: <https://doi.org/10.1029/2018JA025935>
- Lavraud, B., Zhang, Y. C., Vernisse, Y., Gershman, D. J., Dorelli, J., Cassak, P. A., ... Yokota, S. (2016). Currents and associated electron scattering and bouncing near the diffusion region at earth's magnetopause. *Geophysical Research Letters*, 43(7), 3042–3050. Retrieved from <https://agupubs.onlinelibrary.wiley.com/doi/abs/10.1002/2016GL068359> doi: <https://doi.org/10.1002/2016GL068359>
- Mozer, F. S., Bale, S. D., & Phan, T. D. (2002, Jun). Evidence of diffusion regions at a subsolar magnetopause crossing. *Phys. Rev. Lett.*, 89, 015002. Retrieved from <https://link.aps.org/doi/10.1103/PhysRevLett.89.015002> doi: 10.1103/PhysRevLett.89.015002
- Nagai, T., Shinohara, I., Fujimoto, M., Hoshino, M., Saito, Y., Machida, S., & Mukai, T. (2001). Geotail observations of the hall current system: Evidence of magnetic reconnection in the magnetotail. *Journal of Geophysical Research: Space Physics*, 106(A11), 25929–25949. Retrieved from <https://agupubs.onlinelibrary.wiley.com/doi/abs/10.1029/2001JA900038> doi: <https://doi.org/10.1029/2001JA900038>
- Norgren, C., Graham, D. B., Khotyaintsev, Y. V., André, M., Vaivads, A., Chen, L. J., ... Burch, J. L. (2016). Finite gyroradius effects in the electron outflow of asymmetric magnetic reconnection. *Geophysical Research Letters*, 43, 6724–6733. doi: doi:10.1002/2016GL069205
- Øieroset, M., Phan, T. D., Fujimoto, M., Lin, R. P., & Lepping, R. P. (2001). In situ detection of collisionless reconnection in the earth's magnetotail. *Nature*, 412, 414–417. doi: <https://doi.org/10.1038/35086520>
- Paschmann, G., Haaland, S. E., Phan, T., Sonnerup, B., Burch, J., Torbert, R., ... Fuselier, S. (2018). Large-scale survey of the structure of the dayside magnetopause by mms. *Journal of Geophysical Research: Space Physics*, 123, 2018–2033. doi: <https://doi.org/10.1002/2017JA025121>
- Phan, T. D., Eastwood, J. P., Cassak, P. A., Gosling, J. T., Gershman, D. J., Mozer, F. S., ... Wilder, F. D. (2016). Mms observations of electron-scale filamentary currents in the reconnection exhaust and near the x line. *Geophysical Research Letters*, 43. doi: doi:10.1002/2016GL069212
- Pollock, C., Moore, T., Jacques, A., Burch, J., Gliese, U., Saito, Y., ... et al (2016). Fast plasma investigation for magnetospheric multiscale. *Space Science Reviews*, 199, 331–406. doi: <https://doi.org/10.1007/s11214-016-0245-4>
- Rager, A. C., Dorelli, J. C., Gershman, D. J., Uritsky, V., Avannov, L. A., Torbert, R. B., ... Saito, Y. (2018). Electron crescent distributions as a manifestation of diamagnetic drift in an electron-scale current sheet:

- 541 Magnetospheric multiscale observations using new 7.5 ms fast plasma in-  
 542 vestigation moments. *Geophysical Research Letters*, 45, 578-584. doi:  
 543 <https://doi.org/10.1002/2017GL076260>
- 544 Russell, C. T., Anderson, B. J., Baumjohann, W., Bromund, K. R., Dearborn, D.,  
 545 Fischer, D., ... Richter, I. (2016). The magnetospheric multiscale magne-  
 546 tometers. *Space Science Reviews*, 199, 189-256. doi: <https://doi.org/10.1007/s11214-014-0057-3>
- 547 Shay, M. A., Phan, T. D., Haggerty, C. C., Fujimoto, M., Drake, J. F., Malakit,  
 548 K., ... Swisdak, M. (2016). Kinetic signatures of the region surround-  
 549 ing the x line in asymmetric (magnetopause) reconnection. *Geophysical*  
 550 *Research Letters*, 43(9), 4145-4154. Retrieved from [https://agupubs](https://agupubs.onlinelibrary.wiley.com/doi/abs/10.1002/2016GL069034)  
 551 [.onlinelibrary.wiley.com/doi/abs/10.1002/2016GL069034](https://doi.org/10.1002/2016GL069034) doi:  
 552 <https://doi.org/10.1002/2016GL069034>
- 553 Shuster, J. R., Gershman, D. J., Chen, L.-J., Wang, S., Bessho, N., Dorelli, J. C., ...  
 554 Viñas, A. F. (2019). Mms measurements of the vlasov equation: Probing the  
 555 electron pressure divergence within thin current sheets. *Geophysical Research*  
 556 *Letters*, 46. doi: <https://doi.org/10.1029/2019GL083549>
- 557 Shuster, J. R., Gershman, D. J., Dorelli, J. C., Giles, B. L., Wang, S., Bessho, N.,  
 558 ... Torbert, R. B. (2021). Structures in the terms of the vlasov equation ob-  
 559 served at earth's magnetopause. *Nature Physics*. doi: <https://doi.org/10.1038/s41567-021-01280-6>
- 560 Sonnerup, B. U. Ö. (1979). Magnetic field reconnection. In *Solar system plasma*  
 561 *physics* (Vol. 3, p. 45-108).
- 562 Sonnerup, B. U. O., & Scheible, M. (1998). Minimum and maximum variance anal-  
 563 ysis. In *Analysis methods for multi-spacecraft data* (p. 1850). Noordwijk: ESA  
 564 Publication Division.
- 565 Vasyliunas, V. M. (1975). Theoretical models of magnetic field line merg-  
 566 ing. *Reviews of Geophysics*, 13(1), 303-336. Retrieved from [https://](https://agupubs.onlinelibrary.wiley.com/doi/abs/10.1029/RG013i001p00303)  
 567 [agupubs.onlinelibrary.wiley.com/doi/abs/10.1029/RG013i001p00303](https://doi.org/10.1029/RG013i001p00303)  
 568 doi: <https://doi.org/10.1029/RG013i001p00303>
- 569 Webster, J. M., Burch, J. L., Reiff, P. H., Daou, A. G., Genestreti, K. J., Graham,  
 570 D. B., ... Wilder, F. (2018). Magnetospheric multiscale dayside reconnec-  
 571 tion electron diffusion region events. *J. Geophys. Res. Space Physics*, 123,  
 572 4858-4878. doi: <https://doi.org/10.1029/2018JA025245>