

B23G-2491

Modeling Active Layer Depth of Permafrost under Changing Surface Boundary Conditions

<sup>1</sup>Jingfeng Wang ([jingfeng.wang@ce.gatech.edu](mailto:jingfeng.wang@ce.gatech.edu)), <sup>1</sup>Modi Zhu, <sup>1</sup>Husayn El Sharif, <sup>2</sup>Valeriy Y. Ivanov, <sup>3</sup>Aleksey Y. Sheshukov

<sup>1</sup>School of Civil & Environmental Engineering, Georgia Institute of Technology; <sup>2</sup>Department of Civil & Environmental Engineering, University of Michigan; <sup>3</sup>Department of Biological & Agricultural Engineering, Kansas State University

Summary

A physically based model is formulated for the active layer depth of permafrost under changing boundary condition instead of constant boundary condition considered in the traditional Stefan problem. Time-varying ground heat flux is obtained from net radiation and surface temperature using the Maximum Entropy Production (MEP) model as the driver of the active layer melting process. Conductive heat flux at the melting front is approximated in terms of an analytical function of ground heat flux. The simulated active layer depth is in good agreement with the field observations.

Classical Two-Phase Stefan Problem

Assumptions

1.

Semi-infinite slab  $0 \leq x < \infty$
2.

Initial condition  $T(x, 0) = T_s < T_m$  (melting temperature)
3.

Boundary condition  $T(0, t > 0) = T_L > T_m, \lim_{x \rightarrow \infty} T(x, t) = T_s$
4.

Melting front  $T(S(t), t) = T_m, S(0) = 0$   
 $\rho_s \lambda_f \dot{S}(t) = -k_L T_x(S(t), t)$  (Stefan condition)

$\rho_s$  ice density of bulk active layer  
 $C_s$  heat capacity of bulk active layer material  
 $k_L$  thermal conductivity of bulk active layer material  
 $\alpha_L$  thermal diffusivity of bulk active layer material  
 $\lambda_f$  latent heat of fusion ( $3.3 \times 10^5$  J kg<sup>-1</sup>)  
“s” – solid phase, “L” – liquid phase

Constant surface boundary condition  $T(0, t) = T_L$

Heat Transfer Equations

$$\frac{\partial T(x, t)}{\partial t} = \alpha_L \frac{\partial^2 T(x, t)}{\partial x^2}, \quad 0 < x < S(t)$$

active layer depth

Neumann Solution [Alexiades & Solomon, 1993]

$$S(t) = 2\lambda\sqrt{\alpha_L t}$$

$$\sqrt{\pi}\lambda\exp(\lambda^2)\text{erf}(\lambda) = St_L$$

(Stefan number)

Permafrost Active Layer Melting Problem

Conditions

1.

Semi-infinite slab  $0 \leq x < \infty$
2.

Initial condition  $T(x, 0) = T_m$
3.

Boundary condition  $T(0, t > 0) = T_L(t) > T_m,$   
 $\lim_{x \rightarrow \infty} T(x, t) = T_m$
4.

Melting front  $T(S(t), t) = T_m, S(0) = 0$   
 $\rho_s \lambda_f \dot{S}(t) = -k_L T_x(S(t), t)$

Changing surface boundary condition  $T(0, t) = T_L(t)$

Ground Heat Flux  $G$  Based Model of Active Layer Depth

Energy balance of active layer

$$\int_0^t G(\tau) d\tau = \int_0^{S(t)} C_s [T(x, t) - T_m] dx + \rho_s \lambda_f \int_0^{S(t)} \theta(x, t) dx$$

Approximate analytical solution of  $T(x, t)$  [Yang et al., 2017]

$$T(x, t) = T_m + \frac{1}{\sqrt{\pi C_s k_L}} \int_0^t \exp\left[-\frac{x^2}{4\alpha_L(t - \tau)}\right] \frac{G(\tau) d\tau}{\sqrt{t - \tau}}$$

Model Formulation

$$\int_0^t \text{erfc}\left[\frac{S(t)}{2\sqrt{\alpha_L(t - \tau)}}\right] G(\tau) d\tau = \rho_s \lambda_f \bar{\theta} S(t)$$

$\theta(x, t)$  water content of active layer  
 $\bar{\theta}$  depth averaged water content of active layer

MEP Model of ground heat flux [Wang and Bras, 2011]


$G$  over permafrost is parameterized based on the Maximum Entropy Production (MEP) Model of surface (latent  $E$  and sensible  $H$ ) heat fluxes using meteorological data:

$$G = R_n - E - H$$

$$\left[1 + B(\sigma) + \frac{B(\sigma)}{\sigma} \frac{\sqrt{C_s k_L}}{I_0 |H|^{1/6}}\right] H = R_n; E = B(\sigma) H$$

$$\sigma = \frac{\lambda_v^2}{c_p R_v} \frac{q_s}{T_L^2}; B(\sigma) = 6 \left(\sqrt{1 + \frac{11}{36} \sigma} - 1\right)$$


$R_n$  net radiation  
 $q_s$  surface specific humidity  
 $\lambda_v$  latent heat of vaporization  
 $c_p$  specific heat of air  
 $R_v$  gas constant of vapor



MEP modeled vs. field observed surface heat fluxes at a permafrost tundra site in Alaska [El Sharif et al., 2019].

Model Test

Field observations of soil temperature and ground meteorological data collected at an Arctic tundra (66.8163°, 65.5723°) and a tree (66.8526°, 65.6475°) site in the Polar Urals, Yamal-Nenets Autonomous District, Russia [Mazepa, 2005], June-July 2007.



Conclusion

Use of time-varying ground heat flux in modeling active layer depth directly links the permafrost melting process to its energy driver. This proof-of-concept study provides a promising alternative approach of modeling dynamics of permafrost freeze-thaw cycle.

References

▪ Alexiades, V., and A. D. Solomon (1993), *Mathematical Modeling of Melting and Freezing Processes*, Hemisphere Publishing Corp., Washington, 323pp.

▪ El Sharif, H., W. Zhou, V. Ivanov, A. Sheshukov, V. Mazepa, and J. Wang (2019), Surface Energy Budgets in Arctic Permafrost Areas during Growing Season, *J. Geophys. Res. – Atmos.*, 124, 6999 – 7017.

▪ Mazepa, V. S. (2005), Stand density in the last millennium at the upper tree-line ecotone in the Polar Ural Mountains, *Canadian Journal of Forest Research*, 35(9), 2082–2091.

▪ Wang, J., and R. L. Bras (2011), A Model of Evapotranspiration Based on the Theory of Maximum Entropy Production, *Water Resour. Res.*, 47, W03521.

▪ Yang, J., Z.-H. Wang, Q. Li, N. Vercauteren, E. Bou-Zeid, and M. B. Parlange (2017), A novel approach for unraveling the energy balance of water surfaces with a single depth temperature measurement, *Limnol. Oceanogr.* 62, 2017, 89–103.

**Acknowledgement** This study is sponsored by NSF Office of Polar Programs Grants 1724633 (Georgia Tech), 1725654 (University of Michigan) and 1724868 (Kansas State University).