

# Global Driving of Auroral Precipitation : 1. Balance of Sources

Agnit Mukhopadhyay<sup>1,2,3</sup>, Daniel Welling<sup>4</sup>, Michael Liemohn<sup>1</sup>, Aaron Ridley<sup>1</sup>,  
Meghan Burleigh<sup>5</sup>, Chen Wu<sup>1</sup>, Shasha Zou<sup>1</sup>, Hyunju Connor<sup>6,2</sup>, Elizabeth  
Vandegriff<sup>4</sup>, Pauline Dredger<sup>4</sup>, Gabor Tóth<sup>1</sup>

<sup>1</sup>Climate and Space Sciences and Engineering Department, University of Michigan, Ann Arbor, MI, USA

<sup>2</sup>NASA Goddard Space Flight Center, Greenbelt, MD, USA

<sup>3</sup>Department of Physics, American University, Washington D.C., USA

<sup>4</sup>Department of Physics, University of Texas at Arlington, Arlington, TX, USA

<sup>5</sup>Naval Research Laboratory, Washington D.C., USA

<sup>6</sup>Department of Physics, University of Alaska Fairbanks, Fairbanks, AK, USA

## Key Points:

- A semi-physical global modeling approach is used to estimate diffuse & discrete sources of auroral precipitation during the Galaxy15 event.
- Diffuse sources contribute 74% of the total auroral power. Discrete sources are strongly driven by activity & can contribute up to 61%.
- Broadband precipitation contributes 31% of the auroral Pedersen conductance playing a significant role in ionospheric electrodynamics.

## Abstract

The accurate determination of auroral precipitation in global models has remained a daunting and rather inexplicable obstacle. Understanding the calculation and balance of multiple sources that constitute the aurora, and their eventual conversion into ionospheric electrical conductance, is critical for improved prediction of space weather events. In this study, we present a semi-physical global modeling approach that characterizes contributions by four types of precipitation - monoenergetic, broadband, electron and ion diffuse - to ionospheric electrodynamics. The model uses a combination of adiabatic kinetic theory and loss parameters derived from historical energy flux patterns to estimate auroral precipitation from magnetohydrodynamic (MHD) quantities. It then converts them into ionospheric conductance that is used to compute the ionospheric feedback to the magnetosphere. The model has been employed to simulate the April 5 - 7, 2010 "*Galaxy15*" space weather event. Comparison of auroral fluxes show good agreement with observational datasets like NOAA-DMSP and OVATION Prime. The study shows a dominant contribution by electron diffuse precipitation, accounting for  $\sim 74\%$  of the auroral energy flux. However, contributions by monoenergetic and broadband sources dominate during times of active upstream solar conditions, providing for up to 61% of the total hemispheric power. The study also finds a greater role played by broadband precipitation in ionospheric electrodynamics which accounts for  $\sim 31\%$  of the Pedersen conductance.

## Plain Language Summary

The aurora is comprised of electrically charged particles that enter the upper atmosphere from outer space. The entry is driven by diverse processes at different locations of the high-latitude atmosphere; these help define the different sources that constitute the bulk of the aurora. Since the aurora is an important phenomena in the study of near-Earth space physics and space weather, it is important to account for the contribution and balance of each individual source and deduce their impact. In this study, we have introduced a novel modeling approach that is capable of estimating contributions from four diverse sources of aurora, and used this approach to study auroral dynamics during a famous space weather event. Our results indicate that the proportion and strength of each source varies over time, location and activity. Additionally, we identify which sources have a pronounced contribution to the ionosphere's electrical conductance.

## 1 Introduction

High-latitude precipitation of charged particles is a crucial driver of ionospheric electrodynamics (e.g. Kivelson & Russell, 1995). These particles precipitate from the near-Earth plasma environment to form the aurora, and enhance the electrical conductance in the polar regions (e.g. Schunk & Nagy, 2009). Auroral precipitation is broadly defined into two types: diffuse and discrete aurora. Particles scattered into the loss cone by plasma waves create the diffuse aurora (Nishimura, Lessard, et al., 2020 and references therein). Diffuse particles precipitate into the upper atmosphere without the need of acceleration, and can consist of both electrons (e.g. Evans & Moore, 1979) and ions (e.g. Sergeev et al., 1983). Conversely, the discrete aurora is generated by particles that are accelerated into the ionosphere (e.g. Korth et al., 2014). These particles can be accelerated by geomagnetic field-aligned electric fields (monoenergetic; e.g. Evans, 1974; Knight, 1973) or by dispersive Alfvén waves (broadband; e.g., Ergun et al., 1998; Chaston et al., 2003). The conductance enhancements caused by auroral precipitation are important to investigative studies of magnetosphere-ionosphere coupling (e.g. Öztürk et al., 2020), since it regulates the closure of field-aligned currents (FACs; Iijima & Potemra, 1976) and maintain the nonlinear feedback loop between the magnetosphere and the ionosphere (e.g. Merkin et al., 2003; Ridley et al., 2004). Since auroral currents are the dom-



inant drivers of ground-based magnetic perturbations in high-latitude regions (e.g. Welling, 2019), auroral conductance is a crucial regulator of ground-based space weather activity (Hartinger et al., 2017; Mukhopadhyay et al., 2020).

Despite their importance, the computation of auroral precipitation and derived conductances is not trivial in most global models. This is due to several reasons. First, the first-principles-driven calculation of conductance needs to account for ionosphere-thermosphere dynamics like atmospheric chemistry and reaction rates (Yu et al., 2016). Most global models work around the complexity of ionosphere-thermosphere dynamics by the use of empirical relationships like Robinson et al. (1987), Galand et al. (2001) and Kaeppler et al. (2015), that derive perpendicular conductances from precipitating fluxes. This method has limitations, as the empirical relations are based off of limited dataset and have numerous associated uncertainties (Liemohn, 2020; Welling et al., 2017). Even so, most global models assume a two-dimensional ionospheric domain (e.g. Goodman, 1995) which makes usage of an empirical conversion between fluxes and conductances undemanding. Recent work by Burleigh et al. (2019) has sought to incorporate a dedicated ionosphere-thermosphere solver to incorporate realistic chemistry and altitudinal ionization rates to provide for a more accurate estimation of the conductance.

Second, estimating the kinetic description of particle precipitation is not straightforward in a global setup. This is especially challenging in magnetohydrodynamic (MHD) models which do not resolve pitch angle distributions and wave scattering. Fedder et al. (1995) expressed electron auroral energy and number fluxes as functions of MHD parameters using adiabatic kinetic theory. This work was further expanded by successive studies (Raeder et al., 2001; Wiltberger et al., 2009; Gilson et al., 2012; Zhang et al., 2015; Yu et al., 2016, 2018) who sought to improve the original methodology and incorporate multiple types of precipitation in the computation of ionospheric conductance in the aurora. Since adiabatic kinetic theory does not fully account for the kinetic physics of loss cone distributions, models like Ridley et al. (2004) and Mukhopadhyay et al. (2020) pursued a bypass, by using empirical relationships to derive precipitation with FACs. While this simplified the process of estimating auroral precipitation, the models only provide conductances in regions of high FACs and have statistical limits to physical phenomena like auroral expansion during extreme driving.

While the influence of auroral conductance on magnetospheric dynamics, ionospheric electrodynamics, and their coupled nonlinear feedback system are well known (e.g., Ridley et al., 2001; Liemohn et al., 2005; Ebihara et al., 2005; Zheng et al., 2008; Welling & Ridley, 2010; Connor et al., 2016; Ozturk et al., 2017), the contribution of each individual source of precipitation has not been widely studied, especially for variable solar wind driving. This is challenging to do with data, since most measurements of ionospheric conductance have significant underlying challenges and uncertainties (Ohtani et al., 2014). Empirical modeling efforts by Newell et al. (2009, 2014) have thoroughly studied the balance of auroral precipitation through the determination of multiple sources from in-situ observations. Earlier studies by Hardy et al. (1985, 1989) and Brautigam et al. (1991) have sought to provide balance between different sources using upstream and/or space weather conditions. The comparison of observed FACs with in-situ precipitation by Korth et al. (2014) provided further quantification of discrete sources of precipitation. Despite this, empirical approaches are limited by observational findings and lack the global perspective to relate drivers affecting auroral precipitation with quantities that they influence. Such a relationship can more easily be studied through a global first-principles-based modeling approach. Furthermore, with increasing usage of first-principles-based geospace models for operations-grade space weather prediction (e.g. Pulkkinen et al., 2011, 2013; Rastätter et al., 2016; Cash et al., 2018), the need to quantify the impact of multiple sources of conductance on the M-I feedback becomes ever more necessary.

This work describes the development of a novel modeling approach that predominantly uses a semi-physical method to estimate four sources of precipitation - monoen-

ergetic, broadband, electron and ion diffuse - using input variables from the Space Weather Modeling Framework (SWMF; Tóth et al., 2005, 2012). This model has been used to study salient aspects of the April 5 - 7, 2010 space weather event (e.g. Chen et al., 2015; Keese et al., 2014) in order to determine the individual contributions of each source. In this endeavour, the article aims to address the following tasks: (1) quantification of individual contribution by each source of precipitation, (2) comprehension of the impact of upstream solar wind drivers on each source, and (3) discernment of the impact of each source on the net ionospheric conductance. Each task is addressed in Section 4 through the following comparative experiments:

1. Modeled auroral fluxes are validated against both patterns and hemispheric integrated quantities.
2. A case study is investigated that describes changes in each source of precipitation with changing upstream conditions.
3. Precipitation from each source is converted into ionospheric Hall and Pedersen conductance, and their contributions to both conductances are quantified.


## 2 Numerical Methodology

### 2.1 Geospace Numerical Setup of SWMF

The geospace version of SWMF (see model layout in Figure 1), that is currently used for real-time space weather prediction at NOAA-SWPC (Cash et al., 2018), consists of three independent models that are numerically coupled together. The Block Adaptive Tree Solar-Wind Roe Upwind Scheme (BATS-R-US; Powell et al., 1999; Gombosi et al., 2003) model uses ideal semi-relativistic single-fluid magnetohydrodynamic (MHD) equations to simulate the global magnetosphere. BATS-R-US' magnetospheric domain is a three dimensional space in GSM coordinates. In the  $x$  axis, the domain spans  $32 R_E$  on the dayside and  $224 R_E$  on the nightside, while in the  $y$  and  $z$  coordinate axes, the domain spans  $128 R_E$  in either directions. The model uses a block-adaptive Cartesian grid to ensure highest spatial resolution in regions of interest. The grid resolution used in this study is similar to the *Hi-Res SWPC* grid used in the study by Mukhopadhyay et al. (2020), details about which are found in Appendix A of Haiducek et al. (2017). To better capture energy-dependent drift physics in the inner magnetosphere, BATS-R-US is coupled to a dedicated inner magnetosphere model (De Zeeuw et al., 2004). In the present setup, we have used the Rice Convection Model (RCM; e.g. Wolf et al., 1982) which solves for bounce-averaged particle distribution in the ring current region. To drive this, RCM uses flux tube volumes from BATS-R-US and adjusts the MHD pressure and density in return.

SWMF also has a dedicated coupling to an ionospheric solver, the Ridley Ionosphere Model (RIM; e.g. Ridley et al., 2001), which computes ionospheric electrodynamics at an altitude of 110 km. RIM is a finite-difference Poisson solver that computes the electrostatic potential and horizontal currents using field-aligned currents (FACs) as input and a *prescribed* conductance pattern (Goodman, 1995). FACs are mapped down from near the inner boundary of BATS-R-US (typically between  $3.5 R_E$  and  $2.5 R_E$ ) to ionospheric altitudes ( $\sim 110$  km). Additionally, BATS-R-US has optional settings to return plasma pressure and density values mapped near its inner boundary (De Zeeuw et al., 2004), which is then mapped onto the ionospheric grid (Yu et al., 2016). RIM returns the ionospheric potential to both BATS-R-US and RCM, which are subsequently used as inner boundary conditions.

Ridley et al. (2004) describes the computation of conductance in RIM. The solver estimates contributions by multiple sources like solar EUV illumination, auroral precipitation and polar rain. Conductance due to EUV illumination affects the dayside, and is computed as a function of the solar zenith angle (Moen & Brekke, 1993). Enhance-




**Figure 1.** Numerical coupling between models within the geospace setup of SWMF.

ments due to starlight conductance and polar rain generally affect the nightside ionosphere, and are added in as constants. Conductance enhancements due to auroral precipitation is relatively complicated to estimate, since the aurora is driven by upstream driving conditions. In RIM, auroral conductance is typically computed by one of two dedicated models - Ridley Legacy Model (RLM; Ridley et al., 2004) or the Conductance Model for Extreme Events (CMEE; Mukhopadhyay et al., 2020). Both models are based off of assimilative maps (Ridley & Kihn, 2004), and use empirical relationships with FACs to estimate the auroral conductance (see Mukhopadhyay et al., 2020 for details).

## 2.2 The MAGNetosphere-Ionosphere-Thermosphere (MAGNIT) Auroral Precipitation Model

In this study, we introduce the MAGNetosphere-Ionosphere-Thermosphere (MAGNIT) auroral model that computes individual sources of precipitation in SWMF. MAGNIT culminates from a series of modeling developments within RIM (Mukhopadhyay et al., 2018, 2019; Liemohn et al., 2018; Burleigh et al., 2019; Mukhopadhyay, Burleigh, et al., 2021) that replace the existing empirical conductance models with state-of-the-art numerical couplers and solvers to estimate auroral dynamics. Figure 2 describes the ionospheric setup of MAGNIT within RIM. Advanced numerical couplings are introduced to transfer fieldline traced values of bulk quantities like pressure and density from BATSR-US to RIM, mapping the values down from the inner boundary of MHD to 110 km altitude. MAGNIT uses adiabatic kinetic theory to compute auroral fluxes from the MHD state variables (e.g. Fedder et al., 1995). This investigation uses MAGNIT to estimate four different sources of precipitation - electron diffuse, ion diffuse, monoenergetic and broadband - and quantify their individual contributions. The computation of each source is described in the following.



**Figure 2.** Schematic description of auroral conductance and subsequent ionospheric E-field computation within RIM when equipped with MAGNIT. Dotted lines indicate optional couplings when RLM or CMEE is deployed. Note: Colorbar scale is not uniform across dial plots.

### 2.2.1 Electron and Ion Diffuse Precipitation

Parameterization of diffuse fluxes in MAGNIT observes the following formulation:

$$\Phi_{N_{\text{diff}(e,i)}} = \alpha_1(e, i) \times \frac{N_{e,i} T_{e,i}^{1/2}}{\sqrt{2\pi m_{e,i}}} \quad (1)$$

$$\Phi_{E_{\text{diff}(e,i)}} = \alpha_2(e, i) \times \frac{2N_{e,i} T_{e,i}^{3/2}}{\sqrt{2\pi m_{e,i}}} \quad (2)$$

Here,  $\Phi_{N_{\text{diff}}}$  and  $\Phi_{E_{\text{diff}}}$  are the diffuse number and energy fluxes respectively,  $N$  is the particle number density,  $T$  is particle thermal temperature (derived from MHD) and  $m$  is the particle mass. The equations have been denoted for both electrons and ions by subscripts  $e$  and  $i$ , respectively. The multipliers  $\alpha_1(e, i)$  and  $\alpha_2(e, i)$  act as the particle filling rate of the loss cone, and are optimized using data-model comparisons (discussed in more detail in Section 3). This is necessary because global MHD is unable to resolve pitch angle distributions, and therefore cannot accurately predict particle loss from the magnetosphere without further parameterization (e.g. Zhang et al., 2015). The remaining fractions are the standard solution for total particle and energy flux contained in a Maxwellian distribution for all pitch angles (e.g. Gombosi, 1994).

$T_e$  is assumed to be 1/6 of the single-fluid MHD temperature,  $T_{\text{MHD}}$ , based on an electron-to-ion temperature ratio of 1:5, as observed in the plasma sheet (Paschmann et al., 1993; Phan et al., 1994; Fedder et al., 1995). This is a simplistic approximation however; though this relationship is typically valid for low-energy plasma sheet particles in the near-Earth region (Wang et al., 2012), a more accurate representation between the electron and ion temperature is planned in future modeling developments. This will incorporate the independent computation of electron plasma temperature using a two- or multi-fluid MHD approach (e.g. Glocer et al., 2009, 2020) to work around this approximation. Unlike some modeling approaches (e.g. Zhang et al., 2015), the precipitation pattern of either diffuse sources do not undergo any spatial variations. This is done in order to better retain dynamic changes in auroral boundaries across activity levels. Ion precipitation occurs in regions where the field line curvature becomes comparable to the particle gyroradius (Sergeev & Tsyganenko, 1982; Sergeev et al., 1983). To simulate this, we have used a modified version of the model developed by Gilson et al. (2012), where a step function is used as a function of the  $\kappa$ -parameter (e.g. Büchner & Zelenyi, 1987). In lieu of a fieldline-estimated  $\kappa$ , an equatorward and poleward boundary based on the peak strength of fieldline-traced pressure was used to define the bounds of the step function.

After the computation of the ion and electron fluxes, the average energy  $\bar{E}_{\text{diff}(e,i)}$  is computed as  $\Phi_{E_{\text{diff}(e,i)}} / \Phi_{N_{\text{diff}(e,i)}}$ . Using the empirical relationships developed by Robinson et al. (1987), the average energy and energy flux for the electron diffuse source is converted into Hall and Pedersen conductances. Similarly, MAGNIT uses the ion flux and energies to compute the ion-driven conductance via the Galand & Richmond (2001) empirical relationships.

### 2.2.2 Monoenergetic Precipitation

MAGNIT estimates monoenergetic precipitation using the Knight-Fridman-Lemaire (KFL; Knight, 1973; Lyons et al., 1979; Fridman & Lemaire, 1980) relationship that estimates electrons accelerated by a quasi-static parallel potential drop along a magnetic field-line. We broaden our assumption of an isotropic Maxwellian particle distribution by estimating incident electrons upon a field-aligned potential drop  $V$ . Using this modification, we follow the KFL procedure to estimate the first and third moment of the dis-



tribution to compute the downward number and energy flux (e.g. Zhang et al., 2015; Yu et al., 2016).

$$\Phi_{N_{\text{mono}}} = \alpha_3 \times \frac{N_e T_e^{1/2}}{\sqrt{2\pi m_e}} \left[ \frac{B_{iono}}{B_{ps}} - \left( \frac{B_{iono}}{B_{ps}} - 1 \right) e^{-\frac{q_e V B_{ps}}{T_e (B_{iono} - B_{ps})}} \right] \quad (3)$$

$$\Phi_{E_{\text{mono}}} = \alpha_4 \times \frac{2N_e T_e^{1/2}}{\sqrt{2\pi m_e}} \left[ \frac{1 - e^{-\frac{q_e V B_{ps}}{T_e (B_{iono} - B_{ps})}}}{2 \left( 1 + \left( \frac{B_{iono} - B_{ps}}{B_{iono}} \right) e^{-\frac{q_e V B_{ps}}{T_e (B_{iono} - B_{ps})}} \right)} q_e V + T_e \right] \times \left[ \frac{B_{iono}}{B_{ps}} - \left( \frac{B_{iono}}{B_{ps}} - 1 \right) e^{-\frac{q_e V B_{ps}}{T_e (B_{iono} - B_{ps})}} \right] \quad (4)$$

Similar to equations 1 and 2,  $\Phi_{N_{\text{mono}}}$  and  $\Phi_{E_{\text{mono}}}$  stand for the monoenergetic number and energy fluxes respectively,  $N_e$  is the electron number density,  $T_e$  is the electron temperature in the night-side plasma sheet,  $m_e$  is electron mass, and  $q_e$  is the elementary charge.  $B_{iono}$  and  $B_{ps}$  signify the magnetic field strengths at ionospheric altitude (assumed at 110 km in the present model) and at the source region respectively; their ratio  $B_{iono}/B_{ps}$  is the magnetic mirror ratio. The source region in our simulations is assumed to be the plasma sheet (Yu et al., 2016) and the mirror ratio is assumed to be dipolar. The multipliers  $\alpha_3$  and  $\alpha_4$  are parameters that represent the degree of loss-cone filling in the electron source region. Unspecified by the MHD fluid approach, we change these factors to scale the resulting fluxes.

In regions of upward field-aligned current ( $J_{||}$ ), assuming that the current is entirely carried by electrons, the number flux of monoenergetic precipitation can be expressed as:

$$\Phi_{N_{\text{mono}}} = J_{||}/q_e \quad (5)$$

Therefore, the potential drop  $V$  can be expressed in terms of  $J_{||}$  as

$$V = \frac{T_e (B_{ps} - B_{iono})}{q_e B_{ps}} \ln \left( \frac{B_{iono} - B_{ps} \frac{J_{||}}{q_e \alpha_3 N_e T_e^{1/2}}}{B_{iono} - B_{ps}} \right) \quad (6)$$

The above condition is only valid for the logarithm not being zero, which indicates that

$$B_{ps} \leq B_{ps} \frac{J_{||}}{q_e \alpha_3 N_e T_e^{1/2}} \leq B_{iono} \quad (7)$$

must be satisfied for a monoenergetic source of precipitation. Also, the potential structure along the field line must satisfy the relationships described by Chiu & Schulz (1978), in which case a more complicated approach is needed (e.g., Liemohn & Khazanov, 1998). Fortunately, this influence is small for potential drops accelerating electron auroral precipitation (e.g., Khazanov et al., 1998). In the absence of parallel fields ( $V = 0$ ), the above equations are reduced to equations 1 and 2 that model the electron diffuse precipitation. Similar to diffuse precipitation, the average energy for monoenergetic precipitation  $\bar{E}_{\text{mono}}$  is computed as  $\Phi_{E_{\text{mono}}}/\Phi_{N_{\text{mono}}}$ . Using the empirical relation by Robinson et al. (1987), the Hall and Pedersen conductances are computed and combined with the diffuse sources using a vector sum.

### 2.2.3 Broadband Precipitation

Broadband precipitation is driven by low energy electrons that are accelerated by dispersive Alfvén waves (e.g. Ergun et al., 1998). Successive investigations (Chaston et al., 2003; Strangeway, 2010; Zhang et al., 2014, 2015) have characterized a relationship between broadband flux and the Alfvénic Poynting flux. In MAGNIT, we use a similar

approach as in Zhang et al. (2015), to estimate the broadband number and electron flux as an empirical function of the Poynting flux. The relation is shown as

$$\Phi_{N_{\text{bbnd}}} = 3 \times 10^9 \times (\alpha_5 \times S_{||})^{0.47} \quad (8)$$

$$\Phi_{E_{\text{bbnd}}} = 2 \times (\alpha_6 \times S_{||})^{0.5} \quad (9)$$

Similar to previous sources,  $\Phi_{N_{\text{bbnd}}}$  and  $\Phi_{E_{\text{bbnd}}}$  stand for the broadband number and energy fluxes respectively, which are collectively used to define the broadband average energy  $\langle E_{\text{bbnd}} \rangle$  as  $\Phi_{E_{\text{bbnd}}} / \Phi_{N_{\text{bbnd}}}$ .  $S_{||}$  is the Poynting flux into the ionosphere. Unlike the model by Zhang et al. (2015), MAGNIT does not use AC Poynting flux which is a more direct measure of small-scale Alfvénic energy, but instead relies on DC Poynting flux (Yu et al. 2010) derived from Joule heating in the ionosphere (e.g. Rastätter et al., 2016). Due to the empirical nature of the above equations, the multipliers  $\alpha_5$  and  $\alpha_6$  act as empirical moderators of the Poynting energy, since DC Poynting flux typically has higher values than electron precipitation (Janhunen et al., 2005). Broadband fluxes are converted into electron-driven conductances by the Robinson et al. (1987) relationship.

Following Zhang et al. (2015), the broadband contribution is added linearly to the total conductance due to it enhancing ion density in the bottomside F-region rather than the E region of the ionosphere. Since the rest of the sources are added as a vector sum (Wallis & Budzinski, 1981), the final sum of auroral sources to ionospheric conductance is computed as follows.

$$\Sigma_{\text{Aurora}} = \sqrt{\Sigma_{e^- \text{ diff}}^2 + \Sigma_{i^+ \text{ diff}}^2 + \Sigma_{\text{mono}}^2} + \Sigma_{\text{bbnd}} \quad (10)$$

where  $\Sigma_{\text{Aurora}}$  stands for the total auroral conductance (Hall or Pedersen)  $\Sigma_{e^- \text{ diff}}$ ,  $\Sigma_{i^+ \text{ diff}}$ ,  $\Sigma_{\text{mono}}$  and  $\Sigma_{\text{bbnd}}$  stand for Hall or Pedersen conductances from the electron diffuse, ion, monoenergetic and broadband sources, respectively.

### 2.3 Comparisons against Observations & Empirical Models

In order to quantify the contribution by each source to the net auroral flux and energies, the hemispheric power (integrated energy flux over a hemisphere), hemispheric number flux (integrated number flux over a hemisphere), and overall average energy (hemispheric power / hemispheric number flux) were computed. Percent contributions by each source of precipitation ( $\%C_{\text{source}}$ ) were defined as

$$\%C_{\text{source}} = \frac{Q_{\text{source}}}{Q_{\text{aurora}}} \quad (11)$$

where  $Q_{\text{source}}$  stands for a given quantity from an individual source, and  $Q_{\text{aurora}}$  stands for the same quantity from all auroral sources combined, with quantity  $Q$  being either hemispheric power or hemispheric number flux. Determining the contribution to ionospheric conductance is not as straightforward as in the case of energy flux and number flux. This is because the broadband source of conductance is added linearly to the total conductance. Therefore, contribution to the conductance has been defined in two ways - the total contribution, which computes the percentage contribution of a source as a fraction of the linear sum of conductances (similar to Equation 11), and the resultant contribution, which computes the percentage contribution for the sources in the following way:

$$\%C_{\text{source}} = \frac{\Sigma_{\text{source}}^2}{\Sigma_{\text{aurora}}^2} \quad \text{where source} = e^- \text{ Diff}, i^+ \text{ Diff}, \text{ Mono} \quad (12)$$

and

$$\% \text{ C}_{\text{broadband}} = 1 - \frac{(\Sigma_{e^{-}\text{Diff}}^2 + \Sigma_{i^{+}\text{Diff}}^2 + \Sigma_{\text{Mono}}^2)}{\Sigma_{\text{aurora}}^2} \quad (13)$$

where  $e^{-}\text{Diff}$ ,  $i^{+}\text{Diff}$  and Mono indicate electron diffuse, ion diffuse and monoenergetic sources respectively, and  $\Sigma$  is the Hall or Pedersen conductance.

Modeled results of auroral fluxes have been evaluated through comparisons against observations and multiple derived-estimates. The study uses hemispheric power data from in-situ observations measured by the Defense Meteorological Satellite Program (DMSP) and National Oceanic and Atmospheric Administration (NOAA) satellites (Emery et al., 2006, 2008). The observations are hourly-averaged and span the past  $\sim 30$  years (1978 - 2013 for electrons; 1983 - 2013 for ions). The study uses the empirical models, OVA-TION Prime (Newell et al., 2009; shortened to OV Prime) and the AE-driven Feature Tracking of Aurora (FTA; Wu et al., 2021) for comparison of energy flux and average energies in the auroral region. OV Prime is developed from multiple observations from DMSP during the years 1988 - 1998, while FTA is based on 1.5 years of Polar Ultraviolet Imager data. Both OV Prime and MAGNIT account for multiple (and similar) sources of precipitation. This has enabled us to compare individual contributions to hemispheric power, number flux and energies in this study. Multiple empirical functions relating hemispheric power and number flux to space weather indices, auroral electrojet parameters and/or upstream conditions exist. While the usage of all such models is not possible, this study identified and employed five empirical models - Brautigam et al. (1991), Ahn et al. (1983), Lu et al. (1998), Østgaard et al. (2002) and Korth et al. (2014) - to compare modeled predictions. Of these, the first four models provide total hemispheric power estimates - the model by Brautigam et al. (1991) estimates fluxes for electrons and ions separately, and is driven using solar wind inputs; remaining models are driven using AE/AU/AL values, and seek to establish a relationship between the energy deposition by electron precipitation and geomagnetic indices (see Østgaard et al., 2002 for detailed comparisons). The model by Korth et al. (2014) estimates discrete energy fluxes using upward FACs, and was predominantly used to validate modeled monoenergetic and broadband precipitation.

Additionally, simulated cross-polar cap potential (CPCP), integrated FACs (iFACs; Anderson et al., 2017),  $Kp$  and Sym-H were also compared against observations. CPCP values were compared against derived estimates obtained from the AMIE model and the Super Dual Auroral Radar Network (SuperDARN; e.g., Khachikjan et al., 2008). Observations for  $Kp$  and Sym-H were obtained from the Kyoto Observatory, while iFAC observations were acquired from the the Active Magnetosphere and Planetary Electrodynamics Response Experiment (AMPERE) mission (Anderson et al., 2014; Waters et al., 2020).

Evaluation of modeled results against observations have been quantified using two metrics - median absolute percentage error (MAPE) and the exclusion parameter (EP). MAPE provides an absolute value of the relative percentage error, and has been commonly used as a measure of accuracy of prediction (e.g. Morley et al., 2018). EP measures the accuracy of prediction against multiple observation-derived estimates, by accounting for all data that is outside the range of observed values (Mukhopadhyay, Jia, et al., 2021). Low values of both metrics is generally assumed as a good prediction (e.g. Liemohn et al., 2021).

### 3 Parameterization of Sources

Due to its inability to compute loss-cone distributions, MAGNIT uses flux multipliers  $\alpha_s$  (for each source  $s$ ) to regulate the final value of energy and number flux. A comparative study of modeled results to OV Prime and NOAA-DMSP was undertaken to determine  $\alpha_s$ . Initially, a similar procedure to Zhang et al. (2015) was applied - hemi-




Run	$B_{(x,y,z)}$ (nT)	$\frac{d\Phi_{MP}}{dt}$
A	(0, 0, -1)	0.29e+04
B	(0, 0, -2)	0.47e+04
C	(0, 0, -3)	0.61e+04
D	(0, 0, -4)	0.74e+04
E	(0, 0, -5)	0.86e+04
F	(0, 0, -6)	0.97e+04
G	(0, 0, -7)	1.08e+04
H	(0, 0, -8)	1.18e+04
I	(0, 0, -9)	1.28e+04
J	(0, 0, -10)	1.37e+04
K	(0, 0, -11)	1.46e+04
L	(0, 0, -12)	1.54e+04
M	(0, 0, -13)	1.63e+04
N	(0, 0, -14)	1.71e+04
O	(0, 0, -15)	1.79e+04

**Table 1.** IMF conditions used in each test run to determine  $\alpha_s$ . A solar wind velocity of 400 km/s in the x-direction and a number density of  $5 \text{ cm}^{-3}$  were used in each test run. The third column shows the Newell coupling function,  $d\Phi_{MP}/dt = v^{4/3}(B_x^2 + B_y^2)^{1/3} \sin^{8/3} \theta_C$ , where  $\theta_C$  is the IMF clock angle, is listed in the fifth column.

spheric fluxes from each auroral source were compared against OV Prime estimates to initialize  $\alpha_s$  for each source. Both SWMF and OV Prime were run for diverse driving conditions - Table 1 lists idealized solar wind conditions used for these simulations. For each SWMF run, the magnetosphere was preconditioned for 6 hours by first driving with southward IMF  $B_z = -5 \text{ nT}$  for 3 hours, followed by northward IMF  $B_z = +5 \text{ nT}$  for 3 hours. Post preconditioning, the simulation was driven for 3 more hours with the values listed in Table 1. The solar wind velocity, number density and temperature in these runs were kept the same as Zhang et al. (2015) for consistency.

Figure 3a and b compare hemispheric power and number flux predictions respectively from MAGNIT against OV Prime values. Each subplot corresponds to a distinct source. The x-axis in each subplot displays the variation in driving conditions, using the Newell function  $d\Phi_{MP}/dt$  (Newell et al., 2007) and IMF  $B_z$ .  $\alpha_s$  for each source was defined using OV Prime fluxes ( $Q_{OP}$ ) and MAGNIT fluxes ( $Q_{MAGNIT}$ ) as the median of their ratio, i.e.  $\alpha_s = \text{median}(Q_{OP}/Q_{MAGNIT})$ . The final values of MAGNIT-estimated precipitation (denoted in blue) were scaled by multiplication with the initialized values of  $\alpha_s$ , for comparison against OV Prime estimates (in black-red). In parts a(iii - iv) and b(iii - iv) of Fig. 3, monoenergetic and broadband fluxes show good agreement with OV Prime values when scaled. However, for both diffuse sources (parts a(i - ii) and b(i - ii) in Fig. 3), MAGNIT estimates are much higher than OV Prime, and the two sets of prediction diverge from each other beyond an IMF  $B_z$  value of -6 nT. The difference in diffuse fluxes is further elucidated in Figure 3c, where dial plots of electron diffuse energy flux from OV Prime (left) and MAGNIT (right) are compared for Runs A, H and O (see Table 1). With increasing activity, MAGNIT displays stronger auroral precipitation and significant expansion in the auroral oval. By contrast, OVATION Prime caps the max value of the energy flux, and displays minute expansion of the auroral oval. OV Prime has been shown to underestimate hemispheric fluxes during extreme driving (Newell et al., 2014). The oval expansion is an aspect of MAGNIT's usage of MHD pressure to compute diffuse fluxes (see Section 5 for further details).

### Determining $\alpha_s$ - Initial Comparisons against OVATION Prime




**Figure 3.** Initial determination of  $\alpha_s$  by comparing MAGNIT runs against OVATION Prime values. Comparison of (a) hemispheric power and (b) total hemispheric number flux for (i) electron diffuse, (ii) ion diffuse, (iii) monoenergetic and (iv) broadband precipitation through varying driving conditions. Here, IMF  $B_T = \sqrt{B_y^2 + B_z^2}$ . (c) Dial plot comparison of diffuse electron precipitation from both models for (row-wise) Runs A, H and O, indicating change in morphology and auroral strength with increasing activity.

To verify diffuse contributions, the comparative analysis was expanded to include hourly-averaged hemispheric power observed by the NOAA-DMSP satellite chain over the past >30 years (1978 - 2013) (Emery et al., 2006, 2008). This is, in essence, an extension to the comparison against OV Prime, since OV Prime itself is based on multiple observations from DMSP. Unlike OV Prime, the NOAA-DMSP observations do not distinguish between different precipitative sources, instead informing only of contributions made by electrons and ions to the total energy flux respectively. Therefore, the extended comparisons were conducted in two phases - (1) Verifying discrete contributions against Korth et al. (2014), and (2) comparing total electron and ion fluxes against NOAA-DMSP. The empirical relationship given in Korth et al. (2014) relates upward FACs in the dusk-afternoon sector with discrete (mono+broadband) energy flux. Comparisons of discrete hemispheric power in the dusk-afternoon sector for both models are shown in Figure 4(i). The *Korth14* model was driven using FAC estimates from each SWMF run. MAGNIT results display reasonable agreement with the *Korth14* model in Subplot 4(i-c) when scaled using the  $\alpha_s$  values deduced from Fig. 3. Values for  $\alpha_s$  are determined as the median of observed-to-modeled ratio for each source. In Subplot 4(i-c), these values were determined for monoenergetic and broadband sources separately to deduce the total discrete contribution. Comparison of model-model energy flux distributions in the dusk-afternoon sector (parts i-a and i-b of Fig. 4) also exhibit similarities in pattern.


Figure 4(ii) shows final comparisons between NOAA-DMSP observations (in green) and MAGNIT estimates (in blue) over varying IMF  $B_z$ . The values from OV Prime (in black-red) have also been plotted for clarity. The NOAA-DMSP dataset was binned by the Newell function ( $d\Phi_{MP}/dt$ ) to provide median values for each bin. Since the in-situ measurements do not observe number flux of precipitating particles, they were derived using average energies of electrons and ions from Hardy et al. (1985, 1989).  $\alpha_s$  values

## Determining $\alpha_s$ - Comparison against NOAA-DMSP & Empirical Predictions

### (i) Comparison of Discrete Hemispheric Flux



### (ii) Comparisons against NOAA-DMSP



**Figure 4.** Finalization of  $\alpha_s$  values by comparing MAGNIT runs against in-situ observations and empirical predictions. (i) Comparison of discrete HP - (left) Post-noon quarter-dial plots of (a) Korth et al. (2014) and (b) MAGNIT at IMF  $B_T = 8$  nT, (right) Quarter-hemispheric power across variable driving conditions for same models. (ii) Comparison against NOAA-DMSP HP - Comparison of hemispheric power for (a) electron and (b) ion power, and (c-d) their respective number fluxes compared against NOAA-DMSP and OV Prime estimates over varying driving conditions.

Source	$\alpha_s$ (NumFlux)		$\alpha_s$ (EFlux)	
	Variable	Value	Variable	Value
Electron Diffuse	$\alpha_{1,e}$	$0.055 \pm 27.4\%$	$\alpha_{2,e}$	$0.224 \pm 18.3\%$
Ion Diffuse	$\alpha_{1,i}$	$0.038 \pm 13.8\%$	$\alpha_{2,i}$	$0.207 \pm 23.8\%$
Monoenergetic	$\alpha_3$	$0.741 \pm 4.48\%$	$\alpha_4$	$0.995 \pm 4.68\%$
Broadband	$\alpha_5$	$0.244 \pm 4.70\%$	$\alpha_6$	$2.247 \pm 5.12\%$

**Table 2.** Finalized value of  $\alpha_s$  for auroral number and energy flux determined for each precipitative source.

for both electron and ion diffuse precipitation were adjusted based on the median of ratio between NOAA-DMSP and MAGNIT fluxes, and between OV Prime and MAGNIT fluxes. Scaled comparisons of MAGNIT fluxes in Fig. 4(ii-a) and 4(ii-b) show reasonable agreement with NOAA-DMSP hemispheric power values during extreme driving for both electrons and ions. However, in Fig. 4(ii-c) and 4(ii-d), NOAA-DMSP estimates lower number fluxes for stronger driving conditions, during which MAGNIT fluxes match well with OV Prime values.


The finalized values of  $\alpha_s$  (and their associated uncertainties) are presented in Table 2. The associated uncertainty is derived using the standard deviation in  $\alpha_s$  from each run, while the median value has been used as the finalized value for simulations. The uncertainty in  $\alpha_s$  is bound to impact the final modeling results, which would have far-reaching impacts on our space weather results. Investigation of this impact is beyond the scope of the present study, and will be researched in further detail in a future study. The finalized values were retrofitted into MAGNIT to balance each source of precipitation during run-time. This procedure has been applied for the simulation of the *Galaxy15* event described in Section 4.

## 4 Event Simulation

The *Galaxy15* Event is a prominent space weather interval (e.g. Allen, 2010) which has been investigated by multiple studies (e.g. Pulkkinen et al., 2013; Chen et al., 2015; Keese et al., 2014; Anderson et al., 2017; Welling et al., 2017; Mukhopadhyay et al., 2020), and spanned from April 5, 2010 00:00 UT until April 6, 2010 23:59 UT. The event was driven by a fast interplanetary coronal mass ejection (ICME; e.g. Cane & Richardson, 2003) that caused significant dipolarization during the main phase of the storm (e.g. Connors et al., 2011), resulting in a prolonged recovery period (e.g. Möstl et al., 2010). Intense auroral activity along with multiple substorms were observed during this period (Clilverd et al., 2012; Loto'aniu et al., 2015), resulting in the event being reclassified as a supersubstorm (e.g. Nishimura, Lyons, et al., 2020). Figure 5(a - e) display IMF  $\vec{u}$ ,  $\vec{B}$ , number density and temperature, that have been used as input conditions to drive this SMWF run. The solar plasma parameters were obtained from instruments aboard the Advanced Composition Explorer (ACE). Unlike the idealized cases in Section 3, the event was simulated without the necessity of a dedicated preconditioning period. Separate SWMF simulations were run with the empirical conductance models RLM and CMEE to compare model-model auroral dynamics.

Panels 5(f - j) show simulated space weather quantities against respective observations. The quantities compared here include the space weather indices Kp and Sym-H, and ionospheric quantities iFACs and CPCP. Upward and downward iFACs have been accounted for separately in Figures 5(h) and (i) respectively. MAGNIT simulations of Kp and Sym-H exhibit a MAPE value of 8.73% and 19.67% respectively. In contrast, the RLM and CMEE simulation exhibit higher values of MAPE for both Kp (RLM - 11.02%, CMEE - 14.29%) and Sym-H (RLM - 30.46%, CMEE - 40.36%). Modeled predictions for both quantities show reasonable agreement throughout the event, with the exception of modeled Sym-H during the event peak and early recovery period which was overpredicted. All models show excellent agreement with observed upward and downward iFACs. MAGNIT exhibits a  $\sim 16\%$  MAPE value for both up- and downward FACs, followed closely by RLM and CMEE which exhibit a MAPE of  $\sim 16.4\%$  and  $\sim 18.6\%$  respectively. This is partly due to the numerical grid resolution of the global MHD domain, which plays a dominant role in defining FAC magnitude and structure (Ridley et al., 2010; Wiltberger et al., 2017; Welling, 2019; Mukhopadhyay et al., 2020). Comparison of the CPCP to either source of observations do not yield meaningful conclusions, as SuperDARN is clearly underestimating the CPCP while AMIE overpredicts the value (Gao, 2012; Mukhopadhyay, Jia, et al., 2021). So, it is desirable to be between the two, i.e., have a low value of EP. The modeled CPCP by the MAGNIT run had an EP value of  $\sim 27\%$ . In contrast,

## April 5-7 2010 “Galaxy15” Event - SWMF Input Parameters and Results



**Figure 5.** Input conditions and global results from the SWMF run of the *Galaxy15* Event. (a) IMF  $U_x$ , (b)  $U_y$  and  $U_z$ , (c) IMF  $\vec{B}$ , (d) number density  $\rho$ , (e) particle energy. (f) Kp, (g) Sym-H, (h) Upward integrated field aligned currents (iFACs), (i) Downward iFACs, and (j) cross polar cap potential (CPCP) simulated by MAGNIT (in blue), RLM (in red), and CMEE (in green) against observations (in black; Kyoto Observatory for Kp and Sym-H, AMPERE FACs for iFACs). In (j), two observation-derived sources - AMIE and SuperDARN - have been used. For (f-i), the MAPE between the modeled values and observations have been provided. For (j), the EP is provided.

the RLM and CMEE runs had an EP of  $\sim 33\%$  and  $\sim 71\%$  respectively. Since FACs and CPCP stand to be affected the most by a change in auroral precipitation a more exhaustive examination that quantifies modifications in the particle drift velocities and currents could be conducted, but is not the focus of the present study. Subsequent investigations will review the ionospheric electrodynamics results in greater detail.

### 4.1 Balance of Precipitation

Figure 6 compares total and source-wise hemispheric power from MAGNIT against multiple sources of both observations and observation-derived estimates. Fig. 6(a) compares MAGNIT total hemispheric power against observations by DMSP SSUSI, simulated results from OV Prime, FTA and the four empirical models by Brautigam et al. (1991), Ahn et al. (1983), Lu et al. (1998) and Østgaard et al. (2002) (hereinafter referred to as *Brautigam91*, *Ahn83*, *Lu98*, *Ostgaard02* models, respectively). The latter four models have been bundled together to form a light-blue band in the subplot. Solar wind inputs drive both OV Prime and *Brautigam91* models, while Kyoto-observed AE/AL values are used to drive the FTA model and the remaining empirical models. Since several models in this plot were designed to derive hemispheric power from electrons only



(e.g. FTA), the electron contribution from MAGNIT has been plotted as a dot-dashed black line alongside the total contribution. Overall, modeled precipitation by MAGNIT exhibits reasonable agreement with both observed and derived estimates. The hemispheric power peak of 225.2 GW estimated by MAGNIT compares well against the observed peak of 218.3 GW from DMSP, albeit at different times. This is most likely due to the difference in time cadence between both datasets, with DMSP being hourly-averaged and MAGNIT having a cadence of one minute. Modeled HP are greater than OV Prime estimates during the storm peak-time and remain larger except for a short interval during April 6, 2010 04:00 UT to 07:31 UT. Despite this, MAGNIT electron precipitation matches well against the total OV Prime estimates during the long recovery period of the storm, exhibiting an aggregate MAPE value of 24.4%. MAGNIT estimates also have good agreement with the FTA model during the main impulse of the event, but do not match during the period preceding the storm and the recovery period when FTA predicts a larger energy flux deposition. MAGNIT exhibits an 34.2% EP when compared against the range of values formed by OV Prime and FTA, denoting good agreement. Additionally, the modeled values reasonably agree against estimates from the four empirical models (EP 29%), with majority of the overprediction occurring during the main impulse of the storm.


In Panel (b), MAGNIT predicts a larger value of electron diffuse power in comparison to OV Prime, resulting in a median absolute error of 19.95 GW (MAPE = 65.8%). This is expected for two reasons - (1) The finalized  $\alpha_s$  values that regulate the electron diffuse flux in MAGNIT allow for a much higher diffuse flux value during moderate-to-extreme driving conditions (see Figure 4(ii-a)), and (2) OVATION Prime is most likely underpredicting diffuse precipitation during the solar wind enhancement (e.g. Newell et al., 2014). With a maximum peak of 125.68 GW, electron diffuse precipitation accounts for a median 51.4% of the total contribution to the hemispheric power. This causes the magnitude differences between the two models, as seen in Fig. 6(a).

In Panels (c) and (d), monoenergetic and broadband powers display good agreement with the OVATION Prime values, except during the main impulse phase and early recovery period (08:35 UT to ~20:00 UT) of the event, when monoenergetic precipitation is overpredicted while broadband is underpredicted. However, both models predict the double peak in flux values centered around 10:45 UT and 13:30 UT. Discrete (monoenergetic + broadband) hemispheric power in the post-noon pre-dusk (12 - 18 MLT) sector is compared against OV Prime and the empirical relationship given by Korth et al. (2014) in Panel (e). Both models show reasonable agreement with the *Korth14* model and with each other during the latter half of the recovery period of the event. However, significant differences during the event peak are also observed: both MAGNIT and OV Prime estimate a greater discrete precipitation during this time period, compared to the *Korth14* model.

In Panels (f) and (g), the total electron and ion power from DMSP, OV Prime and *Brautigam91* indicate good agreement for electron precipitation, but show a significant disconnect in ion precipitation, with MAGNIT overpredicting the ion fluxes by over 2.5 times. This also increases the contribution of ion precipitation to the total hemispheric energy flux, to a median 22.4%. It is likely that the observed estimates are underpredicting the energy flux due to ions, since both DMSP and OV Prime do not observe particles with average energies >30 keV (e.g. Newell et al., 2009). A closer examination of ion precipitation is necessary, which will be pursued in a future study.

Finally, Panel (h) compares MAGNIT HP against predictions from the empirical conductance models, RLM and CMEE, and NOAA-DMSP, FTA, OV Prime and the four empirical models combined (light-blue band). RLM and CMEE predict higher fluxes than MAGNIT estimates. This is most probably because of the models' usage of empirical adjustments to increase conductances in regions of high FACs. Both RLM and CMEE

## MAGNIT Comparisons : Hemispheric Power Comparisons



**Figure 6.** Comparison of hemispheric power estimated by MAGNIT against multiple observational-derived estimates. Comparison of (a) Total hemispheric power estimated by MAGNIT (black; solid line denotes all sources, dot-dashed denotes electron sources), NOAA-DMSP (red), OV Prime (blue), FTA (green) and four empirical models (lightblue band). (b - d) Electron diffuse, monoenergetic and broadband precipitation by MAGNIT (black) against OVATION Prime (deep blue). (e) Discrete precipitation in the dusk-noon sector by MAGNIT (black) against OV Prime (blue) and Korth et al. (2014) (red). (f) Total electron precipitation, and (g) ion precipitation - MAGNIT (black) vs. OV Prime (blue), NOAA-DMSP (red) and Brautigam et al. (1991)(turquoise) (h) Hemispheric power by RLM and CMEE vs. MAGNIT compared with observation-derived estimates (lightblue band).

overpredict the hemispheric power, with CMEE values reaching  $\sim 500$  GW during the event peak.

Figure 7 compares MAGNIT flux patterns and aggregate contributions from each source against OV Prime. In this figure, panels (a) and (b) compare auroral patterns due to diffuse (ion + electron) and discrete (monoenergetic + broadband) sources respectively in the Northern hemisphere for two time intervals. The first time interval, Epoch 1, is chosen at April 5, 2010 09:10 UT, during the storm onset. The second time interval, Epoch 2, is chosen at April 6, 2010 16:12 UT during the recovery period. In Panel 7(a - i), modeled diffuse precipitation matches the magnitude and location of precipitation predicted by OV Prime. Diffuse sources in MAGNIT account for 37% of the total contribution at Epoch 1, indicating a larger contribution by the discrete sources during this epoch. The inverse is true for OV Prime, where diffuse sources contribute toward 60.7% of the total precipitation making them the dominant contributor. In Panel 7(b - i), modeled discrete energy flux is dominated by the monoenergetic source, which is indicated by the presence of strong auroral activity in regions of upward FACs (indicated by green dotted line). These regions are discontinuous, but contain strong precipitation in both dawn- and dusk-ward sectors. Broadband precipitation is not as strong, and therefore contributes to only 11.3% of the total energy flux. By contrast, OV Prime estimates a wide oval of energy flux spanning the  $70^\circ$  -  $60^\circ$  MLat at 00 MLT, with no visible differences demarcating monoenergetic and broadband precipitation.


Figure 7(a - ii) and (b - ii) display auroral patterns during Epoch 2. In Figure 7(a - ii), MAGNIT predicts a larger diffuse flux contribution in comparison to OV Prime, resulting in a dominant 74% of the total contribution. MAGNIT's energy flux pattern expands beyond  $60^\circ$  MLat, with high fluxes spanning the entire nightside. By contrast, OV Prime predicts a minimal increment in the fluxes with no noticeable expansion in the aurora. In Figure 7(b - ii), modeled discrete precipitation is still dominated by monoenergetic precipitation. But the magnitude of fluxes have reduced significantly. Diffuse fluxes from OV Prime are estimated to be more pole-ward than MAGNIT. Both models predict a discrete auroral power of around  $\sim 34$  GW. However, their percentage contributions are significantly different in each model, with OV Prime estimating a 45% discrete contribution, while MAGNIT predicts a much lower 29% contribution. MAGNIT observes strong monoenergetic precipitation in both dusk and dawn sectors. This is likely because the model predicts strengthened R2 FACs due to the coupling with RCM.

Figure 7(c-g) compare median values of hemispheric power, number flux and average energy for the four sources against OV Prime estimates. Subplots 7(c) and (d) compare the median values of hemispheric power and number flux from each source and the associated uncertainty in the data (indicated by the translucent spread around the marker). Here, the uncertainty is measured using the standard deviation in the data for each source. The figure shows that MAGNIT estimates a higher diffuse power than OV Prime. Modeled estimates for electron and ion diffuse precipitation has a median value of 52.38 GW and 22.82 GW respectively. This is greater than the OV Prime estimates, which predict 32.73 GW from electron diffuse and 6.79 GW from ion precipitation. Conversely, estimates of discrete power contributions by both models are similar. While electron diffuse number fluxes are higher in MAGNIT, both models match ion number flux estimates. OV Prime predicts a higher number flux from their monoenergetic source. However, both models reasonably agree on the median number flux from broadband precipitation.

Subplots 7(e) and (f) compare the median contributions by each source to the total hemispheric power and number flux respectively. The percentage contributions are computed using Equation 11. The figures show higher diffuse contributions by MAGNIT, which subsequently reduces contributions by the discrete sources. In Subplot 7(f), ion precipitation accounts for the smallest contributions ( $\sim 2\%$ ) to the total number flux. Electron diffuse precipitation accounts for a larger percentage of the total number flux than OV Prime prediction, while monoenergetic contribution reduces. Broadband con-



## MAGNIT vs. OVATION : Flux Patterns & Contributions



**Figure 7.** MAGNIT vs. OVATION Prime - Balance of fluxes and comparison of auroral patterns. (Top) Dial-plot comparisons of (a) diffuse energy flux and (b) discrete energy flux at two distinct epochs during the Galaxy15 Event. (Bottom) Source-wise comparison of (c) hemispheric power (in GigaWatts), (d) hemispheric number flux (in particles), (e) power contribution (in percent), (f) number flux contributions (in percent), and (g) average energy (in kilo-electronVolt).

tributions for both models show reasonable agreement with each other. Subplot 7(g) compares average energies of particles from all sources. MAGNIT exhibits stronger average energies for ion (29.64 keV) and monoenergetic precipitation (5.17 keV), deviating sharply from OV Prime estimates (8.96 keV for ions, 2.25 keV for monoenergetic populations) that show a 230% and 130% increase in energies of each respective source. In comparison, MAGNIT's electron diffuse (5.4% increase) and broadband (10.1% decrease) energies are similar to that predicted by OV Prime. MAGNIT estimates a more energetic ion population, compared to OV Prime. This is because of two reasons - (1) MAGNIT estimates lower ion number fluxes and higher energy fluxes, and (2) OV Prime underpredicts the ion average energy (Newell et al., 2009). While higher energies are exhibited by monoenergetic precipitation as well, this is most likely due to the model's dependence on strong FACs to derive bulk of its monoenergetic precipitation, which is a function of activity.

#### 4.1.1 Comparison of Auroral Flux Patterns


Modeled auroral patterns in the Northern hemisphere have been compared against DMSP SSUSI observations in Figure 8. Fig. 8(a) compares energy flux observations by the DMSP F17 spacecraft in the first column against simulations by (chronologically) MAGNIT, RLM and CMEE. Each row indicates a specific time epoch. The first row indicates auroral fluxes at April 5, 2010 at 07:24 UT, before the sudden commencement of the Galaxy15 Event. During this time interval, flux pattern from DMSP shows low auroral activity with limited expansion of the oval. This observation is well reproduced by MAGNIT which match the magnitude of DMSP's energy flux. However, MAGNIT predicts an expanded auroral oval, with its equatorward boundary reaching lower 60°s MLat. Both RLM and CMEE overpredict the magnitude of energy flux, displaying a strong auroral oval with minimal auroral expansion. However, both models predict the auroral oval in similar latitude as DMSP.

Epoch 2 takes place immediately following the main impulse of the event at 10:48 UT. At this time, DMSP observations indicate a significant increase in auroral precipitation in the dawnward sector, with similar but comparatively limited enhancements in the duskward sector. The oval has expanded substantially, with the dawnward peak appearing at 67° - 66° MLat and the duskward peak at 65° MLat. The dawnward flux precipitation is broader and stronger than the duskward sector, most likely caused by heightened electron diffuse precipitation. The enhancement of auroral precipitation is also captured by MAGNIT, which exhibits a higher magnitude and a distinct auroral expansion. Flux magnitudes from MAGNIT compare well against the DMSP observations matching peak regions of precipitation. However, MAGNIT overpredicts the auroral expansion by diffuse sources, estimating an equatorward dawnward peak. The flux peak in the duskward sector matches well with the DMSP observations. This is probably because this stretch of precipitation is being driven solely by monoenergetic precipitation, which follow the upward R1 FACs in the region. MAGNIT also exhibits distinct meso-scale structures poleward of the auroral oval. Both RLM and CMEE do exhibit similar flux levels as DMSP, but do not show any significant expansions in the auroral oval. This is expected from their empirical design, and are limited from expanding the aurora beyond 60° MLat (see Section 3.1 in Mukhopadhyay et al., 2020 for details).


Epoch 3 takes place during the long recovery period of the storm at 17:36 UT on April 5 2010. During this time interval, DMSP observes a broad dawnward flux peak spanning 70° to 60° MLat, and a narrow duskward peak bordering 70° MLat. Despite its capability to predict correct magnitudes, MAGNIT overpredicts the latitudinal extent of the dawnward precipitation. Auroral fluxes predicted by RLM (Column iii) and CMEE (Column iv) exhibit high energy fluxes and distinct FAC-driven structures during these two epochs without much auroral expansion.

## Comparison against DMSP SSUSI

### (a) Energy Flux Patterns



### (b) Average Energy Patterns



**Figure 8.** Comparison of (a) total energy flux and (b) average energy patterns in the Northern hemisphere at discrete time intervals spanning the *Galaxy15* Event. (Column-wise) Dial plots display observations from (left to right) DMSP SSUSI (Column i), and simulated results from SWMF simulations driven using MAGNIT (Column ii), RLM (Column iii) and CMEE (Column iv) at three distinct time epochs. Dotted lines in each dial plot indicates span of SSUSI coverage.

Figure 8(b) compares the average energy for the aforementioned datasets for the same epochs. Unlike part (a), only electron-driven precipitation (electron diffuse, monoenergetic and broadband) have been used to compute the energies in MAGNIT. This was done because the energies from ion precipitation in MAGNIT were much higher than the observational limit of DMSP ( $\geq 25$  keV). DMSP observations and results from MAGNIT display significant morphological similarities. Similar to the previous part, MAGNIT exhibits an extended auroral oval in comparison to DMSP observations, but is able to match average energy magnitudes. By contrast, both RLM and CMEE are unable to produce resembling average energy patterns. The average energy patterns in both are characterized by a high energy region around the geomagnetic poles, with slight reduction in energies as one moves equatorward. This is a drawback in both these models due to their usage of FACs to compute conductances directly, and not fluxes. The average energy, therefore, is a byproduct of a reverse-*Robinson* relationship, where the FAC-derived conductances are converted to energies using the inverse of the relationship given in Robinson et al. (1987) (see Mukhopadhyay et al., 2020 for further details), rather than a physical manifestation of particle energies like in MAGNIT.

## 4.2 Magnetospheric Feedback

Due to its dynamic coupling with the magnetosphere, MAGNIT allows for better reception of magnetospheric feedback into their calculation of auroral fluxes and ionospheric conductance. Figures 9 and 10 show an example of this process. Figure 9(a-i) compares the contribution by each source of precipitation during the *Galaxy15* event. For a majority of the event, the source-wise contributions take a near-constant value, with electron diffuse precipitation providing 51% of the total energy flux. Ion and monoenergetic precipitation contributions are each about half of the electron diffuse contribution, with a median contribution of 22.6% and 18.6% each. The remaining contribution is provided by the broadband source, which provides for a mere 6.5% of the total contribution making it the smallest source of energy flux. However, these contributions are severely distorted during the event peak where the discrete sources overtake diffuse contributions peaking at a combined contribution of 61% at 08:35 UT on April 5, 2010. The zoomed section in part (a-ii) shows this time duration in further detail, and identifies five time intervals - 08:17 ( $t_1$ ), 08:37 ( $t_2$ ), 09:01 ( $t_3$ ), 09:15 ( $t_4$ ) and 09:45 UT ( $t_5$ ) on April 5, 2010 - to investigate the driving factors of the discrete enhancement. The time duration spans the main impulse of the *Galaxy15* event during which the solar wind ram pressure increased, as shown in part (a-ii). The impact of magnetospheric dynamics and subsequent ionospheric interactions during a solar wind enhancement have been previously analyzed in multiple numerical studies like Yu et al. (2010), Ozturk et al. (2018), and more recently in Welling et al. (2021). In this study, we investigate these dynamics in regards to changes in auroral precipitation.

In Figure 9(b), the response of four quantities - (row-wise) FACs, Joule Heating, monoenergetic energy flux and broadband energy flux - have been displayed for the five time intervals. At  $t_1$ , FAC patterns follow the standard southward  $B_z$  model with strong R1 FACs and relatively weaker R2 FACs, and are symmetric about the noon-midnight meridian with an integrated total current of 4.51 MA. Joule heating in the second row is computed using the Pedersen conductivity and electric field (e.g. Rastätter et al., 2016), and appear in the same region as the FACs exhibiting a maximum flux of 13.08 mW/m<sup>2</sup>. Since the model is not coupled to a dedicated ionosphere-thermosphere model, the Joule heating does not incorporate contributions from the neutral wind. The third and fourth rows display monoenergetic and broadband energy flux. Both FACs and Joule heating are the dominant drivers of these two respective sources. Monoenergetic precipitation, which follows the KFL relationship, is strong only in regions of upward FACs and accounts for 22.5% of the total hemispheric power. Similarly, broadband enhancements are derived as an empirical function of the Joule heating, and contributes to 23.3% of the total power.

Time interval  $t_2$  takes place 7 minutes after the main impulse of the event that leads to ram pressure enhancement in the solar wind. This interval is readily characterized by the strengthening of FACs and Joule heating patterns. FACs are still symmetric about the noon-midnight meridian, but exhibit significant enhancement in R1 currents and stronger R2 currents.  $t_2$  observes a doubling of integrated total current to 10.1 MA. A significant dayside peak emerges in the Joule heating pattern, with the maximum flux surging to  $47.3 \text{ mW/m}^2$ . Correspondingly, the discrete fluxes react to these changes with an enhancement in precipitation. Monoenergetic flux rises from 3.9 GW at  $t_1$  to 9.8 GW at  $t_2$ , with a significant contribution in the R2 FACs (dawnward) sector in addition to the upward R1 FACs in the dusk sector. Broadband precipitation increases by 3.35 times, becoming the dominant contributor to auroral precipitation. Discrete contributions account for 60.5% of the total precipitation, in contrast with contributions during the rest of the event.

During  $t_3$  (24 minutes after  $t_2$ , 31 minutes after the pressure enhancement), R1 FACs start expanding in addition to becoming more enhanced in magnitude. The dominant peaks expand beyond  $70^\circ$  MLat, resulting in a further expansion of the R2 currents into the upper  $50^\circ$  MLat. The FAC pattern increasingly becomes asymmetric across the noon-midnight meridian with small scale structures connecting the R1 and R2 FACs appearing in the nightside. Two dominant peaks arise in the Joule heating in the dusk and dawnward flanks, with the duskward peak appearing in the dayside region of the peak at  $t_2$ . The dawnward peak appears in the nightside, with the pattern being symmetric about the 10-22 MLT meridian. The enhancement in monoenergetic precipitation is characterized by the strengthening of FACs, and accounts for 34.9% of the total precipitation. Despite the strengthening of FACs in both the dayside and nightside alike, the monoenergetic population is mostly concentrated on the nightside. This is most likely because the diffuse number flux is low on the dayside, i.e. there is insufficient particles in the dayside to accelerate. Broadband precipitation is concentrated in the Joule heating peaks in the two flanks of the northern hemisphere. However, due to strengthening of the other sources of precipitation, the contribution by broadband precipitation reduces to 15.3%.


Time intervals  $t_4$  and  $t_5$  display conditions at 09:15 UT (14 minutes after  $t_3$ , 45 minutes after the pressure enhancement) and 09:45 UT (44 minutes after  $t_3$ , 1.25 hours after the pressure enhancement). During these time intervals, the FAC expansion observed during  $t_3$  reaches its furthest point and plateaus, with R1 current peaks extending from the upper  $70^\circ$  MLat to mid  $60^\circ$  MLat while R2 currents reach down to mid- $50^\circ$  MLat. Both R1 and R2 FACs are further strengthened and become increasingly asymmetrical. Joule heating patterns have a similar two-peak configuration as seen at  $t_3$ . The peaks lie in the same region as the upward FAC peaks, albeit being much broader than R2 peaks in the dawnward sector. Correspondingly, monoenergetic precipitation increases along with FAC enhancements and expansion, resulting in a total hemispheric power of 39.6 GW during  $t_4$  and 31.1 GW during  $t_5$ . Due to the increase in diffuse flux during these times in the dayside sector, monoenergetic precipitation extends into the dayside in both dawn and duskward sectors. Broadband precipitation follows the two-peak configuration of the Joule heating pattern, and results in a total power of 15.4 GW at  $t_4$  and 11.3 GW at  $t_5$ . Contributions by discrete sources to the total precipitation drops down from 60.5% during  $t_3$  to 41.5% during  $t_4$  and 26.3% during  $t_5$  of the total precipitation, indicating significant enhancement in diffuse precipitation.

Figure 10 describes the impact of the solar wind pressure enhancement on diffuse precipitation. The MHD single-fluid pressure and temperature have been mapped to the ionospheric grid for clarity, and have been displayed in the first and second rows respectively for the same time intervals as in Figure 9(b). Since both these quantities drive diffuse precipitation in MAGNIT, the energy flux from electron and ion precipitation have been plotted in the third and fourth rows. Comparison of the first row across the five time intervals indicate a steady enhancement in the nightside pressure. This enhance-




## Impact of M-I Feedback on Discrete Precipitation

### (a) Comparison of Individual Contributions




### (b) Drivers & Resultant Fluxes of Discrete Precipitation



**Figure 9.** (a) Comparison of contributions by all sources during the Galaxy15 event, with five time intervals highlighted. (b) Comparison of quantities pertaining to discrete precipitation - (i) field-aligned currents, (ii) Joule heating, (iii) monoenergetic energy flux, and (iv) broadband energy flux. Despite the uniformity in the colorbar scale, note that the colorbar range for  $t_1$  are different from other times.

### Impact of M-I Feedback on Diffuse Precipitation



**Figure 10.** Comparison of quantities pertaining to diffuse precipitation - (i) MHD plasma pressure mapped onto the ionospheric domain, (ii) MHD plasma temperature mapped onto the ionospheric domain, (iii) electron diffuse energy flux, and (iv) ion diffuse energy flux - during each time interval identified in Fig. 9(a).

ment in the pressure also brings the pressure peaks in nightside closer to Earth, resulting in an equatorward expansion of the nightside peak on the ionospheric grid. The buildup in nightside pressure is gradual, as can be seen during  $t_1$ ,  $t_2$  and  $t_3$ , and is not slower than the ramp-up in FACs. This is expected, since asymmetric currents that drive FACs in the system build up much quicker than symmetric pressure used here (e.g. Liemohn et al., 2015). Before the solar wind enhancement, the nightside pressure at  $t_1$  is expected to be less. Post-ram pressure enhancement, at  $t_2$ , the increase in pressure is demarcated by the formation of an oval with a peak in the midnight-dusk sector. However, the pressure increases steadily over the next 24 minutes, culminating in a sharp nightside peak with a band spanning the auroral region. The pressure peak expands equatorward from their previous location at  $t_2$ , and quadruples in magnitude. Intervals  $t_4$  and  $t_5$  exhibit further enhancement and expansion of the nightside pressure peak. At  $t_5$ , the pressure peak moves further equatorward reaching the lower  $50^\circ$  MLat. A similar buildup of energy is observed in the second row, when comparing particle temperatures. Contrary to the pressure distribution, the MHD temperature distributions during  $t_3$ ,  $t_4$  and  $t_5$  exhibit strong meso-scale dynamics poleward of the auroral peak formed through the mapping. The dynamic structures in the temperature distribution change with the march in time, occasionally strengthening the peaks in the oval region, like in  $t_3$ .


The electron and ion diffuse energy fluxes are compared in the last two rows of Figure 10. During  $t_1$ , electron and ion diffuse precipitation account for 37.8% and 16.5% of the total energy flux respectively, with the ion precipitation accounting for the lowest contribution. However, energy fluxes from both these sources are comparatively lower and exhibit a diffuse band of precipitation centered around the nightside pressure peak (for ions) and its mirrored location (for electrons). With the solar wind pressure enhancement, the fractional contribution of the diffuse sources reduce, since discrete fluxes enhance faster resulting in the electron diffuse sources to contribute to 27.5% of the total flux during  $t_2$ . At  $t_3$ , the pressure peak in the nightside is sufficiently strong to raise the diffuse hemispheric power to 48.4 GW, accounting for nearly half of the total precipitation. The electron diffuse precipitation at  $t_3$  is characterized by a sharp peak in the midnight-dawnward sector, with the auroral oval expanding beyond  $60^\circ$  MLat. The ion precipitation mirrors this configuration resulting in a total hemispheric power of 14.7 GW. During  $t_4$  and  $t_5$ , both diffuse precipitation enhance in magnitude and expand further equatorward. Electron diffuse precipitation becomes the dominant contributor to the total precipitation, accounting for 51.4% of the total precipitation by  $t_5$ . Contributions by ion precipitation is similarly enhanced, contributing 22.4% of the total precipitation greater than either discrete sources.

### 4.3 Impact on Ionospheric Conductance

Figure 11 presents detailed comparisons of contributions to the ionospheric conductance by each source of precipitation. Figure 11(a) and (b) present individual contributions from each source to the Hall and Pedersen conductance during the *Galaxy15* event. These values are calculated by Equations 12 for monoenergetic, electron and ion diffuse contributions, and by Equation 13 for broadband contributions. Electron diffuse precipitation is the largest contributor to both Hall and Pedersen conductance, accounting for a contribution of  $\sim 34\%$  to both types of conductance. This is closely followed by ion precipitation which accounts for 31% of the Hall conductance, while being the third largest source of Pedersen conductance. Monoenergetic precipitation accounts for 23.3% of the Hall conductance. However, this proportion dwindles down to a mere 5.77% contribution to the Pedersen conductance. The opposite is true for broadband precipitation which accounts for 9.4% of the Hall conductance, and 31.4% of the Pedersen conductance making it the second-largest source to Pedersen conductance after electron diffuse precipitation. The disparity in contributions for the two discrete sources could be explained by the nature of these flux populations and how they interact with the *Robinson* relation when converted into conductance. Monoenergetic precipitation is generally the more



### Resultant Contributions to Hall & Pedersen Conductance



**Figure 11.** Individual contributions to perpendicular conductance - (a) Comparison of individual resultant contributions by each source of precipitation to the Hall Conductance, and (b) Pedersen conductance. (c) Comparison of total and resultant contributions by broadband precipitation to Hall and Pedersen conductance. (d) Polar patterns of total conductance on April 5, 2010 at 08:55 UT, simulated using MAGNIT, RLM and CMEE.

energetic source of electron precipitation. This leads to a higher Hall conductance value through the Robinson relationship, as the Hall conductance term is directly proportional to the average energy  $\bar{E}^{0.85}$ . Broadband precipitation is the least energetic population, and therefore possesses a low average energy but high number flux. Since the *Robinson* relationship defines the Pedersen conductance as being directly proportional to the square root of the energy flux, this increases the contribution of the broadband source. Furthermore, the linear addition of the broadband contribution in Eq. 10 significantly raises its resultant contribution, leading to aforementioned disparity.

The enhancement in the broadband contribution is further explained in Figure 11(c). Here, the contribution by broadband conductance to the total auroral conductance is computed by two methods - (1) as a fraction of the total conductance (Total contribution; computed using Equation 11), and (2) actual contribution due to its linear sum (Resultant contribution; computed using Equation 13). Comparison of both contributions indicate that the resultant contribution of broadband conductance to the total auroral conductance is much higher than expected. The resultant contribution of the broadband source is nearly  $\sim 3$  times their numerical value due to the linear addition. Driving conditions during the early phase of the event causes fluctuating contributions that can contribute to nearly 71% of the total auroral Pedersen conductance. The heightened contribution of broadband precipitation naturally postulates an important role played by this source of precipitation in ionospheric electrodynamics.

Figure 11(d) compares dial plots of Hall and Pedersen conductance patterns simulated by MAGNIT, RLM and CMEE. The conductance patterns are simulated at 08:55 UT on April 5, 2010. Both RLM and CMEE provide higher conductances in the auroral region in comparison to MAGNIT. In MAGNIT, the auroral conductance is mostly concentrated on the night, with minor flanks of the dawnward and duskward fluxes reaching into the dayside sector. By contrast, significant dayside precipitation is visible in both RLM and CMEE, which exhibit a thicker auroral oval. Despite this, MAGNIT exhibits a more expanded auroral oval, with strong contributions by the monoenergetic and electron diffuse sources. RLM follows the FAC pattern closely, creating a strong R1 FAC systems with discontinuities in the conductance where the FACs change polarity. This is similar in CMEE, where the magnitude of conductance is much higher leading to a more dynamic auroral oval featuring several FAC-driven structures in polar regions.

## 5 Discussion

The introduction of MAGNIT to the SWMF environment is a significant step forward towards computing realistic precipitation in the global model. First, the incorporation of advanced coupling mechanisms between BATS-R-US and RIM allow for the computation of multiple sources. These mechanisms establish a solid roadmap for future advancements in computing fluxes and conductances in RIM, that may involve further physics-based couplings with MHD or ring current models. Second, computation of precipitation in MAGNIT is far more realistic in comparison to its empirical predecessors. Both RLM and CMEE estimate higher energy fluxes and incorrect average energies in the auroral region. MAGNIT outperforms both models when predicting energy flux (as shown in Fig. 6h), and results in relatively lower but sharper conductance contributions in the auroral region. The use of MHD variables to compute auroral fluxes also means a flexible activity-driven oval expansion, eradicating the problem observed in the empirical models during active periods (see Section 3.1 in Mukhopadhyay et al., 2020). Third, the ability to compute four individual sources of precipitation allows for the quantification of source-wise contributions to other ionospheric variables. Investigations quantifying the individual impacts of each source on field aligned currents, ionospheric potential and E-fields will soon be presented in a subsequent manuscript. Fourth, this adds to SWMF's capability to quantify the dependence of space weather results on distinct auroral precipitation types (Vandegriff et al., 2020, 2021). Finally, results simulated with MAGNIT

show good agreement with observations and state-of-the-art empirical models. The magnitude of auroral precipitation agree well with both DMSP-NOAA and OV Prime. Further comparisons of more events are underway, and will be presented in future investigations.

Note that there are several modeling caveats to this study. The auroral flux patterns formed in MAGNIT simulations, as seen in Fig. 7(a) and Fig. 8(b), show that the expanded oval of the diffuse aurora is more equatorward than those in observations. In both OV Prime (in reference to Fig. 3) and DMSP (in reference to Fig. 8), the dawn-side peak of auroral precipitation lies in the middle of the 60 - 70° MLat range, while the dawnward sector in MAGNIT is characterized by a much more expanded diffuse oval peaking between 55° and 65°. This is most likely because MAGNIT computes diffuse precipitation using a Maxwellian distribution computed from the single-fluid plasma temperature provided by BATS-R-US. This is problematic, as the MHD temperature is that of ions, and is converted into electron temperature by assuming a 1:5 ratio between electrons and ions. Ion temperature and pressure in global MHD is generally closer to Earth, since plasma sheet ions penetrate deeper into the inner magnetosphere than electrons (e.g. Ejiri et al., 1980; Ganushkina et al., 2000), especially during active periods (Yang et al., 2011; Gkioulidou et al., 2015). When mapped onto an ionospheric grid, these values are situated more equatorward of electron-associated pressure and temperature values. An example of this can be seen in Jordanova et al. (2012) where the difference between ion and electron flux patterns in the nightside shows 15keV ion flux peaks being closer to Earth than for electrons.

Usage of a dedicated electron temperature in the MHD equations would result in a nightside pressure peak that is farther away from the ion pressure peak. This would automatically result in an electron precipitation pattern that is more poleward than current estimates. The dedicated computation of multi-fluid (e.g. Glocer et al., 2009) and multi-species (e.g. Welling & Ridley, 2010) MHD pressure and temperature is possible through BATS-R-US. An alternative solution to improve diffuse calculations is to use particle fluxes computed by the ring current model. Yu et al. (2016) demonstrated the computation of electron fluxes using a coupled version of SWMF with the RAM-SCB model. An extension to this work was performed by Perlongo et al. (2017), where the RAM-SCB model was driven by empirical waves to compute ionospheric conductance in GITM. More recently, work by Lin et al. (2019) and Lin et al. (2021) have sought to use RCM-derived electron diffuse fluxes and MHD-computed monoenergetic fluxes to compute the resultant auroral precipitation in the LFM MHD model. The use of a ring current model also results in more accurate computation of fluxes, since the models are able to account for pitch angle distributions. Recent work by Kang et al. (2019) have shown the incorporation of wave-induced diffuse precipitation with a dedicated coupling between the ring current model CIMI and BATS-R-US. Work towards incorporating such solutions to provide MAGNIT with a more accurate value of diffuse precipitation will increase our physical understanding of auroral dynamics, and is planned for future work.

Computing monoenergetic fluxes using the KFL relationship requires an accurate knowledge of the global magnetic field, especially that in the source region and at the ionosphere. At present, MAGNIT assumes a dipole configuration (resulting in a latitudinally varying magnetic mirror ratio), with the source region for precipitating electrons in the plasma sheet (e.g. Yu et al., 2016). This simplifies the exact contribution by this source; the source region of precipitating plasma is located higher than the equatorial plasma sheet (Hatch et al., 2019). While this assumption doesn't significantly impact the resultant fluxes from the model (most likely due to the high value of  $\alpha_s$ ), it does impact the generation of mesoscale structures in the poleward and equatorward boundaries which could further impact ionospheric electrodynamics. The computation of a realistic magnetic field is possible through the field-line tracing component of SWMF, and is being currently implemented to provide a realistic magnetic field in the calculation of

monoenergetic precipitation in our model. Results pertaining to these developments will be presented in future investigative studies.

The median values of  $\alpha_s$  listed in Table 2 have been used to regulate fluxes in MAGNIT. This has significant disadvantages, since particle scattering rates (which  $\alpha_s$  emulates for diffuse and monoenergetic fluxes) are prone to modifications in the geomagnetic field leading to variable precipitation in different sectors. Furthermore, quantities like FACs and plasma pressure are highly dependent on numerical grid resolution (e.g. Ridley et al., 2010; Haiducek et al., 2017) and the inclusion of a dedicated ring current model (e.g. De Zeeuw et al., 2004). Changes in either of these factors would result in modifications in  $\alpha_s$  values for variable driving conditions. A remedy to such an issue would be to use an activity-driven MLT-wide map of  $\alpha_s$ , which tweaks auroral fluxes in regions of interest for a given upstream driving condition. In retrospect, attempts to estimate physics-based precipitation using extended couplings to the MHD model or the inner magnetospheric model is a more worthwhile solution, since it can provide a realistic physics-derived reasoning for flux outputs.

The computation of the final value of ionospheric conductance is contentious, as it tends to elevate the contribution of broadband precipitation significantly. As shown in Figure 11(c), the resultant contribution of broadband-driven Pedersen conductance jumps to a median 31% of the total. The physics associated with the summation of conductance sources (Wallis & Budzinski, 1981) is challenging to solve in a 2-D ionosphere. In this work, we have followed the example of Zhang et al. (2015) when including broadband precipitation. They state that the conductance due to broadband precipitation adds to the bottomside F-layer of the ionosphere, instead of the E-layer, where the dominant conductivity peaks in the Hall and Pedersen conductance are found (e.g. Schunk & Nagy, 2009). This is most likely because broadband precipitation exhibits lower average energy, resulting in its deposition at an upper layer. To identify this difference in altitude, the broadband-driven conductance was added linearly to the net conductance, as was done by Zhang et al. (2015). To truly solve this issue, a dedicated coupling to a 3D ionosphere-thermosphere solver is necessary. Work by Burleigh et al. (2019) has introduced novel couplings between the geospace version of the SWMF with GITM, specifically when it comes to computing ionospheric conductance realistically. The incorporation of this approach with MAGNIT-driven flux computations leads to a more realistic ionospheric feedback. Future studies by authors will feature the combination and planned usage of this modeling approach prominently in studying terrestrial (and planetary) plasma dynamics during extreme events.

## 6 Summary

A novel modeling approach was developed and used to study the April 5 - 7 *Galaxy15* Event. The model uses mapped MHD pressure and density as inputs to derive four sources of precipitation - electron diffuse, ion, monoenergetic and broadband. Precipitation from each source is regulated using empirical multipliers that ultimately define the balance between each source. The investigation of this modeling study focused on quantifying the contribution of each source of precipitation during the *Galaxy15* event, and found important results through the comparison of auroral fluxes, average energy, and ionospheric conductance to observations and empirical modeling techniques. Important findings are summarized in the following:

1. Electron diffuse precipitation is the dominant source of auroral precipitation during the *Galaxy15* event, accounting for a median 52% of the total hemispheric power. Ion diffuse and monoenergetic precipitation act as secondary sources of auroral precipitation, accounting for 22% and 19% of the total power. Broadband precipitation contributes for 7%, making it the smallest contributor to hemispheric power.

2. Auroral fluxes were converted into ionospheric conductance using empirical relationships, and used in the two-way coupling between RIM and BATS-R-US. The individual contributions of each source to the total conductance were quantified. Despite its small contribution to hemispheric power, the linear addition of broadband-driven conductances result in a 31% contribution to the Pedersen conductance, and 9% to the Hall conductance.
3. Despite the dominance by diffuse precipitation, discrete precipitation accounted for up to 61% of the total hemispheric flux during the main impulse of the *Galaxy15* event.
4. Comparison of hemispheric power against DMSP and OV Prime exhibit a higher ion energy flux in MAGNIT estimates, indicating higher average energy of ions in the model results relative to the observations. The electron diffuse precipitation is also larger than the value predicted by OV Prime.
5. Due to usage of a single-fluid ion pressure and density to derive electron diffuse precipitation, MAGNIT places the dawnward peak too far equatorward, but gets the monoenergetic peak in the dusk sector at the correct location.
6. By basing particle precipitation calculations on MHD state variables that are more tied to the drivers of ion and electron precipitation instead of only FACs, both the computation and understanding of feedback in the nonlinear M-I system are improved by this modeling approach.

## Acknowledgments

Support for this work has been provided by NASA Grants NNX17AB87G, 80NSSC18K1120, and 80NSSC17K0015, and the PREEVENTS NSF Grant AGS-1663770. A.M. gratefully acknowledges support from the Rackham Predoctoral Fellowship provided by the University of Michigan. H. K. C. gratefully acknowledges support from the NASA grants, 80NSSC18K1043, 80NSSC18K1052, 80NSSC19K0844, 80NSSC20K1670, and 80MSFC20C0019, and the NSF grants, OIA-1920965 and AGS-1928883. M.B. was supported by the Office of Naval Research. G.T. gratefully acknowledges support from the PREEVENTS NSF Grant AGS-1663800 and SWQU grant PHY-2027555.

The authors would like to acknowledge high-performance computing support from Pleiades (allocation 1815) provided by NASA's High-End Computing Capability Programme, and Cheyenne (allocation UUSL0016) provided by NCAR's Computational and Information Systems Laboratory, sponsored by the National Science Foundation. The Space Weather Modeling Framework is maintained by the University of Michigan Center for Space Environment Modeling and can be obtained at <http://csem.engin.umich.edu/tools/swmf>. AMIE Results used in this study are maintained at the University of Michigan's Virtual Model Repository (VMR; <http://vmr.engin.umich.edu/>). The authors sincerely thank Barbara Emery and the CEDAR Workshop for providing access to the NOAA-DMSP HPI dataset (<http://cedarweb.vsp.ucar.edu/wiki>), the OVATION Prime team at Johns Hopkins Applied Physics Laboratory (JHU-APL) led by Patrick Newell for maintaining open access to the 2010 version of the model available through SourceForge (<https://sourceforge.net/projects/ovationprime/>), and the AMPERE Science team and the AMPERE Science Center led by Brian Anderson for providing access to the Iridium-derived data products through their website (<http://ampere.jhuapl.edu/index.html>). The authors would also like to thank Larry Paxton and the associated JHU-APL team for maintaining DMSP-SSUSI observations for public access and usage through their website (<https://ssusi.jhuapl.edu/>), and Shibaji Chakrabarty for providing SuperDARN data for this study.

The authors would like to thank Dr. Alex Gloer, Ms. Shannon Hill, Dr. Doğan Su Öztürk, Dr. Zhenguang Huang, Dr. Katherine Garcia-Sage, Dr. Tamas Gombosi, Dr. Tuija Pulkkinen, Dr. James Slavin, Dr. Robert Robinson and Dr. George Khazanov for sharing their expertise and advice in the course of the model development and statistical analysis conducted as part of this study.



Simulation data for all runs have been made available via UM Deep Blue Data archive,  
accessible through <https://doi.org/10.7302/rtmh-x457>.

## References

- Ahn, B.-H., Akasofu, S.-I., & Kamide, Y. (1983). The Joule heat production rate and the particle energy injection rate as a function of the geomagnetic indices AE and AL. *Journal of Geophysical Research: Space Physics*, 88(A8), 6275–6287. Retrieved from <https://agupubs.onlinelibrary.wiley.com/doi/abs/10.1029/JA088iA08p06275> doi: <https://doi.org/10.1029/JA088iA08p06275>
- Allen, J. (2010). The Galaxy 15 anomaly: Another satellite in the wrong place at a critical time. *Space Weather*, 8(6).
- Anderson, B. J., Korth, H., Waters, C. L., Green, D. L., Merkin, V. G., Barnes, R. J., & Dyrud, L. P. (2014). Development of large-scale Birkeland currents determined from the Active Magnetosphere and Planetary Electrodynamics Response Experiment. *Geophysical Research Letters*, 41(9), 3017–3025. Retrieved from <https://agupubs.onlinelibrary.wiley.com/doi/abs/10.1002/2014GL059941> doi: 10.1002/2014GL059941
- Anderson, B. J., Korth, H., Welling, D. T., Merkin, V. G., Wiltberger, M. J., Raeder, J., ... Rastaetter, L. (2017). Comparison of predictive estimates of high-latitude electrodynamics with observations of global-scale birkeland currents. *Space Weather*, 15(2), 352–373. Retrieved from <https://agupubs.onlinelibrary.wiley.com/doi/abs/10.1002/2016SW001529> doi: 10.1002/2016SW001529
- Brautigam, D. H., Gussenhoven, M. S., & Hardy, D. A. (1991). A statistical study on the effects of IMF Bz and solar wind speed on auroral ion and electron precipitation. *Journal of Geophysical Research: Space Physics*, 96(A4), 5525–5538. Retrieved from <https://doi.org/10.1029/91JA00157> doi: <https://doi.org/10.1029/91JA00157>
- Büchner, J., & Zelenyi, L. M. (1987). Chaotization of the electron motion as the cause of an internal magnetotail instability and substorm onset. *Journal of Geophysical Research: Space Physics*, 92(A12), 13456–13466.
- Burleigh, M., Mukhopadhyay, A., Welling, D., Ridley, A., & Liemohn, M. (2019, dec). The Importance of Self-Consistent Conductivity in Coupling Magnetosphere-Ionosphere-Thermosphere Models. In *Agu fall meeting abstracts* (Vol. 2019, pp. SA41B–3168).
- Cane, H. V., & Richardson, I. G. (2003). Interplanetary coronal mass ejections in the near-Earth solar wind during 1996–2002. *Journal of Geophysical Research: Space Physics*, 108(A4). Retrieved from <https://agupubs.onlinelibrary.wiley.com/doi/abs/10.1029/2002JA009817> doi: <https://doi.org/10.1029/2002JA009817>
- Cash, M., Singer, H., Millward, G., Toth, G., Welling, D., & Balch, C. (2018, jul). NOAA SWPC's Operational Geospace Model Performance during Earth-Affecting Events. In *42nd cospar scientific assembly* (Vol. 42, pp. D2.3–37–18).
- Chaston, C. C., Bonnell, J. W., Carlson, C. W., McFadden, J. P., Ergun, R. E., & Strangeway, R. J. (2003). Properties of small-scale Alfvén waves and accelerated electrons from FAST. *Journal of Geophysical Research: Space Physics*, 108(A4). Retrieved from <https://agupubs.onlinelibrary.wiley.com/doi/abs/10.1029/2002JA009420> doi: <https://doi.org/10.1029/2002JA009420>
- Chen, M. W., Lemon, C. L., Guild, T. B., Keesee, A. M., Lui, A., Goldstein, J., ... Anderson, P. C. (2015). Effects of modeled ionospheric conductance and electron loss on self-consistent ring current simulations during the 5–7 April 2010 storm. *Journal of Geophysical Research: Space Physics*, 120(7), 5355–5376. Retrieved from <https://agupubs.onlinelibrary.wiley.com/doi/abs/10.1002/2015JA021285> doi: <https://doi.org/10.1002/2015JA021285>

- Chiu, Y. T., & Schulz, M. (1978). Self-consistent particle and parallel electrostatic field distributions in the magnetospheric-ionospheric auroral region. *Journal of Geophysical Research: Space Physics*, 83(A2), 629–642. Retrieved from <https://agupubs.onlinelibrary.wiley.com/doi/abs/10.1029/JA083iA02p00629> doi: <https://doi.org/10.1029/JA083iA02p00629>
- Clilverd, M. A., Rodger, C. J., Danskin, D., Usanova, M. E., Raita, T., Ulich, T., & Spanswick, E. L. (2012). Energetic particle injection, acceleration, and loss during the geomagnetic disturbances which upset Galaxy 15. *Journal of Geophysical Research: Space Physics*, 117(A12). Retrieved from <https://agupubs.onlinelibrary.wiley.com/doi/abs/10.1029/2012JA018175> doi: <https://doi.org/10.1029/2012JA018175>
- Connor, H. K., Zesta, E., Fedrizzi, M., Shi, Y., Raeder, J., Codrescu, M. V., & Fuller-Rowell, T. J. (2016). Modeling the ionosphere-thermosphere response to a geomagnetic storm using physics-based magnetospheric energy input: OpenGGCM-CTIM results. *Journal of Space Weather and Space Climate*, 6, A25. Retrieved from <http://www.swsc-journal.org/10.1051/swsc/2016019> doi: 10.1051/swsc/2016019
- Connors, M., Russell, C. T., & Angelopoulos, V. (2011). Magnetic flux transfer in the 5 April 2010 Galaxy 15 substorm: an unprecedented observation. *Annales Geophysicae*, 29(3), 619–622. Retrieved from <https://angeo.copernicus.org/articles/29/619/2011/> doi: 10.5194/angeo-29-619-2011
- De Zeeuw, D. L., Sazykin, S., Wolf, R. A., Gombosi, T. I., Ridley, A. J., & Tóth, G. (2004). Coupling of a global MHD code and an inner magnetospheric model: Initial results. *Journal of Geophysical Research: Space Physics*, 109(A12), 1–14. doi: 10.1029/2003JA010366
- Ebihara, Y., Fok, M.-C., Sazykin, S., Thomsen, M. F., Hairston, M. R., Evans, D. S., ... Ejiri, M. (2005). Ring current and the magnetosphere-ionosphere coupling during the superstorm of 20 November 2003. *Journal of Geophysical Research: Space Physics*, 110(A9). Retrieved from <https://agupubs.onlinelibrary.wiley.com/doi/abs/10.1029/2004JA010924> doi: <https://doi.org/10.1029/2004JA010924>
- Ejiri, M., Hoffman, R. A., & Smith, P. H. (1980). Energetic particle penetrations into the inner magnetosphere. *Journal of Geophysical Research: Space Physics*, 85(A2), 653–663. Retrieved from <https://agupubs.onlinelibrary.wiley.com/doi/abs/10.1029/JA085iA02p00653> doi: <https://doi.org/10.1029/JA085iA02p00653>
- Emery, B. A., Coumans, V., Evans, D. S., Germany, G. A., Greer, M. S., Holeman, E., ... Xu, W. (2008). Seasonal, Kp, solar wind, and solar flux variations in long-term single-pass satellite estimates of electron and ion auroral hemispheric power. *Journal of Geophysical Research: Space Physics*, 113(A6).
- Emery, B. A., Evans, D. S., Greer, M. S., Holeman, E., Kadinsky-Cade, K., Rich, F. J., & Xu, W. (2006). The low energy auroral electron and ion hemispheric power after NOAA and DMSP intersatellite adjustments. *Tech. Note NCAR/TN-470+ STR*.
- Ergun, R. E., Carlson, C. W., McFadden, J. P., Mozer, F. S., Delory, G. T., Peria, W., ... Kistler, L. (1998). FAST satellite observations of electric field structures in the auroral zone. *Geophysical Research Letters*, 25(12), 2025–2028. Retrieved from <https://agupubs.onlinelibrary.wiley.com/doi/abs/10.1029/98GL00635> doi: <https://doi.org/10.1029/98GL00635>
- Evans, D. S. (1974). Precipitating electron fluxes formed by a magnetic field aligned potential difference. *Journal of Geophysical Research (1896-1977)*, 79(19), 2853–2858. Retrieved from <https://agupubs.onlinelibrary.wiley.com/doi/abs/10.1029/JA079i019p02853> doi: <https://doi.org/10.1029/JA079i019p02853>
- Evans, D. S., & Moore, T. E. (1979). Precipitating electrons associated with the

- diffuse aurora: Evidence for electrons of atmospheric origin in the plasma sheet. *Journal of Geophysical Research: Space Physics*, 84(A11), 6451–6457.
- Fedder, J. A., Slinker, S. P., Lyon, J. G., & Elphinstone, R. D. (1995, oct). Global numerical simulation of the growth phase and the expansion onset for a substorm observed by Viking. *Journal of Geophysical Research*, 100(A10), 19083. Retrieved from <http://doi.wiley.com/10.1029/95JA01524> doi: 10.1029/95JA01524
- Fridman, M., & Lemaire, J. (1980). Relationship between auroral electrons fluxes and field aligned electric potential difference. *Journal of Geophysical Research: Space Physics*, 85(A2), 664–670. Retrieved from <https://agupubs.onlinelibrary.wiley.com/doi/abs/10.1029/JA085iA02p00664> doi: 10.1029/JA085iA02p00664
- Galand, M., Fuller-Rowell, T. J., & Codrescu, M. V. (2001, jan). Response of the upper atmosphere to auroral protons. *Journal of Geophysical Research: Space Physics*, 106(A1), 127–139. Retrieved from <http://doi.wiley.com/10.1029/2000JA002009> doi: 10.1029/2000JA002009
- Galand, M., & Richmond, A. D. (2001). Ionospheric electrical conductances produced by auroral proton precipitation. *Journal of Geophysical Research: Space Physics*, 106(A1), 117–125. Retrieved from <https://agupubs.onlinelibrary.wiley.com/doi/abs/10.1029/1999JA002001> doi: <https://doi.org/10.1029/1999JA002001>
- Ganushkina, N. Y., Pulkkinen, T. I., Sergeev, V. A., Kubyshkina, M. V., Baker, D. N., Turner, N. E., ... Fritz, T. A. (2000). Entry of plasma sheet particles into the inner magnetosphere as observed by Polar/CAMMICE. *Journal of Geophysical Research: Space Physics*, 105(A11), 25205–25219. Retrieved from <https://agupubs.onlinelibrary.wiley.com/doi/abs/10.1029/2000JA900062> doi: <https://doi.org/10.1029/2000JA900062>
- Gao, Y. (2012). Comparing the cross polar cap potentials measured by superdarn and amie during saturation intervals. *Journal of Geophysical Research: Space Physics*, 117(A8). Retrieved from <https://agupubs.onlinelibrary.wiley.com/doi/abs/10.1029/2012JA017690> doi: 10.1029/2012JA017690
- Gilson, M. L., Raeder, J., Donovan, E., Ge, Y. S., & Kepko, L. (2012). Global simulation of proton precipitation due to field line curvature during substorms. *Journal of Geophysical Research: Space Physics*, 117(A5). Retrieved from <https://doi.org/10.1029/2012JA017562> doi: <https://doi.org/10.1029/2012JA017562>
- Gkioulidou, M., Ohtani, S., Mitchell, D. G., Ukhorskiy, A. Y., Reeves, G. D., Turner, D. L., ... Lanzerotti, L. J. (2015). Spatial structure and temporal evolution of energetic particle injections in the inner magnetosphere during the 14 July 2013 substorm event. *Journal of Geophysical Research: Space Physics*, 120(3), 1924–1938. Retrieved from <https://agupubs.onlinelibrary.wiley.com/doi/abs/10.1002/2014JA020872> doi: <https://doi.org/10.1002/2014JA020872>
- Glocer, A., Tóth, G., Gombosi, T., & Welling, D. (2009). Modeling ionospheric outflows and their impact on the magnetosphere, initial results. *Journal of Geophysical Research: Space Physics*, 114(A5). Retrieved from <https://agupubs.onlinelibrary.wiley.com/doi/abs/10.1029/2009JA014053> doi: <https://doi.org/10.1029/2009JA014053>
- Glocer, A., Welling, D., Chappell, C. R., Toth, G., Fok, M.-C., Komar, C., ... Mouikis, C. (2020). A Case Study on the Origin of Near-Earth Plasma. *Journal of Geophysical Research: Space Physics*, 125(11), e2020JA028205. Retrieved from <https://agupubs.onlinelibrary.wiley.com/doi/abs/10.1029/2020JA028205> doi: <https://doi.org/10.1029/2020JA028205>
- Gombosi, T. I. (1994). *Gaskinetic Theory*. Cambridge University Press. doi: 10.1017/CBO9780511524943
- Gombosi, T. I., De Zeeuw, D. L., Powell, K. G., Ridley, A. J., Sokolov, I. V., Stout, Q. F., & Tóth, G. (2003). Adaptive mesh refinement for global magnetohy-



- drodynamic simulation. In J. Büchner, M. Scholer, & C. T. Dum (Eds.), *Space plasma simulation* (pp. 247–274). Berlin, Heidelberg: Springer Berlin Heidelberg. Retrieved from [https://doi.org/10.1007/3-540-36530-3\\_12](https://doi.org/10.1007/3-540-36530-3_12) doi: 10.1007/3-540-36530-3\_12
- Goodman, M. L. (1995, aug). A three-dimensional, iterative mapping procedure for the implementation of an ionosphere-magnetosphere anisotropic Ohm's law boundary condition in global magnetohydrodynamic simulations. *Annales Geophysicae*, 13(8), 843–853. Retrieved from <https://doi.org/10.1007/s00585-995-0843-z> doi: 10.1007/s00585-995-0843-z
- Haiducek, J. D., Welling, D. T., Ganushkina, N. Y., Morley, S. K., & Ozturk, D. S. (2017). SWMF Global Magnetosphere Simulations of January 2005: Geomagnetic Indices and Cross-Polar Cap Potential. *Space Weather*, 15(12), 1567–1587. Retrieved from <https://agupubs.onlinelibrary.wiley.com/doi/abs/10.1002/2017SW001695> doi: 10.1002/2017SW001695
- Hardy, D. A., Gussenhoven, M. S., & Brautigam, D. (1989). A statistical model of auroral ion precipitation. *Journal of Geophysical Research: Space Physics*, 94(A1), 370–392. Retrieved from <https://doi.org/10.1029/JA094iA01p00370> doi: <https://doi.org/10.1029/JA094iA01p00370>
- Hardy, D. A., Gussenhoven, M. S., & Holeman, E. (1985). A statistical model of auroral electron precipitation. *Journal of Geophysical Research: Space Physics*, 90(A5), 4229–4248. Retrieved from <https://doi.org/10.1029/JA090iA05p04229> doi: <https://doi.org/10.1029/JA090iA05p04229>
- Hartinger, M. D., Xu, Z., Clauer, C. R., Yu, Y., Weimer, D. R., Kim, H., ... Willer, A. N. (2017). Associating ground magnetometer observations with current or voltage generators. *Journal of Geophysical Research: Space Physics*, 122(7), 7130–7141. Retrieved from <https://agupubs.onlinelibrary.wiley.com/doi/abs/10.1002/2017JA024140> doi: 10.1002/2017JA024140
- Hatch, S. M., LaBelle, J., & Chaston, C. C. (2019). Inferring Source Properties of Monoenergetic Electron Precipitation From Kappa and Maxwellian Moment-Voltage Relationships. *Journal of Geophysical Research: Space Physics*, 124(3), 1548–1567. Retrieved from <https://agupubs.onlinelibrary.wiley.com/doi/abs/10.1029/2018JA026158> doi: <https://doi.org/10.1029/2018JA026158>
- Iijima, T., & Potemra, T. A. (1976). The amplitude distribution of field-aligned currents at northern high latitudes observed by Triad. *Journal of Geophysical Research-Space Physics*, 81(13), 2165–2174. Retrieved from <http://onlinelibrary.wiley.com/doi/10.1029/JA081i013p02165/abstract%5Cnpapers3://publication/doi/10.1029/JA081i013p02165> doi: 10.1029/JA081i013p02165
- Janhunen, P., Olsson, A., Tsyganenko, N. A., Russell, C. T., Laakso, H., & Blomberg, L. G. (2005). Statistics of a parallel Poynting vector in the auroral zone as a function of altitude using Polar EFI and MFE data and Astrid-2 EMMA data. *Annales Geophysicae*, 23(5), 1797–1806. Retrieved from <https://angeo.copernicus.org/articles/23/1797/2005/> doi: 10.5194/angeo-23-1797-2005
- Jordanova, V. K., Welling, D. T., Zaharia, S. G., Chen, L., & Thorne, R. M. (2012). Modeling ring current ion and electron dynamics and plasma instabilities during a high-speed stream driven storm. *Journal of Geophysical Research: Space Physics*, 117(A9). Retrieved from <https://agupubs.onlinelibrary.wiley.com/doi/abs/10.1029/2011JA017433> doi: <https://doi.org/10.1029/2011JA017433>
- Kaeppler, S. R., Hampton, D. L., Nicolls, M. J., Strømme, A., Solomon, S. C., Hecht, J. H., & Conde, M. G. (2015). An investigation comparing ground-based techniques that quantify auroral electron flux and conductance. *Journal of Geophysical Research: Space Physics*, 120(10), 9038–9056. Retrieved from <https://agupubs.onlinelibrary.wiley.com/doi/abs/10.1002/2015JA021396> doi: 10.1002/2015JA021396

- 1181 Kang, S.-B., Gloer, A., Komar, C., Fok, M.-C., & Shim, J. (2019). Wave-induced  
1182 particle precipitation into the ionosphere from the inner magnetosphere. In *2019*  
1183 *ursi asia-pacific radio science conference (ap-rasc)* (p. 1). doi: 10.23919/URSIAP  
1184 -RASC.2019.8738674
- 1185 Keesee, A. M., Chen, M. W., Scime, E. E., & Lui, A. T. Y. (2014). Regions of  
1186 ion energization observed during the Galaxy-15 substorm with TWINS. *Journal*  
1187 *of Geophysical Research: Space Physics*, 119(10), 8274–8287. Retrieved from  
1188 <https://agupubs.onlinelibrary.wiley.com/doi/abs/10.1002/2014JA020466>  
1189 doi: <https://doi.org/10.1002/2014JA020466>
- 1190 Khachikjan, G. Y., Koustov, A. V., & Sofko, G. J. (2008). Dependence of super-  
1191 darn cross polar cap potential upon the solar wind electric field and magnetopause  
1192 subsolar distance. *Journal of Geophysical Research: Space Physics*, 113(A9). Re-  
1193 trieved from [https://agupubs.onlinelibrary.wiley.com/doi/abs/10.1029/](https://agupubs.onlinelibrary.wiley.com/doi/abs/10.1029/2008JA013107)  
1194 [2008JA013107](https://doi.org/10.1029/2008JA013107) doi: 10.1029/2008JA013107
- 1195 Khazanov, G. V., Liemohn, M. W., Krivorutsky, E. N., & Moore, T. E. (1998, apr).  
1196 Generalized kinetic description of a plasma in an arbitrary field-aligned poten-  
1197 tial energy structure. *Journal of Geophysical Research: Space Physics*, 103(A4),  
1198 6871–6889. Retrieved from <http://doi.wiley.com/10.1029/97JA03436> doi:  
1199 10.1029/97JA03436
- 1200 Kivelson, M., & Russell, C. (1995). *Introduction to Space Physics*.
- 1201 Knight, S. (1973). Parallel electric fields. *Planetary and Space Science*. doi: 10.1016/  
1202 0032-0633(73)90093-7
- 1203 Korth, H., Zhang, Y., Anderson, B. J., Sotirelis, T., & Waters, C. L. (2014). Sta-  
1204 tistical relationship between large-scale upward field-aligned currents and electron  
1205 precipitation. *Journal of Geophysical Research: Space Physics*, 119(8), 6715–6731.  
1206 Retrieved from [https://agupubs.onlinelibrary.wiley.com/doi/abs/10.1002/](https://agupubs.onlinelibrary.wiley.com/doi/abs/10.1002/2014JA019961)  
1207 [2014JA019961](https://doi.org/10.1002/2014JA019961) doi: 10.1002/2014JA019961
- 1208 Liemohn, M. W. (2020). The case for improving the Robinson formulas. *Journal of*  
1209 *Geophysical Research: Space Physics*, 125(10), e2020JA028332.
- 1210 Liemohn, M. W., Katus, R. M., & Ilie, R. (2015). Statistical analysis of storm-time  
1211 near-Earth current systems. In *Annales geophysicae* (Vol. 33, pp. 965–982).
- 1212 Liemohn, M. W., & Khazanov, G. V. (1998). Collisionless plasma modeling in an ar-  
1213 bitrary potential energy distribution. *Physics of plasmas*, 5(3), 580–589.
- 1214 Liemohn, M. W., Ridley, A. J., Brandt, P. C., Gallagher, D. L., Kozyra, J. U.,  
1215 Ober, D. M., ... DeMajistre, R. (2005, dec). Parametric analysis of nightside  
1216 conductance effects on inner magnetospheric dynamics for the 17 April 2002  
1217 storm. *Journal of Geophysical Research*, 110(A12), A12S22. Retrieved from  
1218 <http://doi.wiley.com/10.1029/2005JA011109> doi: 10.1029/2005JA011109
- 1219 Liemohn, M. W., Shane, A. D., Azari, A. R., Petersen, A. K., Swiger, B. M., &  
1220 Mukhopadhyay, A. (2021). Rmse is not enough: Guidelines to robust data-model  
1221 comparisons for magnetospheric physics. *Journal of Atmospheric and Solar-*  
1222 *Terrestrial Physics*, 218, 105624.
- 1223 Liemohn, M. W., Welling, D. T., Simpson, J. J., Ilie, R., Anderson, B. J., Zou,  
1224 S., ... Xu, H. (2018). CHARGED: Understanding the physics of extreme ge-  
1225 omagnetically induced currents. In *Agu fall meeting abstracts* (Vol. 2018, pp.  
1226 NH31C—0993).
- 1227 Lin, D., Sorathia, K., Wang, W., Merkin, V., Bao, S., Pham, K., ... al., E.  
1228 (2021). The role of diffuse electron precipitation in the formation of subau-  
1229 roral polarization streams. *Earth and Space Science Open Archive*, 20. Re-  
1230 trieved from <https://doi.org/10.1002/essoar.10508315.1> doi: 10.1002/  
1231 [essoar.10508315.1](https://doi.org/10.1002/essoar.10508315.1)
- 1232 Lin, D., Wang, W., Scales, W. A., Pham, K., Liu, J., Zhang, B., ... Maimaiti,  
1233 M. (2019). SAPS in the 17 March 2013 Storm Event: Initial Results From  
1234 the Coupled Magnetosphere-Ionosphere-Thermosphere Model. *Journal of*

- Geophysical Research: Space Physics*, 124(7), 6212–6225. Retrieved from <https://agupubs.onlinelibrary.wiley.com/doi/abs/10.1029/2019JA026698>  
doi: <https://doi.org/10.1029/2019JA026698>
- Loto'aniu, T. M., Singer, H. J., Rodriguez, J. V., Green, J., Denig, W., Biesecker, D., & Angelopoulos, V. (2015). Space weather conditions during the Galaxy 15 spacecraft anomaly. *Space Weather*, 13(8), 484–502. Retrieved from <https://agupubs.onlinelibrary.wiley.com/doi/abs/10.1002/2015SW001239>  
doi: <https://doi.org/10.1002/2015SW001239>
- Lu, G., Baker, D. N., McPherron, R. L., Farrugia, C. J., Lummerzheim, D., Ruohoniemi, J. M., ... Hayashi, K. (1998). Global energy deposition during the January 1997 magnetic cloud event. *Journal of Geophysical Research: Space Physics*, 103(A6), 11685–11694. Retrieved from <https://agupubs.onlinelibrary.wiley.com/doi/abs/10.1029/98JA00897> doi: <https://doi.org/10.1029/98JA00897>
- Lyons, L. R., Evans, D. S., & Lundin, R. (1979). An observed relation between magnetic field aligned electric fields and downward electron energy fluxes in the vicinity of auroral forms. *Journal of Geophysical Research: Space Physics*, 84(A2), 457–461. Retrieved from <https://agupubs.onlinelibrary.wiley.com/doi/abs/10.1029/JA084iA02p00457> doi: <https://doi.org/10.1029/JA084iA02p00457>
- Merkine, V. G., Papadopoulos, K., Milikh, G., Sharma, A. S., Shao, X., Lyon, J., & Goodrich, C. (2003, dec). Effects of the solar wind electric field and ionospheric conductance on the cross polar cap potential: Results of global MHD modeling. *Geophysical Research Letters*, 30(23), n/a–n/a. Retrieved from <http://doi.wiley.com/10.1029/2003GL017903> doi: 10.1029/2003GL017903
- Moen, J., & Brekke, A. (1993, may). The solar flux influence on quiet time conductances in the auroral ionosphere. *Geophysical Research Letters*, 20(10), 971–974. Retrieved from <http://doi.wiley.com/10.1029/92GL02109> doi: 10.1029/92GL02109
- Morley, S. K., Brito, T. V., & Welling, D. T. (2018). Measures of Model Performance Based On the Log Accuracy Ratio. *Space Weather*, 16(1), 69–88. Retrieved from <https://agupubs.onlinelibrary.wiley.com/doi/abs/10.1002/2017SW001669> doi: 10.1002/2017SW001669
- Möstl, C., Temmer, M., Rollett, T., Farrugia, C. J., Liu, Y., Veronig, A. M., ... Biernat, H. K. (2010). STEREO and Wind observations of a fast ICME flank triggering a prolonged geomagnetic storm on 5–7 April 2010. *Geophysical Research Letters*, 37(24). Retrieved from <https://agupubs.onlinelibrary.wiley.com/doi/abs/10.1029/2010GL045175> doi: <https://doi.org/10.1029/2010GL045175>
- Mukhopadhyay, A., Burleigh, M. B., Welling, D. T., Vandegriff, E., Liemohn, M. W., Ridley, A. J., ... Anderson, B. J. (2021). Challenges in Space Weather Prediction : Discerning the Impact of Ionospheric Conductance in Global Simulations. In *101st american meteorological society annual meeting* (p. Paper ID 11.9). American Meteorological Society.
- Mukhopadhyay, A., Jia, X., Welling, D. T., & Liemohn, M. W. (2021). Global Magnetohydrodynamic Simulations: Performance Quantification of Magnetopause Distances and Convection Potential Predictions. *Frontiers in Astronomy and Space Sciences*, 8, 45. Retrieved from <https://www.frontiersin.org/article/10.3389/fspas.2021.637197> doi: 10.3389/fspas.2021.637197
- Mukhopadhyay, A., Welling, D., Burleigh, M., Ridley, A., Liemohn, M., Anderson, B., & Gjerloev, J. (2019, dec). Conductance in the Aurora: Influence of Magnetospheric Contributors. In *Agu fall meeting abstracts* (Vol. 2019, pp. SA41B–3169). Retrieved from <https://ui.adsabs.harvard.edu/abs/2019AGUFMSA41B3169M/abstract> doi: [doi: 10.1002/essoar.10502150.1](https://doi.org/10.1002/essoar.10502150.1)
- Mukhopadhyay, A., Welling, D., Liemohn, M., Zou, S., & Ridley, A. (2018, dec). Challenges in Space Weather Prediction: Estimation of Auroral Conductance. In *Agu fall meeting abstracts* (Vol. 2018, pp. SA33B–3462). Retrieved from

- <https://ui.adsabs.harvard.edu/abs/2018AGUFMSA33B3462M/abstract>
- Mukhopadhyay, A., Welling, D. T., Liemohn, M. W., Ridley, A. J., Chakraborty, S., & Anderson, B. J. (2020). Conductance Model for Extreme Events : Impact of Auroral Conductance on Space Weather Forecasts. *Space Weather*(19), e2020SW002551. doi: 10.1029/2020SW002551
- Newell, P. T., Liou, K., Zhang, Y., Sotirelis, T., Paxton, L. J., & Mitchell, E. J. (2014). OVATION Prime-2013: Extension of auroral precipitation model to higher disturbance levels. *Space Weather*, 12(6), 368–379. Retrieved from <https://doi.org/10.1002/2014SW001056> doi: <https://doi.org/10.1002/2014SW001056>
- Newell, P. T., Sotirelis, T., Liou, K., Meng, C.-I., & Rich, F. J. (2007). A nearly universal solar wind-magnetosphere coupling function inferred from 10 magnetospheric state variables. *Journal of Geophysical Research: Space Physics*, 112(A1). Retrieved from <https://doi.org/10.1029/2006JA012015> doi: <https://doi.org/10.1029/2006JA012015>
- Newell, P. T., Sotirelis, T., & Wing, S. (2009, sep). Diffuse, monoenergetic, and broadband aurora: The global precipitation budget. *Journal of Geophysical Research: Space Physics*, 114(A9). Retrieved from <http://doi.wiley.com/10.1029/2009JA014326> doi: 10.1029/2009JA014326
- Nishimura, Y., Lessard, M. R., Katoh, Y., Miyoshi, Y., Grono, E., Partamies, N., ... Kurita, S. (2020). Diffuse and Pulsating Aurora. *Space Science Reviews*, 216(1), 4. Retrieved from <https://doi.org/10.1007/s11214-019-0629-3> doi: 10.1007/s11214-019-0629-3
- Nishimura, Y., Lyons, L. R., Gabrielse, C., Sivadas, N., Donovan, E. F., Varney, R. H., ... Zhang, S. R. (2020). Extreme Magnetosphere-Ionosphere-Thermosphere Responses to the 5 April 2010 Supersubstorm. *Journal of Geophysical Research: Space Physics*, 125(4), e2019JA027654. Retrieved from <https://agupubs.onlinelibrary.wiley.com/doi/abs/10.1029/2019JA027654> doi: <https://doi.org/10.1029/2019JA027654>
- Ohtani, S., Wing, S., Merkin, V. G., & Higuchi, T. (2014, jan). Solar cycle dependence of nightside field-aligned currents: Effects of dayside ionospheric conductivity on the solar wind-magnetosphere-ionosphere coupling. *Journal of Geophysical Research: Space Physics*, 119(1), 322–334. Retrieved from <http://doi.wiley.com/10.1002/2013JA019410> doi: 10.1002/2013JA019410
- Østgaard, N., Germany, G., Stadsnes, J., & Vondrak, R. R. (2002). Energy analysis of substorms based on remote sensing techniques, solar wind measurements, and geomagnetic indices. *Journal of Geophysical Research: Space Physics*, 107(A9), SMP 9–1–SMP 9–14. Retrieved from <https://agupubs.onlinelibrary.wiley.com/doi/abs/10.1029/2001JA002002> doi: <https://doi.org/10.1029/2001JA002002>
- Öztürk, D. S., Garcia-Sage, K., & Connor, H. K. (2020). All Hands on Deck for Ionospheric Modeling. *Eos*, 101. Retrieved from <https://doi.org/10.1029/2020EO144365> doi: 10.1029/2020EO144365
- Ozturk, D. S., Zou, S., Ridley, A. J., & Slavin, J. A. (2018). Modeling Study of the Geospace System Response to the Solar Wind Dynamic Pressure Enhancement on 17 March 2015. *Journal of Geophysical Research: Space Physics*, 123(4), 2974–2989. Retrieved from <https://agupubs.onlinelibrary.wiley.com/doi/abs/10.1002/2017JA025099> doi: <https://doi.org/10.1002/2017JA025099>
- Ozturk, D. S., Zou, S., & Slavin, J. A. (2017). IMF By effects on ground magnetometer response to increased solar wind dynamic pressure derived from global MHD simulations. *Journal of Geophysical Research: Space Physics*. doi: 10.1002/2017JA023903
- Paschmann, G., Baumjohann, W., Sckopke, N., Phan, T. D., & Lühr, H. (1993). Structure of the dayside magnetopause for low magnetic shear. *Journal of Geophysical Research: Space Physics*, 98(A8), 13409–13422. Retrieved from



- 1343 <https://agupubs.onlinelibrary.wiley.com/doi/abs/10.1029/93JA00646>  
 1344 doi: <https://doi.org/10.1029/93JA00646>
- 1345 Perlongo, N. J., Ridley, A. J., Liemohn, M. W., & Katus, R. M. (2017, apr). The  
 1346 effect of ring current electron scattering rates on magnetosphere-ionosphere  
 1347 coupling. *Journal of Geophysical Research: Space Physics*, 122(4), 4168–  
 1348 4189. Retrieved from <http://doi.wiley.com/10.1002/2016JA023679> doi:  
 1349 10.1002/2016JA023679
- 1350 Phan, T. D., Paschmann, G., Baumjohann, W., Sckopke, N., & Lühr, H. (1994).  
 1351 The magnetosheath region adjacent to the dayside magnetopause: AMPTE/IRM  
 1352 observations. *Journal of Geophysical Research: Space Physics*, 99(A1), 121–141.  
 1353 Retrieved from [https://agupubs.onlinelibrary.wiley.com/doi/abs/10.1029/](https://agupubs.onlinelibrary.wiley.com/doi/abs/10.1029/93JA02444)  
 1354 [93JA02444](https://doi.org/10.1029/93JA02444) doi: <https://doi.org/10.1029/93JA02444>
- 1355 Powell, K. G., Roe, P. L., Linde, T. J., Gombosi, T. I., & Zeeuw, D. L. D.  
 1356 (1999). A Solution-Adaptive Upwind Scheme for Ideal Magnetohydrodynam-  
 1357 ics. *Journal of Computational Physics*, 154(2), 284–309. Retrieved from  
 1358 <http://www.sciencedirect.com/science/article/pii/S002199919996299X>  
 1359 doi: <https://doi.org/10.1006/jcph.1999.6299>
- 1360 Pulkkinen, A., Kuznetsova, M., Ridley, A., Raeder, J., Vapirev, A., Weimer, D.,  
 1361 ... Chulaki, A. (2011, feb). Geospace Environment Modeling 2008-2009  
 1362 Challenge: Ground magnetic field perturbations. *Space Weather*, 9(2), n/a–  
 1363 n/a. Retrieved from <http://doi.wiley.com/10.1029/2010SW000600> doi:  
 1364 10.1029/2010SW000600
- 1365 Pulkkinen, A., Rastätter, L., Kuznetsova, M., Singer, H., Balch, C., Weimer, D., ...  
 1366 Weigel, R. (2013, jun). Community-wide validation of geospace model ground  
 1367 magnetic field perturbation predictions to support model transition to operations.  
 1368 *Space Weather*, 11(6), 369–385. Retrieved from [http://doi.wiley.com/10.1002/](http://doi.wiley.com/10.1002/swe.20056)  
 1369 [swe.20056](http://doi.wiley.com/10.1002/swe.20056) doi: 10.1002/swe.20056
- 1370 Raeder, J., McPherron, R. L., Frank, L. A., Kokubun, S., Lu, G., Mukai, T., ...  
 1371 Slavin, J. A. (2001). Global simulation of the Geospace Environment Model-  
 1372 ing substorm challenge event. *Journal of Geophysical Research-Space Physics*,  
 1373 106(A1), 381–395. doi: 10.1029/2000ja000605
- 1374 Rastätter, L., Shim, J. S., Kuznetsova, M. M., Kilcommons, L. M., Knipp, D. J.,  
 1375 Codrescu, M., ... Welling, D. (2016). GEM-CEDAR challenge: Poynting flux at  
 1376 DMSP and modeled Joule heat. *Space Weather*, 14(2), 113–135. Retrieved from  
 1377 <https://agupubs.onlinelibrary.wiley.com/doi/abs/10.1002/2015SW001238>  
 1378 doi: <https://doi.org/10.1002/2015SW001238>
- 1379 Ridley, A. J., De Zeeuw, D. L., Gombosi, T. I., & Powell, K. G. (2001). Using  
 1380 steady state MHD results to predict the global state of the magnetosphere-  
 1381 ionosphere system. *Journal of Geophysical Research*, 106(A12), 30067. Re-  
 1382 trieved from <http://adsabs.harvard.edu/abs/2001JGR...10630067R> doi:  
 1383 10.1029/2000JA002233
- 1384 Ridley, A. J., Gombosi, T. I., & De Zeeuw, D. L. (2004). Ionospheric control of the  
 1385 magnetosphere: conductance. *Annales Geophysicae*, 22(2), 567–584. Retrieved  
 1386 from <https://hal.archives-ouvertes.fr/hal-00317238/> doi: 10.5194/angeo-  
 1387 -22-567-2004
- 1388 Ridley, A. J., Gombosi, T. I., Sokolov, I. V., Tóth, G., & Welling, D. T. (2010, aug).  
 1389 Numerical considerations in simulating the global magnetosphere. *Annales Geo-*  
 1390 *physicae*, 28(8), 1589–1614. Retrieved from [http://www.ann-geophys.net/28/](http://www.ann-geophys.net/28/1589/2010/)  
 1391 [1589/2010/](http://www.ann-geophys.net/28/1589/2010/) doi: 10.5194/angeo-28-1589-2010
- 1392 Ridley, A. J., & Kihn, E. A. (2004). Polar cap index comparisons with AMIE cross  
 1393 polar cap potential, electric field, and polar cap area. *Geophysical Research Let-*  
 1394 *ters*, 31(7). Retrieved from [https://agupubs.onlinelibrary.wiley.com/doi/](https://agupubs.onlinelibrary.wiley.com/doi/abs/10.1029/2003GL019113)  
 1395 [abs/10.1029/2003GL019113](https://doi.org/10.1029/2003GL019113) doi: 10.1029/2003GL019113
- 1396 Robinson, R. M., Vondrak, R. R., Miller, K., Dabbs, T., & Hardy, D. (1987,



- mar). On calculating ionospheric conductances from the flux and energy of precipitating electrons. *Journal of Geophysical Research*, 92(A3), 2565. Retrieved from <http://doi.wiley.com/10.1029/JA092iA03p02565> doi: 10.1029/JA092iA03p02565
- Schunk, R., & Nagy, A. (2009). *Ionospheres: Physics, Plasma Physics, and Chemistry* (2nd ed.). Cambridge University Press. doi: 10.1017/CBO9780511635342
- Sergeev, V. A., Sazhina, E. M., Tsyganenko, N. A., & Lundblad JÅand Søråas, F. (1983). Pitch-angle scattering of energetic protons in the magnetotail current sheet as the dominant source of their isotropic precipitation into the nightside ionosphere. *Planetary and Space Science*, 31(10), 1147–1155.
- Sergeev, V. A., & Tsyganenko, N. A. (1982). Energetic particle losses and trapping boundaries as deduced from calculations with a realistic magnetic field model. *Planetary and Space Science*, 30(10), 999–1006.
- Strangeway, R. J. (2010). On the relative importance of waves and electron precipitation in driving ionospheric outflows. In *Agu fall meeting abstracts* (Vol. 2010, pp. SM24B—03).
- Tóth, G., Sokolov, I. V., Gombosi, T. I., Chesney, D. R., Clauer, C. R., De Zeeuw, D. L., ... Kóta, J. (2005, dec). Space Weather Modeling Framework: A new tool for the space science community. *Journal of Geophysical Research*, 110(A12), A12226. Retrieved from <http://doi.wiley.com/10.1029/2005JA011126> doi: 10.1029/2005JA011126
- Tóth, G., van der Holst, B., Sokolov, I. V., De Zeeuw, D. L., Gombosi, T. I., Fang, F., ... Opher, M. (2012). Adaptive numerical algorithms in space weather modeling. *Journal of Computational Physics*, 231(3), 870–903. doi: 10.1016/j.jcp.2011.02.006
- Vandegriff, E., Welling, D. T., Dimmock, A. P., & Morley, S. (2020, December). Small Scale dB/dt Fluctuations: Resolving and Exploring Spikes in Global Models. In *Agu fall meeting abstracts* (Vol. 2020, p. SM003-0004).
- Vandegriff, E., Welling, D. T., Mukhopadhyay, A., Dimmock, A. P., & Morley, S. (2021, December). Forecasting of Localized Geomagnetic Disturbances in Global Models: Physics and Numerics. In *Agu fall meeting abstracts* (Vol. 2021, p. SM41A-0004).
- Wallis, D. D., & Budzinski, E. E. (1981). Empirical models of height integrated conductivities. *Journal of Geophysical Research: Space Physics*, 86(A1), 125–137. Retrieved from <https://agupubs.onlinelibrary.wiley.com/doi/abs/10.1029/JA086iA01p00125> doi: <https://doi.org/10.1029/JA086iA01p00125>
- Wang, C., Gkioulidou, M., Lyons, L. R., & Angelopoulos, V. (2012). Spatial distributions of the ion to electron temperature ratio in the magnetosheath and plasma sheet. *Journal of Geophysical Research: Space Physics* (1978–2012), 117(A8). Retrieved from <https://doi.org/10.1029/2012JA017658> doi: 10.1029/2012JA017658
- Waters, C. L., Anderson, B. J., Green, D. L., Korth, H., Barnes, R. J., & Vanhamäki, H. (2020). Science Data Products for AMPERE. In M. W. Dunlop & H. Lühr (Eds.), *Ionospheric multi-spacecraft analysis tools: Approaches for deriving ionospheric parameters* (pp. 141–165). Cham: Springer International Publishing. Retrieved from [https://doi.org/10.1007/978-3-030-26732-2\\_7](https://doi.org/10.1007/978-3-030-26732-2_7) doi: 10.1007/978-3-030-26732-2\_7
- Welling, D. T. (2019). Magnetohydrodynamic models of b and their use in gic estimates. In *Geomagnetically induced currents from the sun to the power grid* (p. 43-65). American Geophysical Union (AGU). Retrieved from <https://agupubs.onlinelibrary.wiley.com/doi/abs/10.1002/9781119434412.ch3> doi: 10.1002/9781119434412.ch3
- Welling, D. T., Anderson, B. J., Crowley, G., Pulkkinen, A. A., & Rastätter, L. (2017, jan). Exploring predictive performance: A reanalysis of the geospace

- model transition challenge. *Space Weather*, 15(1), 192–203. Retrieved from <http://doi.wiley.com/10.1002/2016SW001505> doi: 10.1002/2016SW001505
- Welling, D. T., Love, J. J., Rigler, E. J., Oliveira, D. M., Komar, C. M., & Morley, S. K. (2021). Numerical Simulations of the Geospace Response to the Arrival of an Idealized Perfect Interplanetary Coronal Mass Ejection. *Space Weather*, 19(2), e2020SW002489. Retrieved from <https://agupubs.onlinelibrary.wiley.com/doi/abs/10.1029/2020SW002489> doi: <https://doi.org/10.1029/2020SW002489>
- Welling, D. T., & Ridley, A. J. (2010). Exploring sources of magnetospheric plasma using multispecies MHD. *Journal of Geophysical Research: Space Physics*, 115(A4). Retrieved from <https://agupubs.onlinelibrary.wiley.com/doi/abs/10.1029/2009JA014596> doi: 10.1029/2009JA014596
- Wiltberger, M., Merkin, V., Zhang, B., Toffoletto, F., Oppenheim, M., Wang, W., ... Stephens, G. K. (2017, may). Effects of electrojet turbulence on a magnetosphere-ionosphere simulation of a geomagnetic storm. *Journal of Geophysical Research: Space Physics*, 122(5), 5008–5027. Retrieved from <http://doi.wiley.com/10.1002/2016JA023700> doi: 10.1002/2016JA023700
- Wiltberger, M., Weigel, R. S., Lotko, W., & Fedder, J. A. (2009). Modeling seasonal variations of auroral particle precipitation in a global-scale magnetosphere-ionosphere simulation. *Journal of Geophysical Research: Space Physics*, 114(1), 1–11. doi: 10.1029/2008JA013108
- Wolf, R. A., Harel, M., Spiro, R. W., Voigt, G.-H., Reiff, P. H., & Chen, C.-K. (1982, aug). Computer simulation of inner magnetospheric dynamics for the magnetic storm of July 29, 1977. *Journal of Geophysical Research*, 87(A8), 5949. Retrieved from <http://doi.wiley.com/10.1029/JA087iA08p05949> doi: 10.1029/JA087iA08p05949
- Wu, C., Ridley, A. J., DeJong, A. D., & Paxton, L. J. (2021). FTA: A Feature Tracking Empirical Model of Auroral Precipitation. *Space Weather*, 19(5), e2020SW002629. Retrieved from <https://agupubs.onlinelibrary.wiley.com/doi/abs/10.1029/2020SW002629> doi: <https://doi.org/10.1029/2020SW002629>
- Yang, J., Toffoletto, F. R., Wolf, R. A., & Sazykin, S. (2011). RCM-E simulation of ion acceleration during an idealized plasma sheet bubble injection. *Journal of Geophysical Research: Space Physics*, 116(A5). Retrieved from <https://agupubs.onlinelibrary.wiley.com/doi/abs/10.1029/2010JA016346> doi: <https://doi.org/10.1029/2010JA016346>
- Yu, Y., Jordanova, V. K., McGranaghan, R. M., & Solomon, S. C. (2018). Self-Consistent Modeling of Electron Precipitation and Responses in the Ionosphere: Application to Low-Altitude Energization During Substorms. *Geophysical Research Letters*, 45(13), 6371–6381. Retrieved from <https://agupubs.onlinelibrary.wiley.com/doi/abs/10.1029/2018GL078828> doi: <https://doi.org/10.1029/2018GL078828>
- Yu, Y., Jordanova, V. K., Ridley, A. J., Albert, J. M., Horne, R. B., & Jeffrey, C. A. (2016). A new ionospheric electron precipitation module coupled with ram-scb within the geospace general circulation model. *Journal of Geophysical Research: Space Physics*, 121(9), 8554–8575. Retrieved from <https://agupubs.onlinelibrary.wiley.com/doi/abs/10.1002/2016JA022585> doi: 10.1002/2016JA022585
- Yu, Y., Ridley, A. J., Welling, D. T., & Tóth, G. (2010, aug). Including gap region field-aligned currents and magnetospheric currents in the MHD calculation of ground-based magnetic field perturbations. *Journal of Geophysical Research: Space Physics*, 115(A8). Retrieved from <http://doi.wiley.com/10.1029/2009JA014869> doi: 10.1029/2009JA014869
- Zhang, B., Lotko, W., Brambles, O., Wiltberger, M., & Lyon, J. (2015). Electron precipitation models in global magnetosphere simulations. *Journal of Geophysical Research: Space Physics*, 120(2), 1035–1056. Retrieved from

- 1505 <https://agupubs.onlinelibrary.wiley.com/doi/abs/10.1002/2014JA020615>  
1506 doi: 10.1002/2014JA020615
- 1507 Zhang, B., Lotko, W., Brambles, O., Xi, S., Wiltberger, M., & Lyon, J. (2014).  
1508 Solar wind control of auroral Alfvénic power generated in the magnetotail. *Jour-*  
1509 *nal of Geophysical Research: Space Physics*, 119(3), 1734–1748. Retrieved from  
1510 <https://agupubs.onlinelibrary.wiley.com/doi/abs/10.1002/2013JA019178>  
1511 doi: <https://doi.org/10.1002/2013JA019178>
- 1512 Zheng, Y., Brandt, P. C., Lui, A. T. Y., & Fok, M.-C. (2008). On ionospheric  
1513 trough conductance and subauroral polarization streams: Simulation results.  
1514 *Journal of Geophysical Research: Space Physics*, 113(A4). Retrieved from  
1515 <https://agupubs.onlinelibrary.wiley.com/doi/abs/10.1029/2007JA012532>  
1516 doi: <https://doi.org/10.1029/2007JA012532>