


Estimating the Contribution of Brick Kiln Industry to PM_{2.5} Emission over Northern India

Ardhi A. Arbain¹ (ardhi@aori.u-tokyo.ac.jp) and Ryoichi Imasu¹

¹Atmosphere and Ocean Research Institute, The University of Tokyo, Kashiwa, Japan

1. BACKGROUND

1-1 Introduction



PM_{2.5} or fine particulate matters are tiny aerosols with diameter of 2.5µm or less, which are typically generated from incomplete combustion of fossil fuel in vehicles, biomass In stoves for cooking and heating, coal in small industrial operations and agricultural waste in post-harvest fields. It's also the PM_{2.5} which plays the role of major component of air pollution which is now choking India.


Brick kiln and PM_{2.5} emission

India is the second largest brick producer in the world. Brick kiln (BK) emits fine particles of coal, dust particles, organic matters and gases such as SO₂, NO_x, H₂S, CO etc.



Estimated annual PM_{2.5} emission: 23,300t for Great Dhaka region alone (Schmidt, 2013)

1-2 Key problems




- Limited inventory for PM_{2.5} emission originated from BK.
- In-situ observations using aerial photo, UAV or drone are difficult due to air-space regulations.
- Lack of remote sensing products for identifying BK locations.

1-3 Study objectives

- Develop a novel technique to estimate regional PM_{2.5} emission from BK over Northern India using numerical model simulation and remote sensing data
- Evaluate the effectiveness of countermeasure in reducing PM_{2.5} emission from brick industry in India


2. METHODOLOGY

2-1 Model design




3. RESULTS

3-1 PM_{2.5} simulation



3-2 MODIS land products analysis



2-2 Data used

- ECMWF ERA-Interim (2017-2018, 6-hourly,~80km spatial res.)
- MOZART/WACCM chemistry boundary dataset (2017-2018)
- MODIS LAI/FPAR dataset (2017-2018, 8-daily, 500m spatial res.)
- MODIS annual land cover products (Jan 2018, 500m spatial res.)
- MODIS AOD (2017-2018, daily, 1km spatial res.)
- BK locations (2016, 3887 points from Google map)
- BK locations based on LANDSAT8 and PALSAR2 data identification using GEE cloud platform

2-3 Model parameterization

Configuration	Scheme	Namelist Value	References
Longwave radiation	RRTMG	ra_lw_physics=4	Iacono et. al (2008)
Shortwave radiation	RRTMG	ra_sw_physics=4	Iacono et. al (2008)
Microphysics	Lin et. al	mp_physics=2	Lin et. al (1983)
Cumulus	Grell3D	cu_physics=5	Grell (1993), Grell and Devenyi (2002)
Planetary Boundary Layer	YSU	Bl_bl_physics=1	Hong et. al (2006)
Land surface	NOAH LSM	sf_surface_physics=2	Niu et. al (2011), Yang et. al (2011)
Surface layer	5-layer Thermal Diffusion	sf_sfclay_physics=1	Dudhia (1996)

4. CONCLUSIONS AND FUTURE WORKS

4-1 Conclusions

- More than 60% of BK plantations are located in the silt-rich areas for making brick kiln with more than 80% plantations are situated over cropland regions in the outskirts of urban areas in India.
- Simulated PM_{2.5} emission shows diurnal variations over the locations of BK plantations

4-2 Future works

- PM_{2.5} simulations using BK location generated by GEE cloud platform
- In-situ observations using digital imaging to estimate the emission of BK plumes

Acknowledgement

- Brick kiln locations using GEE cloud platform were provided by Dr. Prakhar Misra and Prof. Wataru Takeuchi of Institute of Industrial Science, The University of Tokyo

References

Anenberg et al., 2012: *Environ. Health Persp.*, 120(6), 831–839
Anenberg et al., 2011: *Atmos. Chem. Phys.*, 11(14), 7253–7267
Du et al., 2007: *Environ. Sci. Technol.*, 41(3), 928-935
Gao et al., 2015: *Environ. Pollut.*, 199, 56–65
Gorelick et al., 2017: *Remote Sens. Environ.*, 202, 18-27
Grell et al., 2005: *Atmos. Environ.*, 39(37), 6957–6975
Habil et al., 2016: *Data Brief*, 6, 495–502
Mc Farland et al., 2004: *J. Air Waste Assoc.*, 54(3), 296-306
Powers et al., 2017: *Bull. Amer. Meteor. Soc.*, 98(8), 1717–1737
Vadrevu et al., 2011: *Environ. Pollut.*, 159(6), 1560–1569