

# The influence of the stratospheric quasi-biennial oscillation on the tropical easterly jet over the Maritime Continent

Yuanpu Li<sup>1\*</sup>, Sihua Huang<sup>1</sup> and Zhiping Wen<sup>1\*</sup>

<sup>1</sup> Institute of Atmospheric Sciences, Fudan University, Shanghai, China

\*Corresponding author:

Yuanpu Li ([liyuanpu@fudan.edu.cn](mailto:liyuanpu@fudan.edu.cn))

ORCID: <https://orcid.org/0000-0003-3047-5675>

Zhiping Wen ([zpwen@fudan.edu.cn](mailto:zpwen@fudan.edu.cn))

ORCID: <https://orcid.org/0000-0003-0178-9835>

## Key Points:

Main point #1:

TEJ is weakened over the Maritime Continent during easterly phase QBO.

Main point #2:

QBO regulates TEJ by modulating the stability near the tropopause and the convection over the Maritime Continent.

Main point #3:

Different convection heights between the east and west sides of the Maritime Continent lead to different responses to QBO.

34    **Abstract**

35    This study investigated the influence of the stratospheric quasi-biennial oscillation  
36    (QBO) on the tropical easterly jet (TEJ). Easterly (EQBO) and westerly (WQBO)  
37    phases of the QBO are defined based on the 50 hPa zonal wind. The climatological  
38    rising movement on the west side of the Maritime Continent can reach higher altitude  
39    than that on the east side, which makes the convection on the west side more  
40    effectively promoted by the reduced stability near the tropopause during EQBO.  
41    Compared with WQBO, during EQBO, the convection on the west (east) side of the  
42    Maritime Continent is enhanced (weakened), and there is a stronger (weaker)  
43    divergence in the upper troposphere. Corresponding to the change of the divergence  
44    field, the TEJ over the Maritime Continent during EQBO is significantly weakened  
45    than WQBO, and the magnitude of the change can reach 11% of the climatological  
46    TEJ.

47

## **Plain Language Summary**

The tropical easterly jet (TEJ) is an essential component of the Asian monsoon system. Quasi-biennial oscillation (QBO) is a phenomenon characterized by quasi-periodic oscillation in the tropical stratospheric wind field. The opposite east-west wind directions in the lower stratosphere during different phases of QBO are accompanied by opposite temperature and stability anomalies around the tropopause. Due to the different heights of the rising movements on the east and west sides of the Maritime Continent, their responses to the change of upper tropospheric stability caused by QBO are different. Responding to the change of the rising movement, the TEJ over the Maritime Continent during easterly phase of QBO is significantly weakened compared with westerly phase of QBO, and the decrease magnitude can reach 11% of the climatological TEJ. Considering that QBO has a relatively robust periodicity, the relationship between QBO and TEJ has potential merit for long-range prediction of TEJ and weather systems related to the Asian monsoon.

**Keywords:** quasi-biennial oscillation; tropical easterly jet; tropical convection

## 1. Introduction

The tropical easterly jet (TEJ) has a strong easterly wind center in the upper troposphere over the Indian Ocean in boreal summer. The maximum wind speed of the Indian Ocean TEJ can reach above 20 m/s (*Huang et al.*, 2020). TEJ results from the thermal contrast between the Asian continent and the Ocean and is enhanced by the local thermal effect of the Tibetan Plateau (*Koteswaram*, 1958). TEJ is an important component of the Asian monsoon system (*Krishnamurti and Bhalme*, 1976) and has an essential effect on the weather and climate in the region influenced by monsoons, e.g., the high-cloud amount in the Asian monsoon region (*Sathiyamoorthy et al.*, 2004), the Tropical Cyclonic Systems in the Bay of Bengal (*Rao et al.*, 2004), Indian summer monsoon rainfall (*Huang et al.*, 2021), summer rainfall of eastern Africa (*Vashisht et al.*, 2021), et al.. The Maritime Continent is the entrance of the Indian Ocean TEJ. The TEJ intensity over the Maritime Continent and tropical cyclone genesis frequency over the Western North Pacific have a positive relationship (*Zhan et al.*, 2022). The TEJ is located in the upper troposphere and can extend to the bottom of the stratosphere. Previous studies have focused on the influence from the troposphere on TEJ (*Chen and van Loon*, 1987; *Huang et al.*, 2019; *Lu and Ding*, 1989; *Pattanaik and Satyan*, 2000; *Tanaka*, 1982), but it is still unclear whether the stratosphere plays a role in it.

The mean zonal winds of the equatorial stratosphere switch between easterly and westerly winds with periods varying from about 24 to 30 months, which is named the quasi-biennial oscillation (QBO) (*Holton and Hakim*, 2013). The associated

stratospheric wind field has zonal uniformity along the equator and is symmetrical about the equator. QBO is driven by tropical atmospheric disturbances such as Kelvin waves, Rossby-gravity waves, and Inertia-gravity waves with periods much smaller than QBO (*Baldwin et al.*, 2001; *Dunkerton*, 1997; *Plumb*, 1977). The fact that QBO is confined near the equator in the stratosphere rather than in the extratropics is due to the smaller Coriolis torque near the equator (*Lindzen and Holton*, 1968; *Scott and Haynes*, 1998). There are many studies on the effects of QBO on the tropical tropospheric weather systems (more detailed introduction are in the latter text), e.g., Madden-Julian oscillation (MJO) (*Back et al.*, 2020; *Martin et al.*, 2021; *Son et al.*, 2017), the boreal summer monsoon (*Giorgetta et al.*, 1999; *Rai and Dimri*, 2017), and tropical deep convection (*Liess and Geller*, 2012; *Nie and Sobel*, 2015).

QBO is one of the most predictable variations of the tropical atmosphere (Hamilton et al. 2015) and TEJ is an important member of the Asian monsoon system, therefore exploring the relationship between QBO and TEJ may have the potential to improve the longer-range forecasting skills of the summer Asian monsoon system. The influence of the stratosphere on the boreal troposphere in winter is extensively studied, for example, the QBO-MJO relationship is more significant in winter (*Wang et al.*, 2019), while the relationship between the summer stratosphere and members of the Asian monsoon system such as TEJ is worth studying.

## 2. Data and method

The summer mean (June to August) of the zonal wind at 50 hPa (U50) observed at

the Singapore radiosonde station ( $1^{\circ}\text{N}/104^{\circ}\text{E}$ ) are used to define the QBO phases. When the standardized U50 is less than half of its standard deviation, then the summer is defined as EQBO. When the standardized U50 is greater than half of its standard deviation, then the summer is defined as WQBO. EQBO and WQBO years are shown in Table 1. The time evolution of the zonal mean zonal wind at 50 hPa at different latitudes is shown in Figure 1. The Meteorological fields analyzed in this study are based on ERA5 provided by ECMWF (*Hersbach et al.*, 2020). NOAA Climate Data Record of Monthly Outgoing Longwave Radiation (OLR) (CDR OLR; Hai-Tien Lee and NOAA Climate Data Record Program, 2018) is used to analyze convection.

A Monte Carlo test has been applied to test the statistical significance of the difference between EQBO and WQBO. We randomly subsampled M and N elements from all years and obtain the difference between the two subsets. M and N are the number of EQBO and WQBO years, respectively. We repeated this random selection process 10,000 times to obtain a probability density function. The significance of the difference was estimated using the density function. Anomalies of EQBO/WQBO from the climatology were also tested by Monte Carlo method.

### 3. Result

The patterns of tropical upper tropospheric zonal wind anomalies of EQBO and WQBO are similar but with the opposite signs. There are westerly wind anomalies at 150 hPa over the Maritime Continent where the climatological entrance of the

summer TEJ locates during EQBO (Figures 2a and 2c), while there are slightly easterly wind anomalies during WQBO (Figures 2b and 2d). Figure 2 implies the TEJ over the Maritime Continent during EQBO is weaker than that during WQBO. Sectioned from 120°E, the vertical structures of the zonal wind anomalies during EQBO and WQBO show that the zonal wind anomalies above 100 hPa in the stratosphere reflect typical QBO characteristics (*Baldwin et al.*, 2001). In the middle and lower stratosphere, the maximum values of the zonal wind anomalies are located at 10 and 50 hPa at the equator. In the upper troposphere, the maximum values of the zonal wind anomalies are located at 150-200 hPa over the Maritime Continent. The wind anomalies in the stratosphere have strong zonally uniformity along the equator, while the wind anomalies in the troposphere do not have such zonally uniformity. The average value of U50 during EQBO is -18.1 m/s, while that during WQBO is 9.9 m/s, and its absolute value is about half of that of EQBO. Similarly, the absolute value of zonal wind anomalies at upper troposphere over the Maritime Continent during WQBO are about half of that during EQBO. The correlation coefficient between the year-to-year time series of U50 and the zonal wind averaged in the box region at 120° E (box region shown in Figures 2c and 2d) is -0.34, significant at the 95% confident level under t-test, indicating a close relationship between QBO and TEJ over Maritime Continent.

The rainfall plays an important role in regulating TEJ variability (*Kanamitsu et al.*, 1972; *Rao and Srinivasan*, 2016; *Sathiyamoorthy et al.*, 2007), which implies the convection and rising movement in the tropics have an important impact on the

variation of TEJ. Figure 3 shows the differences in the convection between EQBO and WQBO. Summer tropical rising and convection are strongest in the Bay of Bengal and Maritime Continent. Both the observed OLR (Figure 3b) and reanalysis  $\omega$  (Figure 3d) show that convections on the west side of the Maritime Continent during EQBO are stronger than that during WQBO, while convections on the east side of the Maritime Continent during EQBO are weaker than that during WQBO.

During EQBO, the convective enhancement area on the west side of the Maritime Continent is accompanied by the increase of the upper tropospheric divergence, and the convective weakening area on the east side of the Maritime Continent is accompanied by the decrease of the upper tropospheric divergence (Figure 3e). Figure 3f shows the differences in atmospheric circulation along with the equatorial profile between EQBO and WQBO. Compared with WQBO, during EQBO, the rising movement on the west side of the Maritime Continent is stronger, and the rising movement on the east side of the Maritime Continent is weakened, resulting in westerly wind anomalies in the upper troposphere, which is consistent with the divergence change in the upper troposphere. The westerly wind anomalies lead to the weakness of the TEJ over the Maritime Continent.

The effects of QBO on tropical convection have been addressed in recent studies on the relationship between QBO and intraseasonal oscillations, including the Madden Julian Oscillation (MJO) (*Jiang et al.*, 2020; *Nishimoto and Yoden*, 2017; *Son et al.*, 2017; *Wang and Wang*, 2021; *Yoo and Son*, 2016) and Boreal Summer Intraseasonal Oscillation (BSISO) (*Wang et al.*, 2019). Previous studies provide some



mechanisms by which QBO regulates tropical convection (*Martin et al.*, 2021). The equatorial westerly and easterly wind shears in the lower stratosphere appear with warm and cold temperature anomalies, respectively (*Holton and Hakim*, 2013). The wind shear anomalies during EQBO would uplift the tropical tropopause and lead to ascension and a cooler tropopause, while the opposite scenario applies for WQBO (*Collimore et al.*, 2003). The QBO-stratification mechanism (*Collimore et al.*, 2003; *Giorgetta et al.*, 1999; *Gray et al.*, 1992; *Liess and Geller*, 2012; *Nie and Sobel*, 2015) states that EQBO could destabilize the upper troposphere and lower stratosphere, promoting more vigorous deep convection. If the vertical extent of the convection related with the QBO increases, there would be larger convective horizontal extent and more cloud amount (*Gray et al.*, 1992). Figure 4 shows that the tropospheric stability above 150 hPa during EQBO is less than that during WQBO, which are the cases on both sides of the Maritime Continent.. Although Figure 4 analyzed the effect of QBO on the stability in summer, the results here are consistent with the mechanism of QBO affecting stability in winter, and the model simulations also support this result (*Martin et al.*, 2019; *Nie and Sobel*, 2015).

During EQBO, the stability above 150 hPa is reduced consistently on both sides of the Maritime Continent, but the convection and rising movement on the east and west sides of the Maritime Continent change inconsistently. Only about 1% of tropical convective systems can reach above 150 hPa (*Liu and Zipser*, 2005). *Nishimoto and Yoden* (2017) stated that only the deep convection developed very high can be affected by the stability conditions near the tropopause. Inspired by their work, the

difference in the heights of the rising movement between the east and west sides of the Maritime Continent is investigated. Figure 3f shows that the climatological rising movement west of 120°E in the equatorial region can reach above 150 hPa, while the rising movement east of 120°E cannot reach 150 hPa. This implies that even if EQBO causes a decrease in stability around the tropopause, only convections on the west side of the Maritime Continent can effectively take advantage of this condition to furtherly develop.

In addition to the impact of the climatological rising height difference between the two sides of the Maritime Continent, there are differences in stability at 300-150 hPa between the two sides of the Maritime Continent. The stability at 300-150 hPa at 90°E (west of the Maritime Continent) during EQBO is smaller than that of WQBO (Figure 4a), while the stability at 300-150 hPa at 140°E (east of the Maritime Continental) during EQBO is slightly larger than that of WQBO (Figure 4b). The smaller stability contributes to the development of convection, so the rising movement on the west side of the Maritime Continent is enhanced while the rising movement on the east side of the Maritime Continent is weakened during the EQBO compared with the WQBO.

#### **4. Summary**

In this study, the effects of different phases of stratospheric QBO on the summer TEJ were investigated. During EQBO, the rising movement on the west side of the Maritime Continent is enhanced while that on the east side of the Maritime Continent is weakened. The changes in the rising movement lead to the enhancement of the

218 divergence on the west side of the Maritime Continent and the decrease of the  
 219 divergence on the east side of the Maritime Continent in the upper equatorial  
 220 troposphere, which results in the decrease of TEJ over the Maritime Continent. While  
 221 the opposite scenario applies for WQBO. EQBO has a greater effect on TEJ over the  
 222 Maritime Continent than WQBO, because the easterly wind at 50 hPa at the equator  
 223 can reach a maximum of 24 m/s during EQBO summer, while the westerly wind at 50  
 224 hPa can only reach a maximum of 13 m/s during WQBO summer. The contribution of  
 225 QBO to the changes of TEJ is important, since the difference in TEJ over the  
 226 Maritime Continent between EQBO and WQBO can reach 17% of the climatological  
 227 intensity of TEJ (see Table 1, calculated by  $[\text{TEJ}_{\text{EQBO}} - \text{TEJ}_{\text{WQBO}}] / \text{TEJ}_{\text{CLIMATE}}$ ). The  
 228 correlation coefficient between the U50 of QBO and the strength of TEJ over the  
 229 Maritime Continent is -0.34, although it is not very high, it passed the significance  
 230 test, which implies that QBO is a potential contribution factor for seasonally  
 231 forecasting TEJ.

232 About 0.1% of tropical convective systems can penetrate above 100 hPa (*Liu and*  
 233 *Zipser, 2005*), due to the weak vertical movement around the tropopause, although the  
 234 TEJ is located in the upper troposphere and is very close to the QBO wind field in the  
 235 lower stratosphere, it is difficult to directly connect the QBO wind anomaly with the  
 236 TEJ through the momentum exchange of the vertical movement. We propose that  
 237 QBO affects TEJ by changing the upper tropospheric stability and tropical  
 238 convections. The mechanism of the relationship between QBO and convection has  
 239 been proposed and confirmed by many previous studies. The question that this study

needs to explain is why the convections on the east and west sides of the Maritime Continent have opposite changes during different phases of QBO. We found that the climatological rising movement on the west side of the Maritime Continent (west of 120°E) can reach above 150 hPa, but that on the east side of the Maritime Continent cannot. This climatological characteristic makes the convection on the west side more effective in taking advantage of the reduced stability near the tropopause during EQBO. In addition, EQBO reduces the stability at 300-150 hPa on the west side of the Maritime Continent rather than the east side, which is also conducive to the rising movement development on the west side of the Maritime Continent.

## Acknowledgments

We are grateful to the groups and agencies for providing the datasets used in this study. ERA5 data can be downloaded from <https://cds.climate.copernicus.eu/cdsapp#!/dataset/reanalysis-era5-pressure-levels?tab=form>. The QBO series of Singapore data can be obtained from <https://www.geo.fu-berlin.de/en/met/ag/strat/produkte/qbo/>. The monthly OLR data can be downloaded from <https://www.ncei.noaa.gov/products/climate-data-records/outgoing-longwave-radiation-monthly>. This work was supported jointly by the National Natural Science Foundation of China (42030601, 41805029, 41875087), and China Postdoctoral Science Foundation (2018M641913).

## Competing interests

The Authors declare no Competing Financial or Non-Financial Interests.

## References

- Back, S.-Y., J.-Y. Han, and S.-W. Son (2020), Modeling Evidence of QBO-MJO Connection: A Case Study, *Geophysical Research Letters*, *47*(20), e2020GL089480.
- Baldwin, M. P., et al. (2001), The quasi-biennial oscillation, *Reviews of Geophysics*, *39*(2), 179-229.
- Chen, T.-C., and H. van Loon (1987), Interannual Variation of the Tropical Easterly Jet, *Monthly Weather Review*, *115*(8), 1739-1759.
- Collimore, C. C., D. W. Martin, M. H. Hitchman, A. Huesmann, and D. E. Waliser (2003), On The Relationship between the QBO and Tropical Deep Convection, *Journal of Climate*, *16*(15), 2552-2568.
- Dunkerton, T. J. (1997), The role of gravity waves in the quasi-biennial oscillation, *Journal of Geophysical Research: Atmospheres*, *102*(D22), 26053-26076.
- Giorgetta, M. A., L. Bengtsson, and K. Arpe (1999), An investigation of QBO signals in the east Asian and Indian monsoon in GCM experiments, *Climate Dynamics*, *15*(6), 435-450.
- Gray, W. M., J. D. Sheaffer, and J. A. Knaff (1992), Influence of the Stratospheric QBO on ENSO Variability, *Journal of the Meteorological Society of Japan. Ser. II*, *70*(5), 975-995.
- Hersbach, H., et al. (2020), The ERA5 global reanalysis, *Quarterly Journal of the Royal Meteorological Society*, *146*(730), 1999-2049.
- Hai-Tien Lee and NOAA CDR Program (2018): NOAA Climate Data Record (CDR) of Monthly

284 Outgoing Longwave Radiation (OLR), Version 2.7. NOAA National Centers for  
 285 Environmental Information. <https://doi.org/10.7289/V5W37TKD>

286 Holton, J. R., and G. J. Hakim (2013), Chapter 12 - Middle Atmosphere Dynamics, in *An*  
 287 *Introduction to Dynamic Meteorology (Fifth Edition)*, edited by J. R. Holton and G. J.  
 288 Hakim, pp. 413-452, Academic Press, Boston.

289 Huang, S., B. Wang, and Z. Wen (2020), Dramatic Weakening of the Tropical Easterly Jet  
 290 Projected by CMIP6 Models, *Journal of Climate*, *33*(19), 8439-8455.

291 Huang, S., B. Wang, Z. Wen, and Z. Chen (2021), Enhanced Tropical Eastern Indian Ocean  
 292 Rainfall Breaks down the Tropical Easterly Jet-Indian Rainfall Relationship, *Journal of*  
 293 *Climate*, 1-44.

294 Huang, S., Z. Wen, Z. Chen, X. Li, R. Chen, and Y. Guo (2019), Interdecadal change in the  
 295 relationship between the tropical easterly jet and tropical sea surface temperature  
 296 anomalies in boreal summer, *Climate Dynamics*, *53*(3), 2119-2131.

297 Jiang, X., Á. F. Adames, D. Kim, E. D. Maloney, H. Lin, H. Kim, C. Zhang, C. A. DeMott, and N.  
 298 P. Klingaman (2020), Fifty Years of Research on the Madden-Julian Oscillation: Recent  
 299 Progress, Challenges, and Perspectives, *Journal of Geophysical Research: Atmospheres*,  
 300 *125*(17), e2019JD030911.

301 Kanamitsu, M., T. N. Krishnamurti, and C. Depradine (1972), On Scale Interactions in the  
 302 Tropics During Northern Summer, *Journal of Atmospheric Sciences*, *29*(4), 698-706.

303 Koteswaram, P. (1958), The Easterly Jet Stream in the Tropics, *Tellus*, *10*(1), 43-57.

304 Krishnamurti, T. N., and H. N. Bhalme (1976), Oscillations of a Monsoon System. Part I.  
 305 Observational Aspects, *Journal of Atmospheric Sciences*, *33*(10), 1937-1954.

306 Liess, S., and M. A. Geller (2012), On the relationship between QBO and distribution of  
 307 tropical deep convection, *Journal of Geophysical Research: Atmospheres*, 117(D3).  
 308 Lindzen, R. S., and J. R. Holton (1968), A Theory of the Quasi-Biennial Oscillation, *Journal of*  
 309 *Atmospheric Sciences*, 25(6), 1095-1107.  
 310 Liu, C., and E. J. Zipser (2005), Global distribution of convection penetrating the tropical  
 311 tropopause, *Journal of Geophysical Research: Atmospheres*, 110(D23).  
 312 Lu, J., and Y. Ding (1989), Climatic study on the summer tropical easterly jet at 200 hPa,  
 313 *Advances in Atmospheric Sciences*, 6(2), 215-226.  
 314 Martin, Z., S. Wang, J. Nie, and A. Sobel (2019), The Impact of the QBO on MJO Convection  
 315 in Cloud-Resolving Simulations, *Journal of the Atmospheric Sciences*, 76(3), 669-688.  
 316 Martin, Z., S.-W. Son, A. Butler, H. Hendon, H. Kim, A. Sobel, S. Yoden, and C. Zhang (2021),  
 317 The influence of the quasi-biennial oscillation on the Madden–Julian oscillation, *Nature*  
 318 *Reviews Earth & Environment*, 2(7), 477-489.  
 319 Nie, J., and A. H. Sobel (2015), Responses of Tropical Deep Convection to the QBO:  
 320 Cloud-Resolving Simulations, *Journal of the Atmospheric Sciences*, 72(9), 3625-3638.  
 321 Nishimoto, E., and S. Yoden (2017), Influence of the Stratospheric Quasi-Biennial Oscillation  
 322 on the Madden–Julian Oscillation during Austral Summer, *Journal of the Atmospheric*  
 323 *Sciences*, 74(4), 1105-1125.  
 324 Pattanaik, D. R., and V. Satyan (2000), Fluctuations of Tropical Easterly Jet during contrasting  
 325 monsoons over India: A GCM study, *Meteorology and Atmospheric Physics*, 75(1), 51-60.  
 326 Plumb, R. A. (1977), The Interaction of Two Internal Waves with the Mean Flow: Implications  
 327 for the Theory of the Quasi-Biennial Oscillation, *Journal of Atmospheric Sciences*, 34(12),

328 1847-1858.

329 Rai, P., and A. P. Dimri (2017), Effect of changing tropical easterly jet, low level jet and  
330 quasi-biennial oscillation phases on Indian summer monsoon, *Atmospheric Science*  
331 *Letters*, *18*(2), 52-59.

332 Rao, B. R. S., D. V. B. Rao, and V. B. Rao (2004), Decreasing trend in the strength of Tropical  
333 Easterly Jet during the Asian summer monsoon season and the number of tropical  
334 cyclonic systems over Bay of Bengal, *Geophysical Research Letters*, *31*(14).

335 Rao, S., and J. Srinivasan (2016), The impact of latent heating on the location and strength of  
336 the tropical easterly jet, *Meteorology and Atmospheric Physics*, *128*(2), 247-261.

337 Sathiyamoorthy, V., P. K. Pal, and P. C. Joshi (2004), Influence of the Upper-Tropospheric  
338 Wind Shear upon Cloud Radiative Forcing in the Asian Monsoon Region, *Journal of*  
339 *Climate*, *17*(14), 2725-2735.

340 Sathiyamoorthy, V., P. K. Pal, and P. C. Joshi (2007), Intraseasonal variability of the Tropical  
341 Easterly Jet, *Meteorology and Atmospheric Physics*, *96*(3), 305-316.

342 Scott, R. K., and P. H. Haynes (1998), Internal interannual variability of the extratropical  
343 stratospheric circulation: The low-latitude flywheel, *Quarterly Journal of the Royal*  
344 *Meteorological Society*, *124*(550), 2149-2173.

345 Son, S.-W., Y. Lim, C. Yoo, H. H. Hendon, and J. Kim (2017), Stratospheric Control of the  
346 Madden–Julian Oscillation, *Journal of Climate*, *30*(6), 1909-1922.

347 Tanaka, M. (1982), Interannual Fluctuations of the Tropical Easterly Jet and the Summer  
348 Monsoon in the Asian Region, *Journal of the Meteorological Society of Japan. Ser. II*,  
349 *60*(3), 865-875.



350 Vashisht, A., B. Zaitchik, and A. Gnanadesikan (2021), ENSO Teleconnection to Eastern  
351 African Summer Rainfall in Global Climate Models: Role of the Tropical Easterly Jet,  
352 *Journal of Climate*, 34(1), 293-312.

353 Wang, F., and L. Wang (2021), An exploration of the connection between quasi-biennial  
354 oscillation and Madden-Julian oscillation, *Environmental Research Letters*, 16(11),  
355 114021.

356 Wang, S., M. K. Tippett, A. H. Sobel, Z. K. Martin, and F. Vitart (2019), Impact of the QBO on  
357 Prediction and Predictability of the MJO Convection, *Journal of Geophysical Research:*  
358 *Atmospheres*, 124(22), 11766-11782.


359 Yoo, C., and S.-W. Son (2016), Modulation of the boreal wintertime Madden-Julian oscillation  
360 by the stratospheric quasi-biennial oscillation, *Geophysical Research Letters*, 43(3),  
361 1392-1398.

362 Zhan, R., Y. Wang, and Y. Ding (2022), Impact of the Western Pacific Tropical Easterly Jet on  
363 Tropical Cyclone Genesis Frequency over the Western North Pacific, *Advances in*  
364 *Atmospheric Sciences*, 39(2), 235-248.

365


366

367 **Figures**




368

369 **Fig. 1** 50 hPa zonal mean zonal wind during June to August derived from ERA5. The  
 370 resolution of the abscissa is the day, only the corresponding year is marked. The QBO  
 371 phases are labeled next to the years.



**Fig. 2** Summer mean zonal wind anomalies from climatology at 150 hPa during (a) EQBO and (b) WQBO. Zonal wind anomalies at 120°E during (c) EQBO and (d) WQBO. The climatological zonal wind is shown by purple contour lines. The regions with empty dots exceed the 95% confidence level under the Monte Carlo test. The range of the black box in (c) and (d) is 10°N-10°S, 150-200 hPa.

381




382

383

384 **Fig. 3** (a) Climatology of OLR and (b) difference in OLR between EQBO and WQBO.  
 385 (c) Climatology 200 hPa  $\omega$  and (d) difference in 200 hPa  $\omega$  between EQBO and  
 386 WQBO. (e) Climatology of divergence at 150 hPa (contours with values of 3 and 18  
 387  $10^{-6} \text{ s}^{-1}$ ) and difference in divergence between EQBO and WQBO (filled color). The  
 388 black lines mark 90°E and 140°E. (f) Differences in  $u$  and  $\omega$  along the equator  
 389 between EQBO and WQBO (vectors). Differences in  $\omega$  are statistically tested.  
 390 Climatology of  $\omega$  along the equator (filled color). The regions with empty dots exceed  
 391 the 95% confidence level under the Monte Carlo test.

392

393



394

395 **Fig. 4** Difference in  $\frac{\partial\theta}{\partial z}$  (filled color, unit:  $\text{K m}^{-1}$ ) and  $\omega$  (vectors, unit:  $10^{-3} \text{ Pa s}^{-1}$ )

396 between EQBO and WQBO at (a)  $90^\circ\text{E}$  and (b)  $140^\circ\text{E}$ . Differences in  $\frac{\partial\theta}{\partial z}$  above the

397 95% confidence level under the Monte Carlo test are shown by empty dots.

398

EQBO	zonal	Standardized	box region TEJ	WQBO	zonal	Standardized	box region TEJ
year	wind	Index		year	wind	Index	
1987	-24.1	-1.6	-10.6	1988	10.2	0.9	-13.1
1992	-23.9	-1.6	-11.9	1990	9.4	0.9	-16.5
1994	-24.3	-1.7	-14.6	1993	9.0	0.8	-10.6
1996	-17.6	-1.2	-9.3	1995	10.8	1.0	-9.7
1998	-20.5	-1.4	-8.3	1997	13.1	1.1	-13.8
2001	-14.1	-0.9	-14.3	1999	9.5	0.9	-15.5
2007	-9.7	-0.6	-10.0	2000	13.1	1.1	-12.7
2010	-23.0	-1.6	-9.6	2002	8.8	0.8	-15.0
2015	-10.0	-0.6	-11.0	2004	9.8	0.9	-16.6
2016	-14.3	-0.9	-13.7	2006	9.1	0.8	-13.3
2018	-21.3	-1.4	-16.4	2008	8.3	0.8	-10.0
2020	-13.9	-0.9	-7.4	2009	9.3	0.8	-15.6
				2011	11.7	1.0	-14.8
				2013	7.6	0.7	-13.8
				2014	7.5	0.7	-13.3
				2017	11.8	1.0	-13.8
				2019	8.6	0.8	-13.4

average	-18.1	-11.4	average	9.9	-13.6
---------	-------	-------	---------	-----	-------

401

402 **Table 1.** The summer mean zonal wind at 50 hPa (U50, unit: m/s), standardized U50  
 403 observed at the Singapore radiosonde station (1°N/104°E), and averaged TEJ zonal  
 404 wind (unit: m/s) in the black box region in Figure 2 of EQBO and WQBO years.

405

406